1. A laboratory technologist working in a hospital’s clinical chemistry laboratory adds accelerator during total bilirubin test procedure only, but not during direct bilirubin test procedure.

Why does NOT the technologist add accelerator during direct bilirubin test?

(A) The direct bilirubin reacts with bilirubin reagent rapidly

(B) The bilirubin reagent directly interacts with director solutions

(C) The direct bilirubin test kit contains an accelerator solution

(D) The yellow color icterus sample does not affect direct bilirubin test

The correct answer to the question is (A) The direct bilirubin reacts with bilirubin reagent rapidly. Here is a detailed explanation:

- Bilirubin is a yellow pigment that is produced when red blood cells break down. It is normally processed by the liver and excreted in bile.

- There are two types of bilirubin: **direct** and **indirect**. Direct bilirubin is also called **conjugated** bilirubin, because it is combined with a substance called glucuronic acid in the liver. Indirect bilirubin is also called **unconjugated** bilirubin, because it is not combined with anything.

- A **total bilirubin test** measures the amount of both direct and indirect bilirubin in the blood. A **direct bilirubin test** measures only the amount of direct bilirubin in the blood. An **indirect bilirubin test** is calculated by subtracting the direct bilirubin from the total bilirubin.

- To measure the total bilirubin, a reagent called **diazotized sulfanilic acid** is added to the blood sample. This reagent reacts with both direct and indirect bilirubin, forming a colored compound called **azo-bilirubin**. The intensity of the color is proportional to the amount of total bilirubin in the sample.

- To measure the direct bilirubin, the same reagent is used, but an **accelerator** is also added. The accelerator is a substance that enhances the reaction between the reagent and the direct bilirubin, but not the indirect bilirubin. This way, only the direct bilirubin forms azo-bilirubin, and the color intensity reflects the amount of direct bilirubin in the sample.

- The reason why an accelerator is not needed for the direct bilirubin test is that the direct bilirubin reacts with the reagent **rapidly** and **completely**, even without the accelerator. The indirect bilirubin, on the other hand, reacts with the reagent **slowly** and **incompletely**, even with

the accelerator. Therefore, the accelerator is only useful for the total bilirubin test, where it helps to measure both types of bilirubin. For the direct bilirubin test, the accelerator is unnecessary and may even interfere with the accuracy of the result.

2. One of the routine procedures performed by a laboratory technologist working in a clinical chemistry laboratory is preparation of working solutions. Accordingly, the technologist needs to prepare ALT (GPT) working solutions by mixing 4 part from reagent 1 (R1) and 1 part from reagent 2 (R2). He needs to prepare 1 ml of this working solution for the test.

What will be the proportions of R1 and R2 in the solutions respectively?

(A) 200 µl and 800 µl

(B) 400 µl and 600 µl

(C) 600 µl and 400 µl

(D) 800 µl and 200 µl
```

The correct answer to question number 2 is (D) 800 µl and 200 µl. Here is a detailed explanation:

- The question asks about the proportions of R1 and R2 in the working solution for ALT (GPT) test, which is prepared by mixing 4 parts of R1 and 1 part of R2.

- To find the proportions of R1 and R2 in 1 ml of the working solution, we need to divide 1 ml by the total number of parts, which is 4 + 1 = 5. This gives us the volume of one part, which is 1/5 ml or 200 µl.

- Then, we need to multiply the volume of one part by the number of parts of each reagent. For R1, we have 4 parts, so we multiply 200 µl by 4, which gives us 800 µl. For R2, we
```

have 1 part, so we multiply 200 µl by 1, which gives us 200 µl.

- Therefore, the proportions of R1 and R2 in the working solution are 800 µl and 200 µl respectively.

3. A laboratory technologist collected blood specimens from a 35-year-old female patient at the peripheral laboratory. The specimen was collected for glucose measurement and there is no instrument used for measuring glucose at that laboratory. Therefore, the specimen should be preserved and transported to the central laboratory.

What is the best preservative of the specimen for this case?

(A)EDTA (B)Citrate (C)Oxalate (D)Fluoride

```
```

The correct answer to question number 3 is (D) Fluoride. Here is a detailed explanation:

- The question asks about the best preservative for blood specimens collected for glucose measurement, when there is no instrument available at the peripheral laboratory and the specimen needs to be transported to the central laboratory.

- Glucose is a substance that can be metabolized by enzymes or microorganisms in the blood, leading to a decrease in its concentration over time. This can affect the accuracy of the glucose measurement and cause false low results.

- To prevent this, a preservative is added to the blood specimen to inhibit the glycolysis process and stabilize the glucose level. Different preservatives have different mechanisms and effects on glucose preservation.

- **EDTA** is an anticoagulant that binds to calcium ions and prevents blood clotting. It does not have a direct effect on glucose preservation, but it may indirectly reduce glycolysis by lowering the pH and inhibiting some enzymes. However, EDTA is not very effective and can only preserve glucose for up to 8 hours at room temperature[1].

- **Citrate** is another anticoagulant that chelates calcium ions and prevents clotting. It also does not have a direct effect on glucose preservation, but it may indirectly reduce glycolysis by lowering the pH and inhibiting some enzymes. However, citrate is also not very effective and can only preserve glucose for up to 8 hours at room temperature[1].

- **Oxalate** is a third anticoagulant that precipitates calcium ions and prevents clotting. It also does not have a direct effect on glucose preservation, but it may indirectly reduce glycolysis by lowering the pH and inhibiting some enzymes. However, oxalate is

also not very effective and can only preserve glucose for up to 8 hours at room temperature[1].

- **Fluoride** is the best preservative for glucose measurement, as it has a direct effect on inhibiting glycolysis. Fluoride acts as an enzyme inhibitor that blocks the activity of glycolytic enzymes, such as hexokinase and enolase, and stops the breakdown of glucose. Fluoride can preserve glucose for up to 24 hours at room temperature, and up to 48 hours at 4°C[2][3].

Therefore, fluoride is the best preservative for blood specimens collected for glucose measurement, when there is no instrument available at the peripheral laboratory and the specimen needs to be transported to the central laboratory.

4. A laboratory technologist received blood sample collected from a 16-year-old male

patient for glucose analysis using spectrophotometer method. The instrument was under maintenance and the whole blood sample was left at room temperature for more than two hours prior to analysis.

What would be the effect of delayed analysis on glucose concentration?

(A) Decrease glucose level due to glycolysis at room temperature

(B) Decrease glucose level due to lipogenesis at room temperature

(C) Decrease glucose level due to glycogenesis at room temperature

(D) Decrease glucose level due to oxidation reaction at room temperature

```
```

The correct answer is (A) Decrease glucose level due to glycolysis at room temperature. Here is a detailed explanation:

- Glycolysis is a metabolic pathway that occurs in the cytoplasm where glucose is converted into pyruvate, leading to a decrease in the concentration of glucose. If the blood sample is left at room temperature, enzymes responsible for glycolysis remain active, continuing this process and reducing the amount of measurable glucose.

- Lipogenesis is a metabolic pathway that occurs in the cytosol and endoplasmic reticulum where acetyl-CoA is converted into fatty acids, leading to an increase in the concentration of lipids. This process does not directly affect the glucose level, but it may indirectly affect it by consuming acetyl-CoA, which is also a product of glycolysis. However, lipogenesis is not a major factor in glucose preservation, as it is regulated by hormonal and nutritional factors.

- Glycogenesis is a metabolic pathway that occurs in the cytosol and liver where glucose is converted into glycogen, leading to a decrease in the concentration of glucose. This process is also regulated by hormonal and nutritional factors, and it is not a major factor in glucose preservation, as it is limited by the availability of glycogen storage.

- Oxidation reaction is a chemical reaction that involves the loss of electrons or the increase of oxidation state. This process does not directly affect the glucose level, but it may indirectly affect it by producing reactive oxygen species (ROS), which can damage the blood cells and the enzymes involved in glycolysis. However, oxidation reaction is not a major factor in glucose preservation, as it is counteracted by antioxidant mechanisms.

Therefore, the main factor that affects the glucose level in a blood sample left at room temperature is glycolysis, which decreases

the glucose level due to the breakdown of glucose by enzymes.

5. A laboratory technologist collected blood specimen for ALP measurement using clinical chemistry analyzer. The instrument printed the enzyme activity result as 0.2mmol/l. The laboratory technologist was interested to report the result in µmol/l.

What is the enzyme activity result in µmol/l?

(A) 20 µmol/l

(B) 200 µmol/l

(C) 2,000 µmol/l

(D) 20,000 µmo1/1.

```

The correct answer is (B) 200 µmol/l. Here is a detailed explanation:
```

- ALP stands for **alkaline phosphatase**, which is an enzyme that catalyzes the hydrolysis of phosphate esters at alkaline pH. ALP is found in various tissues, such as liver, bone, kidney, and intestine, and its activity can be measured in the blood as a marker of various diseases or conditions.

- The enzyme activity of ALP is expressed in units of **moles of substrate hydrolyzed per liter of blood per minute**. This unit is abbreviated as **mol/L/min** or **mmol/L/min**. The latter is more commonly used, as it represents a smaller and more convenient scale.

- However, sometimes the enzyme activity of ALP is reported in units of **micromoles of substrate hydrolyzed per liter of blood per minute**. This unit is abbreviated as **µmol/L/min** or **µmol/L**. The latter is used when the enzyme activity is measured at a fixed time interval, such as 1 minute.

- To convert the enzyme activity of ALP from mmol/L to µmol/L, we need to multiply by 1000, since there are 1000 micromoles in a millimole. For example, if the enzyme activity of ALP is 0.2 mmol/L, then it is equivalent to 0.2 x 1000 = 200 µmol/L. This is the same as multiplying by 10^3^, which is the conversion factor between the two units.

- Therefore, the enzyme activity result of 0.2 mmol/L is equal to 200 µmol/L. This is the answer to the question.

6. A laboratory technologist working in a clinical chemistry laboratory performed Creatinine measurement using spectrophotometric method. The reaction was stable and an absorbance of 0.2 was recorded from 0.1 mg/dL of standard creatinine solution. Another unknown sample was performed for creatinine measurement and an absorbance of 0.3 was recorded.

What was the concentration of creatinine in the unknown sample?

(A) 0.1 mg/dL

(B) 0.06 mg/dL

(C) 0.15 mg/dL

(D) 0.21 mg/dL

```

The correct answer is (C) 0.15 mg/dL. Here is a detailed explanation:

- Creatinine is a waste product of muscle metabolism that is excreted by the kidneys. Its concentration in the blood can be used to assess kidney function and diagnose kidney diseases.

- To measure the creatinine concentration, a spectrophotometric method is used, which involves a chemical reaction that produces a colored product that absorbs light at a
```

specific wavelength. The amount of light absorbed by the product is proportional to the amount of creatinine in the sample, according to the Beer-Lambert law: $A = \epsilon l c$, where A is the absorbance, ϵ is the molar absorptivity, l is the path length, and c is the concentration.

- To calibrate the spectrophotometer, a standard solution of known creatinine concentration is used. The absorbance of the standard solution is measured and used as a reference point. Then, the absorbance of the unknown sample is measured and compared to the absorbance of the standard solution, using the following formula: $c_s = \frac{A_s}{A_u} c_u$, where c_s is the concentration of the standard solution, A_s is the absorbance of the standard solution, c_u is the concentration of the unknown sample, and A_u is the absorbance of the unknown sample.

- In this question, the standard solution has a concentration of 0.1 mg/dL and an

absorbance of 0.2. The unknown sample has an absorbance of 0.3. To find the concentration of the unknown sample, we plug in the values into the formula: $c_u = \frac{A_u}{A_s} c_s = \frac{0.3}{0.2} \times 0.1 = 0.15 \text{ mg/dL}$.

- Therefore, the concentration of creatinine in the unknown sample is 0.15 mg/dL. This is the answer to the question.

7. A laboratory technologist working in a clinical chemistry laboratory collected blood specimens from a 78-year-old female patient for glucose measurement using glucose oxidase test method. The technologist used an accelerator in the procedure to shorten the reaction time.

What was the role of the accelerator?

(A) Accelerate the reaction by converting α- L glucose to β- L glucose

(B) Accelerate the reaction by converting β- L glucose to α- L glucose

(C) Accelerate the reaction by converting β- D glucose to α- D glucose

(D) Accelerate the reaction by converting α- D glucose to β- D glucose
```

The correct answer is (D) Accelerate the reaction by converting α- D glucose to β- D glucose. Here is a detailed explanation:

- Glucose oxidase is an enzyme that catalyzes the oxidation of β- D glucose to D-gluconolactone and hydrogen peroxide. This reaction is used to measure the concentration of glucose in blood samples, as the amount of hydrogen peroxide produced is proportional to the amount of glucose present.

- However, glucose exists in two anomeric forms: α- D glucose and β- D glucose. These
```

forms differ in the orientation of the hydroxyl group on the first carbon atom. Glucose oxidase is specific for β- D glucose and does not react with α- D glucose. Therefore, the glucose concentration measured by glucose oxidase may not reflect the true glucose concentration, as some glucose may be in the α- D form.

- To overcome this problem, an accelerator is added to the reaction mixture. The accelerator is a substance that converts α- D glucose to β- D glucose, making it available for the glucose oxidase reaction. This way, the reaction rate and the accuracy of the glucose measurement are increased. One example of an accelerator is mutarotase, an enzyme that catalyzes the interconversion of α- D glucose and β- D glucose.

- Therefore, the role of the accelerator is to accelerate the reaction by converting α- D glucose to β- D glucose. This is the answer to the question.

8. A laboratory technologist working in a clinical chemistry laboratory collected blood specimens from a 78-year-old female patient for glucose measurement using glucose oxidase test method. The technologist used an accelerator in the procedure to shorten the reaction time.

What was the role of the accelerator?

(A) Accelerate the reaction by converting α- L glucose to β- L glucose

(B) Accelerate the reaction by converting β- L glucose to α- L glucose

(C) Accelerate the reaction by converting β- D glucose to α- D glucose

(D) Accelerate the reaction by converting α- D glucose to β- D glucose

```
```

The correct answer is (D) Accelerate the reaction by converting α- D glucose to β- D glucose. Here is a detailed explanation:

- Glucose oxidase is an enzyme that catalyzes the oxidation of β- D glucose to gluconic acid and hydrogen peroxide. This reaction is used to measure the concentration of glucose in blood samples, as the amount of hydrogen peroxide produced is proportional to the amount of glucose present.

- However, glucose exists in two anomeric forms: α- D glucose and β- D glucose. These forms differ in the orientation of the hydroxyl group on the first carbon atom. Glucose oxidase is specific for β- D glucose and does not react with α- D glucose. Therefore, the glucose concentration measured by glucose oxidase may not reflect the true glucose concentration, as some glucose may be in the α- D form.

- To overcome this problem, an accelerator is added to the reaction mixture. The accelerator is a substance that converts α- D glucose to β- D glucose, making it available for the glucose oxidase reaction. This way, the reaction rate and the accuracy of the glucose measurement are increased. One example of an accelerator is mutarotase, an enzyme that catalyzes the interconversion of α- D glucose and β- D glucose.

- According to the web search results, Biosystems reagents use the mutarotase method for direct determination of LDL-C[12]. Therefore, the role of the accelerator is to accelerate the reaction by converting α- D glucose to β- D glucose. This is the answer to the question.

9. A 25-year-old man with confirmed hyperlipidemia brought a request paper to the central laboratory to check his blood LDL cholesterol (LDL-C) level. The technologist collected blood specimen for the analysis and

measured LDL-C level directly using Biosystems reagents.

What method was used by the technologist?

(A) By an empirical Friedewald method

(B) By ultracentrifugation to mask other cholesterols

(C) By precipitation of other cholesterol using polyanion

(D) By masking other cholesterols using polyclonal antibody

```

The correct answer is (D) By masking other cholesterols using polyclonal antibody. Here is a detailed explanation:

- LDL-C is a type of lipoprotein that carries cholesterol in the blood. It is often called "bad" cholesterol because high levels of LDL-
```

C can increase the risk of cardiovascular diseases, such as atherosclerosis and coronary artery disease.

- To measure the LDL-C level, there are two main methods: **indirect** and **direct**. The indirect method estimates the LDL-C level by using the Friedewald formula, which subtracts the HDL-C (high-density lipoprotein, or "good" cholesterol) and VLDL-C (very-low-density lipoprotein) from the total cholesterol. The direct method measures the LDL-C level by using a specific reagent that reacts only with LDL-C and not with other lipoproteins.

- Biosystems reagents use the direct method for LDL-C measurement. The reagent contains a polyclonal antibody that binds to the apolipoprotein B-100, which is a protein component of LDL-C. This antibody masks the other cholesterols, such as HDL-C and VLDL-C, and prevents them from interfering with the reaction. The reagent also contains a chromogen that produces a color change

when it reacts with the cholesterol in LDL-C. The intensity of the color is proportional to the amount of LDL-C in the sample.

- Therefore, the method used by the technologist is by masking other cholesterols using polyclonal antibody. This is the answer to the question.

10. A physician physically diagnosed a 45-year-old male patient with azotemia and send blood sample to the laboratory assessment. The laboratory technologist analyzed the specimen for renal function tests. The technologist recorded 90 mg/dl urea and 1.20 mg/dl creatinine.

Calculate blood urea nitrogen to creatinine ratio?

(A) 13.50

(B) 35.04

(C) 70.46

(D) 75.00
```

The correct answer is (B) 35.04. Here is a detailed explanation:

- Azotemia is a condition where there is an abnormally high level of nitrogenous waste products in the blood, such as urea and creatinine. It can indicate impaired kidney function or dehydration.

- Renal function tests are used to assess the kidney function and diagnose kidney diseases. They include measuring the levels of urea, creatinine, and other substances in the blood and urine.

- Blood urea nitrogen (BUN) is a measure of the amount of urea in the blood. Urea is a waste product of protein metabolism that is excreted by the kidneys. BUN can reflect the balance between the production and
```

excretion of urea. High BUN can indicate kidney dysfunction, dehydration, high protein intake, or liver disease. Low BUN can indicate malnutrition, liver failure, or overhydration.

- Creatinine is a measure of the amount of creatinine in the blood. Creatinine is a waste product of muscle metabolism that is excreted by the kidneys. Creatinine can reflect the glomerular filtration rate (GFR), which is the rate at which the kidneys filter the blood. High creatinine can indicate kidney dysfunction, muscle damage, or dehydration. Low creatinine can indicate muscle wasting, malnutrition, or overhydration.

- BUN to creatinine ratio (BUN/Cr) is a ratio that compares the levels of BUN and creatinine in the blood. It can help to differentiate the causes of azotemia and evaluate the kidney function. Normal BUN/Cr ratio ranges from 10:1 to 20:1. High BUN/Cr ratio (>20:1) can indicate prerenal azotemia,

which is caused by reduced blood flow to the kidneys due to dehydration, heart failure, shock, or bleeding. Low BUN/Cr ratio (<10:1) can indicate intrarenal azotemia, which is caused by damage to the kidney tissues due to infection, inflammation, or toxins. Normal BUN/Cr ratio can indicate postrenal azotemia, which is caused by obstruction of the urine flow due to kidney stones, enlarged prostate, or tumors.

- To calculate the BUN/Cr ratio, we need to convert the urea concentration to BUN using the formula: BUN = Urea / 2.14. This is because urea contains two nitrogen atoms, while BUN measures only one nitrogen atom. Then, we divide the BUN by the creatinine to get the ratio. For example, if the urea is 90 mg/dL and the creatinine is 1.20 mg/dL, then the BUN/Cr ratio is:

BUN = Urea / 2.14 = 90 / 2.14 = ~42 mg/dL

BUN/Cr = BUN / Creatinine = ~42 / 1.20 = ~35

Therefore, the BUN/Cr ratio is ~35, which is closest to option (B) 35.04. This is the answer to the question.

11. A 41-year-old male patient with a history of tachycardia, hypertension, and diabetic mellitus was admitted to a hospital. Different chemistry tests and urinalysis were requested for him by the physician. The laboratory technologist analyzed the specimens and reported the following result: high AST, high LDH, high CK-MB isoenzyme, high urea, high creatinine in serum and few WBC and RBC in urine.

What is the most likely condition that could cause the above alterations?

(A) Renal acute kidney injury

(B) Intrinsic acute kidney injury

(C) Pre-renal acute kidney injury

(D) Post-renal acute kidney injury

```

The correct answer is (C) Pre-renal acute kidney injury. Here is a detailed explanation:

- Acute kidney injury (AKI) is a sudden and temporary loss of kidney function that can be caused by various factors affecting the blood flow, the kidney tissue, or the urine output[1][2].

- Pre-renal AKI is caused by a decrease in blood flow to the kidneys, which impairs their ability to filter waste products and maintain fluid and electrolyte balance. Pre-renal AKI can be triggered by conditions such as dehydration, heart failure, shock, blood loss, or sepsis[1][2].

- Intrinsic AKI is caused by damage to the kidney tissue itself, which affects the structures and functions of the nephrons, the
```

basic units of the kidney. Intrinsic AKI can be caused by conditions such as glomerulonephritis, acute tubular necrosis, interstitial nephritis, or vascular disorders[1][2].

- Post-renal AKI is caused by an obstruction in the urinary tract, which prevents the urine from leaving the kidneys and causes a backflow of pressure and toxins. Post-renal AKI can be caused by conditions such as kidney stones, enlarged prostate, tumors, or blood clots[1][2].

- The laboratory results of the patient suggest that he has pre-renal AKI, because:

    - High AST, LDH, and CK-MB indicate muscle damage and heart injury, which can reduce the blood flow to the kidneys[3][4].

    - High urea and creatinine indicate impaired kidney function and reduced glomerular filtration rate, which can result from low blood pressure or volume[5] .

    - Few WBC and RBC in urine indicate no significant inflammation or bleeding in the

kidney tissue, which would be expected in intrinsic or post-renal AKI .

Therefore, the most likely condition that could cause the above alterations is pre-renal acute kidney injury. This is the answer to the question.

12. A 46-year-old woman with thyrotoxicosis, goiter, ophthalmopathy and dermopathy visited the laboratory for thyroid function tests. The laboratory technologist collected, processed and analyzed the serum sample using chemiluminescence principle. Her laboratory finding showed high free T4 and T3, and low TSH. The thyroid stimulating immunoglobulins (TSIs) and TSH receptor antibodies were also positive.

What disease causes the above alterations?

(A) Graves' disease

(B) Postpartum thyroiditis

(C) Hashimoto's thyroiditis

(D) Iodine deficiency disease
```

The correct answer is (A) Graves' disease. Here is a detailed explanation:

- Thyrotoxicosis is a condition where there is an excess of thyroid hormones in the body, causing symptoms such as increased heart rate, weight loss, nervousness, and heat intolerance. Goiter is an enlargement of the thyroid gland, which may cause difficulty in swallowing or breathing. Ophthalmopathy is an eye disorder that causes bulging eyes, dryness, irritation, and vision problems. Dermopathy is a skin disorder that causes redness and swelling, usually on the shins or feet.
```

- Thyroid function tests are blood tests that measure the levels of thyroid hormones (T4 and T3) and thyroid stimulating hormone (TSH) in the body. TSH is a hormone produced by the pituitary gland that regulates the thyroid gland. T4 and T3 are hormones produced by the thyroid gland that regulate the metabolism and other functions of the body. TSIs and TSH receptor antibodies are antibodies that bind to and stimulate the TSH receptors on the thyroid gland, causing it to produce more thyroid hormones.

- Graves' disease is an autoimmune disorder that causes the body to produce TSIs and TSH receptor antibodies, which mimic the action of TSH and stimulate the thyroid gland to produce excessive amounts of thyroid hormones. This leads to thyrotoxicosis, goiter, ophthalmopathy, and dermopathy. Graves' disease is the most common cause of hyperthyroidism, which is a condition where the thyroid gland is overactive.

- Postpartum thyroiditis is a temporary inflammation of the thyroid gland that occurs after pregnancy. It can cause fluctuations in thyroid hormone levels, leading to hyperthyroidism or hypothyroidism, which is a condition where the thyroid gland is underactive. Postpartum thyroiditis usually resolves within a year and does not cause goiter, ophthalmopathy, or dermopathy.

- Hashimoto's thyroiditis is an autoimmune disorder that causes the body to produce antibodies that attack and damage the thyroid gland, causing it to produce less thyroid hormones. This leads to hypothyroidism, which can cause symptoms such as fatigue, weight gain, depression, and cold intolerance. Hashimoto's thyroiditis can also cause goiter, but not ophthalmopathy or dermopathy.

- Iodine deficiency disease is a condition where there is not enough iodine in the diet, which is essential for the synthesis of thyroid hormones. This leads to hypothyroidism and

goiter, as the thyroid gland tries to compensate for the lack of iodine by enlarging. Iodine deficiency disease does not cause ophthalmopathy or dermopathy.

Therefore, the disease that causes the above alterations is Graves' disease. This is the answer to the question.

13. A 26-year-old woman with a history of breast mass, breast pain, and skin discoloration was referred to a regional laboratory for CA 15-3 tumor marker test. Then, the laboratory technologist collected and analyzed blood specimen using chemiluminescence immunoassay. Her laboratory result showed low level of CA 15-3 in serum.

What condition is associated with the above result?

(A) Malignant tumor

(B) Benign breast tumor

(C) Free from breast cancer

(D) Last stage breast cancer

```

The correct answer is (C) Free from breast cancer. Here is a detailed explanation:

- CA 15-3 is a type of protein that is produced by some breast cancer cells and released into the blood. It is used as a tumor marker, which is a substance that can indicate the presence or progression of cancer. CA 15-3 is not specific for breast cancer, as it can also be elevated in other conditions, such as liver disease, ovarian cancer, or pregnancy .

- Chemiluminescence immunoassay is a method that uses antibodies and a chemical reaction that produces light to detect and
```

measure the amount of CA 15-3 in the blood. The intensity of the light is proportional to the amount of CA 15-3 in the sample.

- The normal range of CA 15-3 in the blood is less than 30 U/mL. A low level of CA 15-3 in the serum indicates that there is no or minimal breast cancer activity in the body. A high level of CA 15-3 in the serum indicates that there is a large or advanced breast cancer that is spreading or growing .

- In this question, the woman has a low level of CA 15-3 in the serum, which suggests that she does not have breast cancer or has a very early stage of breast cancer that is not detectable by this test. This is consistent with her symptoms of breast mass, breast pain, and skin discoloration, which can also be caused by benign breast conditions, such as fibrocystic breast disease, mastitis, or eczema .

- Therefore, the condition that is associated with the above result is free from breast cancer. This is the answer to the question.

14. A group of students conducted a project on plasma proteins level assessment among different age groups. They used Kjeldahl method for total protein determination. What element was determined by this method?

(A) Carbon

(B) Hydrogen

(C) Nitrogen

(D) Sulfur

```

The correct answer is (C) Nitrogen. Here is a detailed explanation:

- Kjeldahl method is a chemical method that is used to measure the amount of nitrogen in organic compounds, such as proteins. It is based on the principle that nitrogen in organic compounds can be converted to
```

ammonium sulfate by digestion with sulfuric acid and a catalyst. The ammonium sulfate can then be converted to ammonia gas by adding sodium hydroxide. The ammonia gas can be collected and titrated with a standard acid solution to determine the amount of nitrogen in the original sample.

- Proteins are organic compounds that consist of amino acids, which are molecules that contain nitrogen, carbon, hydrogen, oxygen, and sometimes sulfur. The nitrogen content of proteins varies depending on the type and number of amino acids, but it is usually around 16%. Therefore, by measuring the nitrogen content of a protein sample using the Kjeldahl method, the total protein content can be estimated by multiplying the nitrogen content by a factor of 6.25.

- Therefore, the element that is determined by the Kjeldahl method is nitrogen. This is the answer to the question.

15. A laboratory technologist working in a hospital clinical chemistry laboratory

performed serum protein electrophoresis. The result showed normal serum electrophoresis fraction values with similar patterns.

Which serum protein fraction was observed in highest concentrations?

(A) α1-Globulin

(B) α2-Globulin

(C) β-Globulin

(D) γ-Globulin

```

The correct answer is (A) α1-Globulin. Here is a detailed explanation:

- Serum protein electrophoresis is a test that separates the proteins in the blood serum into different fractions based on their size, shape, and charge. The fractions are
```

albumin, α1-globulin, α2-globulin, β-globulin, and γ-globulin. Each fraction contains different types of proteins that have different functions in the body.

- Albumin is the most abundant protein in the serum, accounting for about 60% of the total protein. It helps maintain the fluid balance and transport various substances in the blood. α1-Globulin is the second most abundant protein fraction, accounting for about 15% of the total protein. It contains proteins such as alpha-1 antitrypsin, which protects the lungs from damage by enzymes, and haptoglobin, which binds to free hemoglobin and prevents its loss in the urine. α2-Globulin accounts for about 10% of the total protein and contains proteins such as ceruloplasmin, which carries copper in the blood, and alpha-2 macroglobulin, which inhibits various proteases. β-Globulin accounts for about 8% of the total protein and contains proteins such as transferrin, which transports iron in the blood, and

complement proteins, which are involved in the immune system. γ-Globulin accounts for about 7% of the total protein and contains immunoglobulins, which are antibodies that fight infections.

- Therefore, the serum protein fraction that is observed in the highest concentrations is α1-globulin, followed by albumin, α2-globulin, β-globulin, and γ-globulin. This is the answer to the question.

16. A laboratory technologist working in a hospital clinical chemistry laboratory performed Alanine transaminase (ALT) test based on the principle of the following test reaction:

Alanine + α-Ketoglutarate ↔ ALT ↔ Pyruvate + glutamate

Pyruvate + NADH+H ↔ LD ↔ Lactate + NAD+

What does the technologist understand from the above reaction?

(A) LD is incorporated in the reagent test kit

(B) ALT is incorporated in the reagent test kit

(C) Pyruvate is incorporated in the reagent test kit

(D) The amino group from α-Ketoglutarate is transferred to pyruvate

```

The correct answer is (D) The amino group from α-Ketoglutarate is transferred to pyruvate. Here is a detailed explanation:

- Alanine transaminase (ALT) is an enzyme that is mainly found in the liver and catalyzes the reversible transfer of an amino group from alanine to α-ketoglutarate, producing pyruvate and glutamate. This reaction is part of the amino acid metabolism and the
```

gluconeogenesis pathway, which converts amino acids and other substrates into glucose[1][2].

- The above reaction shows the two steps involved in the ALT test. The first step is the ALT reaction, where the enzyme ALT converts alanine and α-ketoglutarate into pyruvate and glutamate. The second step is the lactate dehydrogenase (LD) reaction, where the enzyme LD converts pyruvate and NADH+H into lactate and NAD+. The LD reaction is coupled with the ALT reaction to regenerate NAD+, which is required for the ALT reaction to proceed. The LD reaction also produces a color change that can be measured by a spectrophotometer to determine the amount of ALT activity in the sample[3][4].

- From the above reaction, the technologist can understand that the amino group from α-ketoglutarate is transferred to pyruvate by the enzyme ALT. This is the main function of ALT and the basis of the ALT test. The

technologist can also understand that LD, NADH+H, and NAD+ are part of the reagent test kit, as they are needed for the LD reaction that is coupled with the ALT reaction. Pyruvate and glutamate are not part of the reagent test kit, as they are the products of the ALT reaction. Therefore, option (D) is the correct answer to the question.

17. A laboratory technologist working in a hospital clinical chemistry laboratory collected blood sample from a 35-year-old male patient for measurement of serum alkaline phosphatase (ALP) activity. The clinician indicated obstruction of patient's bile duct in the request paper. The laboratory analysis also showed that the serum activity of ALP was increased.

What is the next most appropriate confirmatory test of bile duct obstruction?

(A) Unconjugated bilirubin

(B) Acid phosphatase (ACP)

(C) Alanine transaminase (ALT)

(D) γ-Glutamyltransferase (GGT)

```

The correct answer is (D) γ-Glutamyltransferase (GGT). Here is a detailed explanation:

- Alkaline phosphatase (ALP) is an enzyme that is found in various tissues of the body, such as the liver, bone, intestine, and placenta. It is involved in the metabolism of phosphate and organic compounds. ALP can be measured in the blood serum to assess the function of these tissues and diagnose various diseases[1][2].

- Bile duct obstruction is a condition where the ducts that carry bile from the liver and
```

gallbladder to the intestine are blocked by gallstones, tumors, inflammation, or injury. This causes bile to accumulate in the liver and interfere with its function. Bile duct obstruction can lead to jaundice, which is a yellowing of the skin and eyes due to the buildup of bilirubin, a waste product of red blood cell breakdown[34].

- Serum ALP activity can be increased in bile duct obstruction, as the liver cells release more ALP into the bloodstream due to the damage and inflammation caused by the bile backup. However, serum ALP activity can also be increased in other conditions that affect the liver, bone, or other tissues, such as liver cirrhosis, bone fractures, or pregnancy. Therefore, serum ALP activity alone is not a specific indicator of bile duct obstruction[12].

- γ-Glutamyltransferase (GGT) is another enzyme that is found mainly in the liver and biliary system. It is involved in the transport and metabolism of amino acids and

glutathione, an antioxidant. GGT can also be measured in the blood serum to assess the function of the liver and biliary system and diagnose various diseases .

- Serum GGT activity can be increased in bile duct obstruction, as the biliary cells release more GGT into the bloodstream due to the damage and inflammation caused by the bile backup. Unlike ALP, serum GGT activity is more specific for bile duct obstruction, as it is not affected by bone or other tissue disorders. Therefore, serum GGT activity is a more reliable confirmatory test of bile duct obstruction .

Therefore, the next most appropriate confirmatory test of bile duct obstruction is γ-Glutamyltransferase (GGT). This is the answer to the question.

19. A laboratory technologist working in a hospital clinical chemistry laboratory received CSF sample of a 33-year-old female patient for potassium electrolyte test. The technologist measured CSF potassium level and reported the result without including the reference interval.

What is the estimation value of CSF potassium level compared to plasma potassium level?

(A) CSF and plasma potassium values are the same

(B) CSF potassium is 70% of plasma potassium

(C) CSF potassium is 20% of plasma potassium

(D) CSF potassium is 50% of plasma potassium

The correct answer is (B).

Explanation:

- CSF is the fluid that surrounds the brain and spinal cord, and it has a similar composition to plasma, but with lower protein and glucose levels[1].

- Potassium is an electrolyte that is important for nerve and muscle function, and it is regulated by the kidneys[2].

- The normal range of potassium in plasma is about 3.5 to 5.0 mmol/L, while the normal range of potassium in CSF is about 2.5 to 3.5 mmol/L[3][4].

- Therefore, CSF potassium is about 70% of plasma potassium, or 0.7 times the plasma potassium level.

- For example, if the plasma potassium level is 4.0 mmol/L, the CSF potassium level would be about 2.8 mmol/L (0.7 x 4.0).

- This ratio is relatively constant and independent of the absolute values of potassium in plasma and CSF[3].

- The other choices are incorrect because they do not reflect the correct proportion of CSF to plasma potassium.

21. A hospital bought a new Clinical Chemistry machine. During calibration of the machine, the laboratory technologist noticed that some analytes are measured after mixing the sample with appropriate reagents and

then incubated for few minutes. The final product will be stable only for few minutes and finally will be deteriorated. He then calibrated the machine to read absorbance after incubation exactly at the beginning of product formation and at the start of deterioration times.}$$

What was the type of method described in the above scenario?

(A) End point method

(B) Fixed kinetics method

(C) Molar absorptivity method

(D) Continuous monitoring method

The correct answer is (B) Fixed kinetics method.

Explanation:

- A **fixed kinetics method** is a type of kinetic method that measures the change in absorbance or other signal at fixed time intervals during a chemical reaction[1].

- This method is useful for reactions that have a non-linear relationship between the signal and the concentration of the analyte or the product[1].

- In this scenario, the technologist calibrated the machine to read the absorbance at two fixed time points: one at the beginning of product formation and one at the start of product deterioration. This ensures that the absorbance readings are proportional to the product concentration and not affected by the instability of the product[2].

- A **end point method** is a type of kinetic method that measures the signal only at the end of the reaction, when the reaction reaches equilibrium or completion[1]. This method is not suitable for this scenario because the product is unstable and deteriorates quickly, so the end point signal would not reflect the true product concentration.

- A **molar absorptivity method** is not a type of kinetic method, but a property of a substance that relates its absorbance to its concentration and path length[3]. This method is not relevant to this scenario because it does not involve measuring the rate of a reaction.

- A **continuous monitoring method** is a type of kinetic method that measures the signal continuously throughout the reaction, without any fixed time intervals[1]. This method is not suitable for this scenario because it would require a more complex and

expensive instrument that can record the signal changes in real time.

22. A 55-year-old male patient brought a request paper for serum thyroid function tests from endocrinology department. A laboratory technologist made appropriate preparation when to collect blood sample, and how to interpret the results.

Which one is correct about TSH?

(A) TSH concentration is higher in men than women

(B) TSH concentration is the same in men and women

(C) TSH concentration is very low between 2:00 and 4:00 AM

(D) TSH is stable for a month when stored at 2 to 8°C.

The correct answer is (C) TSH concentration is very low between 2:00 and 4:00 AM.

Explanation:

- TSH is a hormone produced by the pituitary gland that stimulates the thyroid gland to produce thyroid hormones T3 and T4[1].

- TSH has a **circadian rhythm**, meaning it varies throughout the day and night according to a regular pattern[23].

- TSH levels are usually **highest** around **midnight** and **lowest** between **2:00 and 4:00 AM**[23]. This is because TSH

secretion is influenced by light exposure, melatonin, and other factors[23].

- The circadian rhythm of TSH is important for maintaining normal thyroid function and metabolism[23].

- The other choices are incorrect because:

  - (A) TSH concentration is **not** higher in men than women. TSH levels are similar in both sexes, although they may vary slightly with age[4].

  - (B) TSH concentration is **not** the same in men and women. TSH levels are similar, but not identical, in both sexes. There may be some differences due to factors such as pregnancy, menopause, or thyroid disease[4].

  - (D) TSH is **not** stable for a month when stored at 2 to 8°C. TSH is a protein hormone that can degrade over time, especially at higher temperatures. According to one study, TSH is stable for only 7 days when stored at 2 to 8°C[5].

9. A 72-year-old male patient brought a request paper for lipid profile tests to the laboratory. The laboratory technologist collected blood specimen to perform the above tests. The results of the lipid profile tests showed high total cholesterol, low HDL cholesterol and high LDL cholesterol.

What is the most likely underlying disease of the patient?

(A) Renal disease

(B) Pulmonary disease

(C) Cardiovascular disease

(D) Gastrointestinal disease

```

The correct answer is **(C) Cardiovascular disease**.
```

- High total cholesterol, low HDL (high-density lipoprotein), and high LDL (low-density lipoprotein) are classic indicators of an increased risk for cardiovascular diseases. HDL is often referred to as "good" cholesterol because it helps remove other forms of cholesterol from your bloodstream, hence higher levels are better. On the other hand, LDL is sometimes called "bad" cholesterol because high levels can lead to plaque buildup in arteries and result in heart diseases and stroke.

- Renal disease (A) can affect the levels of cholesterol and triglycerides in the blood, but it is not the most likely cause of the patient's lipid profile. Renal disease would also cause other symptoms and signs such as proteinuria, hematuria, edema, and hypertension.

- Pulmonary disease (B) is not directly related to the lipid profile of the patient. Pulmonary disease would affect the respiratory system

and cause symptoms such as dyspnea, cough, wheeze, and cyanosis.

- Gastrointestinal disease (D) can also affect the lipid profile of the patient, but it is not the most likely cause of the patient's lipid profile. Gastrointestinal disease would also cause other symptoms and signs such as abdominal pain, nausea, vomiting, diarrhea, and weight loss.

Here is the answer and explanation for the question:

```text
18. A 45-year-old male cardiac patient brought a request paper for the diagnosis of creatinine kinase (CK) isoenzymes profile as part of cardiac function test. The technologist reported normal electrophoresis result of CK iso-enzymes as shown below.
```

Cathode O O O O O Anode

A B C

Which letter properly matches with the isolated fractions of CK?

(A) Symbol "B" represent CK-MM

(B) Symbol "A" represent CK-MB

(C) Symbol "C" represent CK-BB

(D) Symbol "A" represent CK-BB

```

The correct answer is **(A) Symbol "B" represent CK-MM**.

- Creatine kinase (CK) is an enzyme found in various tissues and cell types, especially in muscles. It catalyzes the conversion of creatine and ATP to phosphocreatine and
```

ADP, and vice versa. This helps to maintain the energy balance in the cells. [1]

- CK has three different isoenzymes, which are CK-MM, CK-MB, and CK-BB. They differ in their subunit composition and tissue distribution. CK-MM is mainly found in skeletal muscle and heart, CK-MB is mainly found in heart, and CK-BB is mainly found in brain and smooth muscle. [12]

- Electrophoresis is a technique that separates molecules based on their size, shape, and charge. In the case of CK isoenzymes, they are separated by their charge and migrate towards the opposite poles of an electric field. CK-MM has the highest negative charge and moves towards the anode (positive pole), CK-BB has the lowest negative charge and moves towards the cathode (negative pole), and CK-MB has an intermediate charge and stays in the middle. [34]

- Therefore, in the normal electrophoresis result of CK isoenzymes, symbol A represents

CK-MB, symbol B represents CK-MM, and symbol C represents CK-BB. [34]

Source: Conversation with Bing, 1/2/2024

(1) Creatine Kinase with Isoenzymes (Blood) - Health Encyclopedia .... https://www.urmc.rochester.edu/encyclopedia/content.aspx?ContentTypeID=167&ContentID=creatine_kinase_isoenzyme_serum.

(2) Creatine kinase - Wikipedia. https://en.wikipedia.org/wiki/Creatine_kinase.

(3) Creatine Kinase Isoenzymes (CK Isoenzymes) with Total CK. https://testdirectory.questdiagnostics.com/test/test-detail/4451/creatine-kinase-isoenzymes-ck-isoenzymes-with-total-ck?cc=MASTER.

(4) San Diego Hospital, Healthcare - Health Library. https://myhealth.ucsd.edu/Library/TestsProcedures/167,creatine_kinase_isoenzyme_serum.

Here's the question with its choices, presented in a code box for clarity:

Question 24:

A laboratory technologist received CSF specimen of a 55-year-old female patient for chemistry, serology, hematology, and microbiology tests. The technologist was not able to analyze the sample immediately and needed to preserve it. The technologist divided the CSF specimen into three independent tubes to preserve separately.

How should the technologist preserve the specimen for hematology, chemistry and serology, and microbiology tests respectively?

(A) Room Temperature, Refrigeration, Frozen

(B) Frozen, Room Temperature, Refrigeration

(C) Refrigeration, Room Temperature, Frozen

(D) Refrigeration, Frozen, Room Temperature

Explanation:

The correct answer is (C) Refrigeration, Room Temperature, Frozen.

Here's a breakdown of the appropriate preservation methods for each test:

1. **Hematology:**

- Refrigeration at 2-8°C is generally recommended for hematological testing of CSF. This helps maintain cell integrity and prevent degradation of cellular components.

2. **Chemistry and Serology:**

- Most chemistry and serology tests on CSF can be performed on specimens stored at room temperature (18-25°C) for up to 24 hours. However, some specific tests might require refrigeration or freezing, so it's best to consult laboratory guidelines.

3. **Microbiology:**

- Freezing at -20°C or lower is essential for preserving CSF specimens for microbiological testing, especially for culture and identification of bacteria, viruses, or fungi. Freezing prevents growth and overgrowth of microorganisms, ensuring accurate results.

Key points:

- CSF is a delicate specimen that requires careful handling and preservation.
- Different tests have specific preservation requirements to ensure accurate results.
- Following laboratory guidelines is crucial for proper specimen management.
- Refrigeration is suitable for hematology testing to maintain cell integrity.
- Room temperature is often acceptable for chemistry and serology tests for a limited time.
- Freezing is necessary for microbiology testing to prevent microbial growth and overgrowth.

Sure, here is the full question in a code block:

```

24. A laboratory technologist received CSF specimen of a 55-year-old female patient for chemistry, serology, hematology and microbiology tests. The technologist was not able to analyze the sample immediately and he should preserve the sample. Then, the technologist divided the CSF specimen into three independent tubes to preserve separately.

How should the technologist preserve the specimen for hematology, chemistry and serology and microbiology tests respectively?

(A) Room Temperature , Refrigeration, Frozen

(B) Frozen, Room Temperature ,Refrigeration

(C) Refrigeration ,Room Temperature ,Frozen

(D) Refrigeration,Frozen ,Room Temperature

```

Question 24 is:

"How should the technologist preserve the specimen for hematology, chemistry and serology and microbiology tests respectively?

(A) Room Temperature , Refrigeration, Frozen

(B) Frozen, Room Temperature ,Refrigeration

(C) Refrigeration ,Room Temperature ,Frozen

(D) Refrigeration,Frozen ,Room Temperature"

The question is about the preservation of a Cerebrospinal Fluid (CSF) specimen for various tests. The correct preservation method depends on the type of test:

1. **Hematology**: The CSF sample for hematology should be kept at room temperature[2].

2. **Chemistry and Serology**: The CSF sample for chemistry and serology tests should be refrigerated[2][3].

3. **Microbiology**: The CSF sample for microbiology tests should not be refrigerated. Instead, it should be kept at a temperature of 35°C to 37°C[3].

Therefore, the correct answer is **(D) Refrigeration, Frozen, Room Temperature**[2][3]. This ensures that the samples are preserved in a way that maintains their integrity for the respective tests. Please note that these are general guidelines and specific tests may require different preservation methods. Always follow the instructions provided by the laboratory or test kit manufacturer.

Source: Conversation with Bing, 1/3/2024

(1) CSF Specimen Collection for Analysis - UI Health Care.

https://www.healthcare.uiowa.edu/path_handbook/Appendix/TechServ/COLLECTION_OF_CSF.html.

(2) Unit 4 Cerebrospinal fluid (CSF) - Open University of Sri Lanka. https://www.ou.ac.lk/apps/downloads/downloadfile/CSF%20I%20%283%20files%20merged%29.pdf.

(3) Cerebrospinal Fluid (CSF) Analysis - procedure, test, blood, tube .... https://www.surgeryencyclopedia.com/Ce-Fi/Cerebrospinal-Fluid-CSF-Analysis.html.

(4) Collection and storage of CSF samples for CSF 14-3-3/RT-QuIC analysis. https://www.cjd.ed.ac.uk/sites/default/files/collectionandstorage.pdf.

(5) Cerebrospinal Fluid (CSF) Sample Requirements. https://www.gloshospitals.nhs.uk/our-services/services-we-offer/pathology/tests-and-investigations/cerebral-spinal-fluid-csf-sample-requirements/.

Here is the full question in a code block:

```

26. A 33-year-old man came to the laboratory to check his blood and urine glucose status. The technologist performed the analysis using specific glucose oxidase reagent strip tests. The patient’s blood glucose status was normal while urinary glucose level was significantly increased.

What could be the most probable cause of the false positive result?

(A) The urine container may contain strong reducing agent

(B) The urine container may be contaminated with boric acid

(C) The urine container may contain strong oxidizing agents
```

(D) The container may be contaminated with formaldehyde solution
```

The question is asking for the most probable cause of a false positive result in a urinary glucose test, where the patient's blood glucose levels were normal but urinary glucose levels were significantly increased.

According to the information I found[1], false-positive results can be attributed to interfering substances in the environment where the strips are stored or used, such as hydrogen peroxide (H2O2) or bleach (hypochlorite). These substances are strong oxidizing agents, which can interfere with the glucose oxidase reagent strip tests. Therefore, the most probable cause of the false positive result in this case is that the urine container may contain strong oxidizing agents. So, the answer is **(C) The urine
```

container may contain strong oxidizing agents**[1].

Source: Conversation with Bing, 1/3/2024

(1) False-Positive and False-Negative Urine Glucose Results. https://www.labce.com/spg506344_false_positive_and_false_negative_urine_glucose_re.aspx.

(2) Urine test strip analysis, concentration range and its interpretations .... https://gsconlinepress.com/journals/gscbps/sites/default/files/GSCBPS-2021-0091.pdf.

(3) False-Positive and False-Negative Results - LabCE. https://www.labce.com/spg506366_false_positive_and_false_negative_results.aspx.

(4) undefined. https://doi.org/10.30574/gscbps.2023.22.2.0091.

Here is the full question in a code block:

```plaintext
23. A 35-year-old male patient with history of chronic renal problem came to the laboratory for glomerular filtration test. The patient arrived to the laboratory a few hours after taking a diet having heavy meat. The laboratory technologist collected both blood and 24 hours urine specimens to determine creatinine concentration.

What was the best blood specimen collection time?

(A) Before 24 hours urine collection

(B) Just after 24 hours urine collection

(C) In 4 hours of 24 hour urine collection

(D) In 2 hours of 24 hour urine collection
```

The question is asking about the best time to collect a blood specimen for a glomerular filtration test, which is used to determine creatinine concentration. Creatinine is a waste product that comes from normal wear and tear on your muscles. If your kidneys aren't working well, creatinine can build up in your blood[1].

According to the information I found[3], the blood sample must be taken within 24 hours of the day that you collect your urine sample. Therefore, the best blood specimen collection time would be **just after the 24 hours urine collection**. So, the answer is **(B) Just after 24 hours urine collection**[3].

Source: Conversation with Bing, 1/3/2024

(1) Glomerular Filtration Rate (GFR) Test - MedlinePlus. https://medlineplus.gov/lab-tests/glomerular-filtration-rate-gfr-test/.

(2) Creatinine Clearance - Testing.com. https://www.testing.com/tests/creatinine-clearance/.

(3) Kidney Function Tests: Types, Results & Follow Up - Cleveland Clinic. https://my.clevelandclinic.org/health/diagnostics/21659-kidney-function-tests.

(4) Glomerular Filtration Rate (Estimated) | ARUP Laboratories Test Directory. https://ltd.aruplab.com/Tests/Pub/3005478.

Here is the full question in a code block:

```plaintext
32. A 45-year-old male patient with chronic glomerulonephritis and long-lasting extremities edema came to the laboratory for serum albumin and urinalysis tests. The laboratory technologist received and analyzed the sample and result was serum
```

creatinine of 1.7 mg/dl (RI: 0.7-1.4g/dl) and serum albumin of 15g/l (RI: 30-45g/l).

What would happen to the concentration of urinary protein?

(A) It will be normal

(B) It will be decreased

(C) It will be increased

(D) There will be no protein

```

The question is asking about the concentration of urinary protein in a 45-year-old male patient with chronic glomerulonephritis and long-lasting extremities edema. The patient's lab results show a serum creatinine level of 1.7 mg/dl, which is above the reference interval (RI: 0.7-1.4g/dl), indicating impaired kidney function. The serum albumin level is 15g/l, which is
```

below the reference interval (RI: 30-45g/l), indicating hypoalbuminemia, or low levels of albumin in the blood.

Chronic glomerulonephritis is a condition that can lead to kidney damage[2]. In this condition, the glomeruli, which are tiny filters in the kidneys, become inflamed, impairing their ability to filter waste and excess fluids from the blood[2]. This can lead to an increase in the concentration of proteins in the urine, a condition known as proteinuria[4].

Therefore, given the patient's condition and lab results, the concentration of urinary protein would likely be increased. So, the answer is **(C) It will be increased**[4].

Source: Conversation with Bing, 1/3/2024

(1) Glomerulonephritis - Symptoms and causes - Mayo Clinic.

https://www.mayoclinic.org/diseases-conditions/glomerulonephritis/symptoms-causes/syc-20355705.

(2) Glomerulonephritis > Fact Sheets > Yale Medicine. https://www.yalemedicine.org/conditions/glomerulonephritis.

(3) What is Nephritis and its possible symptoms, causes, risk and prevention methods?. https://www.msn.com/en-us/health/condition/Nephritis/hp-Nephritis?source=conditioncdx.

(4) Glomerulonephritis - Kidney and Urinary Tract Disorders - MSD Manual ..... https://www.msdmanuals.com/en-gb/home/kidney-and-urinary-tract-disorders/kidney-filtering-disorders/glomerulonephritis.

(5) Glomerulonephritis - NHS. https://www.nhs.uk/conditions/glomerulonephritis/.

Certainly! Here's the question and its choices in a code block, followed by a detailed explanation:

```plaintext
33. A laboratory technologist working at district laboratory received semen specimen of a 45-year-old man. He started to analyze the specimen two hours after collection but the semen was still clotted and the sperm cell motility was abnormally decreased.

Which procedure was missed by the technologist?

(A) Mixing

(B) Centrifugation

(C) Smearing

(D) Liquefaction
```

The correct answer is **(D) Liquefaction**. Here's why:

- **Liquefaction** is a process that normally occurs within **15-30 minutes** after ejaculation. During this process, enzymes from the prostate gland act on the coagulated semen to liquefy it. This is important because it allows the sperm cells to become motile and capable of fertilizing an egg.

- In the scenario described, the semen was still clotted two hours after collection, which suggests that liquefaction did not occur as it should have. The lack of liquefaction would result in decreased sperm cell motility, as observed by the technologist.

- The other options, (A) Mixing, (B) Centrifugation, and (C) Smearing, are procedures that might be performed after liquefaction for further analysis but are not responsible for the initial liquefaction of the semen.

Therefore, the missed procedure that led to the semen remaining clotted and the sperm cell motility being abnormally decreased is **liquefaction**. It's a crucial step in semen analysis that ensures the sample is in a proper state for accurate assessment of sperm health and motility.

Source: Conversation with Bing, 1/4/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/question-3-according-text-basic-premise-household-production-model-people-productive-two-p-q39095419.

(2) . https://www.toppr.com/ask/question/lane-lo-noctane-x-benzene-to-biphenyl912-explain-whyv-the-dipole-moment-of-chlorobenzene-is/.

(3) . https://www.chegg.com/homework-help/questions-and-answers/question-3-25-

essay-questions-critically-discuss-major-threats-biological-diversity-around-q104282250.

(4) . https://www.chegg.com/homework-help/questions-and-answers/2-small-section-hypothetical-geologic-map-use-information-given-map-produce-geological-str-q73570807.

(5) . https://www.chegg.com/homework-help/questions-and-answers/11-following-true-reaction-mechanism-intermediate-identical-activated-complex-b-know-react-q27004649.

The question from the image is:

```text
34. A junior medical laboratory technologist in a clinical chemistry unit wants to prepare an acidic buffer solution for spectrophotometric measurement. Therefore,
```

the technologist intends to start his activity by purifying the water. However, the technologist was not sure about the type of instrument supposed to be used.

What is the recommended type of instrument to be used in this case?

(A) Liquid chromatography instrument

(B) Electrophoresis instrument

(C) Distillation instrument

(D) Deionizer instrument
```

Now, let's discuss each option in detail:

- **(A) Liquid chromatography instrument**: This instrument is used to separate a mixture into its individual components. It wouldn't be the best choice for purifying water.
```

- **(B) Electrophoresis instrument**: This instrument is used to separate macromolecules like DNA, RNA, or proteins. It's not typically used for water purification.

- **(C) Distillation instrument**: Distillation is a process used to purify liquids, including water. It involves heating the liquid to create vapor and then condensing the vapor back into liquid. This could be a good option for purifying water.

- **(D) Deionizer instrument**: A deionizer instrument removes ions from water through ion exchange process, which can effectively purify the water. This would be the most appropriate instrument for the technologist to use in this case.

So, the recommended type of instrument to be used in this case is **(D) Deionizer instrument**. This instrument would effectively purify the water, making it suitable for preparing an acidic buffer solution for spectrophotometric measurement. The technologist should use a deionizer instrument to ensure the water is pure and free of ions before starting the preparation of the buffer solution. This will help to ensure the accuracy and reliability of the spectrophotometric measurements.

The question from the image is:

```text
35. A laboratory quality officer working at an advanced diagnostic clinical laboratory wants to check the performance of all chemistry analyzers. For that matter, the officer wants to prepare the standard and control solutions to undertake control test runs.
```

What is the right type of pipette need to be used to prepare these reagents?

(A) Mohr pipette

(B) Measuring pipette

(C) Serological pipette

(D) Graduated-folin pipette

```

Now, let's discuss each option in detail:

- **(A) Mohr pipette**: This is a type of pipette used for measuring exact volumes of a solution. It's not typically used for preparing reagents.

- **(B) Measuring pipette**: This is a type of pipette that allows the user to measure a volume of solution extremely precisely. It
```

could be used for preparing reagents, but there might be a better option.

- **(C) Serological pipette**: This type of pipette is often used in biology or medical labs to transport a measured volume of liquid, often as a media dispenser. It's a good choice for preparing reagents.

- **(D) Graduated-folin pipette**: This type of pipette is used for precise measurements of small amounts of liquid. It's not typically used for preparing reagents.

So, the right type of pipette to be used to prepare these reagents is **(C) Serological pipette**. This pipette would allow the laboratory quality officer to accurately measure and prepare the standard and control solutions needed to check the performance of all chemistry analyzers. The officer should use a serological pipette to

ensure the accuracy and reliability of the control test runs.

The question from the image is:

```text
36. A laboratory technologist working at a clinical chemistry laboratory received blood specimens of an 18-year-old male patient for biochemical analysis. Those blood specimens had maximum absorbance in UV light regions.

What light source should preferably be used for the above reaction?

(A) Tungsten lamp

(B) Incandescent lamp

(C) Thermo-tidalde lamp

(D) Deuterium discharge lamp
```

```

Now, let's discuss each option in detail:

- **(A) Tungsten lamp**: This type of lamp emits light in the visible spectrum and is not suitable for UV light regions.

- **(B) Incandescent lamp**: This type of lamp also primarily emits light in the visible spectrum, making it unsuitable for UV light regions.

- **(C) Thermo-tidalde lamp**: There seems to be a mistake here as there's no such thing as a "Thermo-tidalde lamp". It might be a typo or a misunderstanding.

- **(D) Deuterium discharge lamp**: This type of lamp emits light in the ultraviolet
```

(UV) range and is commonly used in spectroscopy, a technique used in biochemical analysis.

So, the right light source to be used for the above reaction, considering the specimens had maximum absorbance in UV light regions, is **(D) Deuterium discharge lamp**. This lamp would provide the necessary UV light for the biochemical analysis of the blood specimens. The technologist should use a deuterium discharge lamp to ensure the accuracy and reliability of the analysis.

Sure, here is the question and its choices:

```text
37. A laboratory technologist working at regional laboratory received CSF specimen from a 34-year-old male patient for electrophoretic separation. The technologist
```

applied the required sample on the sample well and run the electrophoresis. After the electrophoretic separation was accomplished, the technologist left the support in the electrophoresis apparatus and the separated substances start to diffuse.

Which procedure should be applied to minimize diffusion?

(A) Fixation

(B) Change buffer

(C) Change electric current

(D) Changing support medium

```

Now, let's discuss each choice in detail:

**(A) Fixation**: This process is used to preserve tissues from degradation, and to maintain the structure of the cell and of sub-
```

cellular components such as cell organelles (e.g., nucleus, endoplasmic reticulum, mitochondria). Fixation process is not typically used to prevent diffusion in electrophoresis.

(B) Change buffer: The buffer in electrophoresis serves several purposes including maintaining pH balance, providing ions that carry a current, and acting as a cooling agent against heat generated during the process. If substances begin diffusing after separation has been achieved, refreshing or changing the buffer can help maintain ion strength and pH levels ensuring that molecules stay put.

(C) Change electric current: Changing the electric current would affect the speed at which the molecules move during the electrophoresis process, but it would not

prevent diffusion after the process has completed.

(D) Changing support medium: The support medium (like agarose or polyacrylamide) is where the samples are loaded and it provides a matrix through which the samples can move. Changing the support medium would not necessarily prevent diffusion.

So, the correct answer is **(B) Change buffer**. This is because changing or refreshing the buffer can help maintain the environment needed for the molecules to stay put, thus minimizing diffusion.

Source: Conversation with Bing, 1/4/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/explain-two-mechanism-polymerisation-reactions-clear-

examples-4-marks-b-draw-different-typ-q106555536.

(2) . https://askfilo.com/user-question-answers-chemistry/7-according-to-le-chateliers-principle-adding-heat-to-a-373830353738.

(3) . https://www.chegg.com/homework-help/questions-and-answers/question-3-31-differentiate-reliability-maintainability-means-examples-explain-need-reliab-q98919111.

(4) . https://brainly.in/question/34697326.

(5) . https://www.chegg.com/homework-help/questions-and-answers/1-define-resistivity-explain-resistivity-works-depending-rocks-minerals-accordance-differe-q111103704.

Sure, here is the question and its choices:

```text
```

38. A researcher at a given core research laboratory wants to separate a mixture of proteins using chromatographic technique. For that matter, the researcher wants to separate the proteins using ion exchange chromatography. The researcher wants the cationic proteins to be eluted out more than the anionic one.

Which stationary phase should the researcher use in this case?

(A) Positively charged stationary phase

(B) Positively charged solid phase

(C) Cation exchangers

(D) Anion exchanger

```

Now, let's discuss each choice in detail:
```

(A) Positively charged stationary phase: This would attract anionic (negatively charged) proteins, not cationic (positively charged) ones. So, this is not the correct choice.

(B) Positively charged solid phase: Similar to option A, this would attract anionic proteins, not cationic ones. So, this is also not the correct choice.

(C) Cation exchangers: Cation exchangers have a negatively charged stationary phase, which would attract cationic proteins. This is the correct choice if the researcher wants the cationic proteins to be eluted out more than the anionic ones.

(D) Anion exchanger: Anion exchangers have a positively charged stationary phase, which would attract anionic proteins, not

cationic ones. So, this is not the correct choice.

So, the correct answer is **(C) Cation exchangers**. This is because cation exchangers have a negatively charged stationary phase that would attract cationic proteins, allowing them to be eluted out more than the anionic ones.

Source: Conversation with Bing, 1/4/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/1-measure-period-small-amplitudes-spring-mass-system-station-2-question-review-circular-mo-q113614301.

(2) . https://www.chegg.com/homework-help/questions-and-answers/question-2-context-discussion-module-4-means-decipher-nature-phrase-used-especially-contex-q87060515.

(3) . https://www.chegg.com/homework-help/questions-and-answers/tanner-sitting-cafe-noticed-man-stopped-along-sidewalk-pick-litter-place-trash--tanner-tho-q88107061.

(4) . https://www.chegg.com/homework-help/questions-and-answers/following-describe-idea-reproductive-barrier-correct-feature-geography-behavior-morphology-q106964227.

(5) . https://www.chegg.com/homework-help/questions-and-answers/circumstances-listed-indicate-appropriate-letter-following-types-opinions-rendered-entity--q116602442.

Bacteriology

You're welcome! I'm glad to be of assistance. 😊

Here is the answer to the question in the image:

```text

39. Tuberculosis Lymphadenitis is suspected from an eight-year-old female patient and fine needle aspirated (FNA) sample is sent for Acid fast staining (AFS) to confirm Mycobacterium tuberculosis. After fixing the FNA sample on the microscopic slide, the laboratory technologist performs the AFS staining correctly except forgetting to add the primary stain.

What will the microscopy result look like?

(A) Blue rod bacteria with red background

(B) Red rod bacteria with blue background

(C) Blue rod bacteria with colorless background

(D) Colorless rod bacteria with blue background

```

**The correct answer is **(D) Colorless rod bacteria with blue background**.**

****Explanation:****

**- The primary stain in AFS is **carbol fuchsin**, which is a red dye that binds to the **mycolic acid** in the cell wall of **Mycobacterium tuberculosis**.**

**- The secondary stain in AFS is **methylene blue**, which is a blue dye that stains the **background** and the **other bacteria** that do not have mycolic acid.**
```

- If the primary stain is forgotten, the **Mycobacterium tuberculosis will not be stained by carbol fuchsin, and will remain **colorless** under the microscope.**

- The background and the other bacteria will still be stained by methylene blue, and will appear **blue under the microscope.**

- Therefore, the microscopy result will look like **colorless rod bacteria with blue background.**

**I hope this helps you understand the question and the answer better. If you have any more questions, please send me another photo and I will try to answer it. **

Here is the answer to the question in the image:

```text

40. A laboratory technologist is performing gram stain. After staining all 15 wound sample smears and positive control samples, there is no gram negative bacterium found where as there are gram positive cocci and colorless rods during the microscopic examination. The laboratory technologist considers that there is a quality problem on the gram staining reagent.

Which gram staining reagent is a potential source of such error?

(A) Acetone-alcohol

(B) Lugol's iodine

(C) Crystal violet

(D) Safranin

```

The correct answer is **(A) Acetone-alcohol.**

Explanation:

- Gram stain is a differential staining technique that distinguishes bacteria based on the **thickness and **chemical composition** of their **cell walls**.**

- Gram-positive bacteria have a **thick layer of **peptidoglycan** in their cell walls, which retains the **primary stain** of **crystal violet** even after being treated with a **decolorizing agent** such as **acetone-alcohol**.**

- Gram-negative bacteria have a **thin layer of **peptidoglycan** and an outer **lipopolysaccharide** layer in their cell walls, which loses the **primary stain**

after being treated with a **decolorizing agent, and can be counterstained with a **secondary stain** such as **safranin**.**

- The steps of gram staining are as follows:

- **Fixation: The bacterial smear is heat-fixed on a slide to adhere the cells and prevent them from washing off.**

- **Crystal violet: The primary stain is applied to the smear, which stains all the cells purple.**

- **Lugol's iodine: The mordant is applied to the smear, which enhances the binding of the primary stain to the cell walls.**

- **Acetone-alcohol: The decolorizing agent is applied to the smear, which removes the primary stain from the gram-negative cells, but not from the gram-positive cells.**

- **Safranin: The secondary stain is applied to the smear, which stains the gram-negative cells pink or red, while the gram-positive cells remain purple.**

- If the **acetone-alcohol is contaminated or expired, it may not be able to **decolorize** the gram-negative cells properly, resulting in **false-positive** results. This means that the gram-negative cells will appear **purple** instead of **pink** or **red** under the microscope, and no gram-negative bacterium will be found in the samples.**

- Therefore, the **acetone-alcohol is a potential source of such error.**

Here is the question with its choices in a code block:

```
```

41. A laboratory technologist plans to inoculate ear swab in to a culture media. The interest of the technologist is to identify discrete colonies of each organism present in the specimen.

What is the most appropriate media to be used for this purpose?

(A) Semisolid media

(B) Liquid media

(C) Solid media

(D) Agar media

```

The most appropriate medium for isolating and identifying discrete colonies of each organism present in a specimen depends on the type of organisms expected and their growth requirements. Solid media, option (C), is typically used for this purpose because it allows individual colonies to grow separately from one another, making it easier
```

to identify and characterize different organisms present in the sample.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/question-4-using-words-list-describe-four-different-security-mechanisms-used-enforce-secur-q107236431.

Thank you! Here is the question with its choices in a code block:

```
42. During solid culture medium preparation, a laboratory technologist plans to kill contaminant micrograms in the culture media. The manufacturer instructs the medium to be sterilized by steam under pressure.
```

What method of sterilization is appropriate for the medium?

(A) Autoclaving

(B) Boiling

(C) Filtration

(D) Steaming

```

The appropriate method of sterilization for the medium is **autoclaving**, option (A). Autoclaving is a process that uses steam under pressure to kill microorganisms and spores[1]. It is the most common method of sterilization used in microbiology laboratories. The manufacturer's instructions indicate that the medium should be sterilized by steam under pressure, which is consistent with the autoclaving process[1].

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/42-chapter-5-electrophilic-aromatic-substitution-part-b-
```

nitration-methyl-benzoate-objectiv-q52411518.

Sure! Here is the question with its choices in a code block:

```

43. An 18-year-old female patient came to the laboratory with a request of wound abscess culture. The laboratory technologist collected the wound abscess and inoculated to mannitol salt agar, MacConkey agar and Blood agar. On gram staining from mannitol salt agar, gram positive cocci in cluster of grapes were observed.

What is the next appropriate test to be performed to confirm if they are Staphylococci?

(A) Catalase

(B) Coagulase

(C) DNase
```

(D) Hyaluronidase
```

The next appropriate test to be performed to confirm if the bacteria present are Staphylococci is **coagulase**, option (B). Coagulase is an enzyme produced by Staphylococcus aureus that causes blood plasma to clot. It is used as a diagnostic test to differentiate S. aureus from other staphylococcal species. Since the gram-positive cocci observed in the sample were in clusters of grapes, it is possible that they are Staphylococci. The other tests listed are not specific for Staphylococci.

Sure! Here is the question with its choices in a code block:
```

44. A pure colony of Neisseria gonorrhoeae is isolated from sexually active 37-year-old woman. The laboratory technologist continues to perform antimicrobial

susceptibility testing using Kirby-Bauer disk diffusion techniques.

What is the choice of culture medium to perform the test?

(A) Blood agar

(B) Nutrient agar

(C) Chocolate agar

(D) Mueller Hinton agar

```

The appropriate choice of culture medium to perform the test is **Chocolate agar**, option (C). Chocolate agar is a type of enriched agar that is used for the cultivation of fastidious bacteria, including Neisseria gonorrhoeae[2]. It is made by adding lysed red blood cells to a nutrient agar base, which provides the necessary nutrients for bacterial growth[2]. The other options listed are not appropriate for this purpose.
```

Source: Conversation with Bing, 1/8/2024

(1) Thayer–Martin agar - Wikipedia. https://bing.com/search?q=Neisseria+gonorrhoeae+culture+medium.

(2) Modified Thayer-Martin Agar: Preparation, Uses • Microbe Online. https://microbeonline.com/thayer-martin-agar-composition-preparation-uses-colony-characteristics/.

(3) Neisseria gonorrhoeae – Culture, Reference Identification and .... https://www.publichealthontario.ca/en/laboratory-services/test-information-index/neisseria-gonorrhoeae-culture.

(4) Thayer–Martin agar - Wikipedia. https://en.wikipedia.org/wiki/Thayer%E2%80%93Martin_agar.

(5) SPECIMEN COLLECTION PROCEDURES FOR GONORRHEA TESTING. https://www.dhss.delaware.gov/dph/lab/gonorrheascp.html.

(6) undefined. http://www.microbiologyinpictures.com/%29.

Sure! Here is the question with its choices in a code block:

```

45. A 49-year-old woman who was in good health until three days before hospitalization developed fever, chills, increased weakness, and a productive cough with pleuritic chest pain. Blood and respiratory cultures were positive for S. pneumoniae. Her nine-year-old child acquired infection and developed symptoms of disease that is possibly caused by the bacteria four days later.

What is the most likely disease the child has developed?

(A) Pertussis

(B) Diphtheria

(C) Meningitis
```

(D) Rheumatoid fever
```

The most likely disease the child has developed is **meningitis**, option (C). Meningitis is an inflammation of the membranes surrounding the brain and spinal cord, which can be caused by a variety of bacteria, including S. pneumoniae. The symptoms of meningitis include fever, headache, stiff neck, and a rash. Given that the child developed symptoms of disease that is possibly caused by the bacteria four days later, it is possible that the child has developed meningitis. The other options listed are not consistent with the symptoms described.

Here is the question and its choices:

```plaintext

47. A 30-year-old male driver developed generalized musculature spasms, cardiac arrhythmias, fluctuations in blood pressure, profound sweating, and dehydration a week after car accident injury. Biological sample was taken from the wound and cultured. Growth is detected on agar media as a film over the surface of the agar rather than discrete colonies and unable to ferment carbohydrates.

What is the most possible bacterium with such features?

(A) C. botulinum

(B) C. perfringens

(C) C. septicum

(D) C. tetani

```
```

The symptoms described in the question are characteristic of tetanus, a bacterial infection caused by **Clostridium tetani**. This bacterium produces a toxin that affects the nervous system, leading to muscle stiffness and spasms — the major signs of tetanus.

The fact that the growth was detected on agar media as a film over the surface of the agar rather than discrete colonies and was unable to ferment carbohydrates also aligns with the characteristics of **C. tetani**.

Therefore, the most likely answer to this question would be **(D) C. tetani**. However, it's important to note that definitive diagnosis should always be made by a healthcare professional based on a comprehensive clinical evaluation. This explanation is intended for educational purposes only.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/34-natural-selection-favor-highest-proportion-juxtamedullary-nephrons-following-species-ri-q98291343.

Here is the question and its choices:

```text
48. After bioterrorism challenge to a community living in a city, many people came to the hospital with rapid onset of sepsis with fever, edema, and lymphadenopathy; meningeal symptoms in half of the patients, and confirmed hemorrhagic pneumonia in most patients which led them to pass away. Appropriate samples were collected and examination revealed gram positive spore forming nonhemolytic colonies.
```

What is the most likely disease the people have developed?

(A) Pneumonic plague

(B) Inhalational anthrax

(C) Pyogenic pneumonia

(D) Primary Tuberculosis

```

Now, let's analyze each choice:

- **(A) Pneumonic plague**: This typically presents with rapid onset of illness characterized by high fever, chills, headache, body aches, weakness, vomiting, and nausea. However, it doesn't align well with the symptoms described in the question.

- **(B) Inhalational anthrax**: This also has an acute onset but is characterized by flu-like symptoms followed by severe respiratory
```

distress. The presence of gram-positive spore-forming bacteria found upon examination aligns with Bacillus anthracis, the causative agent for anthrax. This makes it a likely answer.

- **(C) Pyogenic pneumonia**: This involves bacterial infection leading to pus formation but does not align well with all presented symptoms.

- **(D) Primary Tuberculosis**: This generally has a slower onset and does not match the acute presentation described.

Considering all these factors, the most likely disease the people have developed is **Inhalational anthrax (B)**. This is due to the alignment of symptoms, the acute onset, and the presence of gram-positive spore-forming bacteria. However, a definitive

diagnosis would require further medical tests.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/3-agglutination-test-eight-serial-dilutions-determine-antibody-titer-set-tube-1-contained--q39506360.

Here is the question and its choices:

```text
49. A two-day-old neonate developed purulent conjunctivitis. Eye discharges were collected and examined in the laboratory which revealed gram negative intracellular diplococci observed and gram stained smear and mixed smooth and opaque colonies on chocolate agar; hence, the technologist concluded the causative agent was N. gonorrhoeae.
```

What is the most likely means of transmission?

(A) Contamination while umbilical stamping

(B) Contamination from the hospital environment

(C) Vertical transmission while the fetus is in the uterus

(D) Contamination while passing through the birth canal
```

Now, let's analyze each choice:

- **(A) Contamination while umbilical stamping**: This is unlikely as the infection is in the eyes and not near the umbilical area.
```

- **(B) Contamination from the hospital environment**: While possible, it's less likely given the specific identification of N. gonorrhoeae, a sexually transmitted infection.

- **(C) Vertical transmission while the fetus is in the uterus**: N. gonorrhoeae is not typically transmitted this way. It's usually an infection that is acquired during birth.

- **(D) Contamination while passing through the birth canal**: This is the most likely scenario. If the mother has a gonorrhea infection, the bacteria can infect the baby as it passes through the birth canal. This can lead to a condition called ophthalmia neonatorum, which can cause eye discharge and potentially lead to blindness if not treated.

So, the most likely means of transmission is **(D) Contamination while passing through the birth canal**. However, a definitive diagnosis would require further medical tests.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/46-following-statements-true-domestic-monetary-shock-less-disruptive-floating-exchange-rat-q96541988.

(2) . https://www.chegg.com/homework-help/questions-and-answers/muscles-normally-active-quiet-inspiration-external-internal-intercostal-muscles-b-external-q110516588.

(3) . https://www.chegg.com/homework-help/questions-and-answers/bioencrgy-b-state-three-3-conversion-technologics-exploration-biomass-valuable-energy-brie-q117961837.

Here is the question and its choices:

```text
50. A three-year-old child developed serous rhinorrhea, sneezing, anorexia, and low-grade fever progressing to the repetitive coughs followed by inspiratory whoops which end up with vomiting and exhaustion. There was marked lymphocytosis in complete blood count analysis. Nasopharyngeal swab was taken and inoculated in culture media enriched with charcoal starch blood and albumin; the result revealed extremely small gram-negative coccobacilli.

What is the most likely etiologic agent for the patient condition?

(A) Bordetella pertussis

(B) Legionella pneumophila
```

(C) Mycoplasma pneumonia

(D) Chlamydophila pneumoniae
```

Now, let's analyze each choice:

- **(A) Bordetella pertussis**: This bacterium causes whooping cough, a highly contagious respiratory disease. The symptoms described, including the "whoop" sound made when trying to take a breath, align with this condition. The presence of gram-negative coccobacilli also matches Bordetella pertussis.

- **(B) Legionella pneumophila**: This bacterium causes Legionnaires' disease, a form of atypical pneumonia. However, the symptoms described in the question are not typical for this disease.
```

- **(C) Mycoplasma pneumonia**: This bacterium can cause walking pneumonia, a milder form of pneumonia. But the symptoms and lab findings don't match well with this option.

- **(D) Chlamydophila pneumoniae**: This bacterium causes Chlamydophila pneumonia, which typically presents as a mild respiratory infection. The severe symptoms described in the question are not typical for this disease.

Considering all these factors, the most likely etiologic agent for the patient's condition is **(A) Bordetella pertussis**. However, a definitive diagnosis would require further medical tests.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/clade-hairy-smooth-flagella-includes-water-molds-diatoms-golden-brown-algae-clade-subsurfa-q20851463.

(2) . https://www.chegg.com/homework-help/questions-and-answers/korean-bakery-manufacturer-supplies-freshly-made-packaged-bread-pastry-supermarkets-conven-q90932653.

(3) . https://www.chegg.com/homework-help/questions-and-answers/jeremy-crilly-7-year-old-boy-diagnosed-attention-deficit-hyperactivity-disorder-adhd-provi-q77657713.

(4) . https://www.chegg.com/homework-help/questions-and-answers/gram-negative-bacteria-different-gram-positive-bacteria-gram-negative-bacteria-contain-one-q61308047.

Here is the question and its choices:

```text
51. A 35-year-old individual came to the clinic with a painful ulcer on the penis following unprotected sexual contact. The ulcer was clean-based and surrounded by an erythematous halo. Darkfield examination of the ulcer exudate revealed spirochetes.

What is the most likely disease the patient has developed?

(A) Chancroid

(B) Genital herpes

(C) Gonorrhea

(D) Syphilis
```

Now, let's analyze each choice:

- **(A) Chancroid**: This sexually transmitted infection is characterized by painful sores on the genitalia. However, it's caused by the bacterium Haemophilus ducreyi, not by spirochetes.

- **(B) Genital herpes**: This is a common sexually transmitted infection marked by genital pain and sores. But it's caused by the herpes simplex virus, not by spirochetes.

- **(C) Gonorrhea**: This sexually transmitted infection typically results in painful urination and abnormal discharge. It's caused by the bacterium Neisseria gonorrhoeae, not by spirochetes.

- **(D) Syphilis**: This is a sexually transmitted infection caused by the bacterium Treponema pallidum, which is a spirochete. The symptoms described in the question, including the ulcer (or chancre) and

the presence of spirochetes, align with this disease.

Considering all these factors, the most likely disease the patient has developed is **(D) Syphilis**. However, a definitive diagnosis would require further medical tests.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/26-differentiate-hegar-chadwick-goodell-signs-27-differentiate-probable-presumptive-signs--q33560687.

(2) . https://www.chegg.com/homework-help/questions-and-answers/48-year-old-woman-arrives-er-unconscious-unresponsive-sensory-stimuli-brain-mri-reveals-ex-q90390251.

Here is the question and its choices:

```text
52. A medical laboratory technologist wants to culture both blood and cerebrospinal fluid specimens which are collected from a patient with bacterial meningitis. The patient has been suspected of being infected with H.influenzae. The technologist wants to perform Satellitism test to confirm the diagnosis.

What is the most likely bacterial species used to do the test?

(A) S.aureus

(B) S.pyogenes

(C) S.epidermidis

(D) S.saprophyticus
```

Now, let's analyze each choice:

- **(A) S.aureus**: This bacterium is not typically used in a satellitism test for H.influenzae.

- **(B) S.pyogenes**: This bacterium is also not typically used in a satellitism test for H.influenzae.

- **(C) S.epidermidis**: This bacterium is not typically used in a satellitism test for H.influenzae.

- **(D) S.saprophyticus**: This bacterium is not typically used in a satellitism test for H.influenzae.

In a satellitism test for H.influenzae, the most commonly used bacterial species is actually **Staphylococcus aureus**. This is because

H.influenzae requires factors V (NAD) and X (heme) for growth, and S.aureus provides these factors. Therefore, the correct answer should be **(A) S.aureus**. However, a definitive diagnosis would require further medical tests.

Source: Conversation with Bing, 1/8/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/following-true-o--regular-exercise-proper-nutrition-adequate-sleep-effective-stress-manage-q90515941.

Here is the question and its choices:

```text
53. A six-year-old male patient visited the pediatrics emergency with symptoms of typhoid fever. The physician requested for confirmation of the disease from stool specimen by culture. Then the laboratory
```

technologist received stool specimen from the patient.

What are the appropriate primary culture media for the diagnosis?

(A) CLED and nutrient agar

(B) XLD and MacConkey agar

(C) Chocolate agar and tryptone soya broth

(D) Blood agar plate and Mueller Hinton agar
```

Now, let's analyze each choice:

- **(A) CLED and nutrient agar**: These media are not typically used for the isolation of Salmonella typhi, the causative agent of typhoid fever.
```

- **(B) XLD and MacConkey agar**: These are selective media used for the isolation of enteric pathogens, including Salmonella typhi. XLD (Xylose Lysine Deoxycholate) agar is both selective and differential, and MacConkey agar is used for the differentiation of gram-negative enteric bacteria.

- **(C) Chocolate agar and tryptone soya broth**: Chocolate agar is used for the cultivation of fastidious organisms, and tryptone soya broth is a general-purpose medium. They are not typically used for the isolation of Salmonella typhi.

- **(D) Blood agar plate and Mueller Hinton agar**: Blood agar is a general-purpose medium, and Mueller Hinton agar is typically used for antibiotic susceptibility testing. They are not typically used for the isolation of Salmonella typhi.

Considering all these factors, the most appropriate primary culture media for the diagnosis of typhoid fever are **(B) XLD and MacConkey agar**. However, a definitive diagnosis would require further medical tests.

Here is the question and its choices:

```text
54. A 32-year-old male patient with the symptom of urinary tract infection came to the hospital. The midstream urine is collected for microbiological analysis based on the request. The laboratory technologist inoculated the specimen on Cystine lactose electrolyte-deficient (CLED) agar by using sterile calibrated wire loop that hold 0.002mL.
```

What is the number of colonies to consider as significant growth?

(A) More than 5 colonies

(B) More than 20 colonies

(C) More than 50 colonies

(D) More than 100 colonies
```

Now, let's analyze each choice:

- **(A) More than 5 colonies**: This is too low to be considered significant in a urine culture. It could easily be due to contamination.

- **(B) More than 20 colonies**: This is also generally considered too low for a urine culture.
```

- **(C) More than 50 colonies**: While this might be considered significant in some cases, it's not the standard threshold.

- **(D) More than 100 colonies**: This is generally considered the threshold for a significant growth in a urine culture. If a single type of bacteria grows at this level, it usually indicates a urinary tract infection.

So, the number of colonies to consider as significant growth is **(D) More than 100 colonies**. However, a definitive diagnosis would require further medical tests.

Source: Conversation with Bing, 1/9/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/question-2-20-marks-waiting-time-receiving-occupational-therapy-treatment-rehabilitation-c-q88666454.

(2) . https://www.chegg.com/homework-help/questions-and-answers/help-question-1-results-poll-organized-tallying-number-times-choices-ranked-order-majority-q99415631.

(3) . https://www.chegg.com/homework-help/questions-and-answers/observed-variations-practice-patterns-different-regions-country-difficult-eliminatebecause-q18031962.

Here is the question and its choices:

```text
55. A 46-year-old female patient with lesion characterized by many erythematous macules, papules, nodules, extensive destruction of nasal cartilage and ears, and sensory loss on the extremities came to a hospital. On laboratory investigation, Ziehl-Neelsen stain from the lesion showed red bacilli.
```

What is the most likely disease caused by the pathogen?

(A) Lepromatous leprosy

(B) Extra pulmonary TB

(C) Tuberculoid leprosy

(D) Pulmonary TB

```

Now, let's analyze each choice:

- **(A) Lepromatous leprosy**: This form of leprosy is characterized by numerous lesions, including macules, papules, and nodules, as well as extensive destruction of nasal cartilage and ears. Sensory loss on the extremities is also a common symptom. The presence of red bacilli in a Ziehl-Neelsen stain is indicative of Mycobacterium leprae, the bacterium that causes leprosy.
```

- **(B) Extra pulmonary TB**: This refers to tuberculosis that occurs outside of the lungs. While it can cause a wide range of symptoms depending on the affected area, the symptoms described in the question are more characteristic of leprosy.

- **(C) Tuberculoid leprosy**: This is a milder form of leprosy with fewer lesions. The extensive symptoms described in the question are more consistent with lepromatous leprosy.

- **(D) Pulmonary TB**: This refers to tuberculosis that affects the lungs. The symptoms described in the question, such as the destruction of nasal cartilage and ears, are not typical of this disease.

Considering all these factors, the most likely disease caused by the pathogen is **(A) Lepromatous leprosy**. However, a definitive diagnosis would require further medical tests.

Source: Conversation with Bing, 1/9/2024

(1) . https://www.numerade.com/ask/question/11-the-surface-tension-of-the-layer-of-liquid-lining-the-alveoli-tends-to-pull-the-walls-inwards-so-alveoli-could-collapse-which-statements-could-explain-how-this-is-prevented-alveolar-fluid-27834/.

(2) . https://www.chegg.com/homework-help/questions-and-answers/question-2-uniform-bar-mass-m-length-l-pivoted-point-o-shown-figure-2--point-mass-m-attach-q98550486.

Here is the question and its choices:

```text
56. A 32-year-old HIV positive patient with a symptom of productive cough, fever and night sweating came to the hospital. The physician suspected for pulmonary tuberculosis and requested for the sputum AFB test. A laboratory technologist prepared Ziehl-Neelsen stain from the specimen.

What is the feature of the most probable pathogen causing the disease?

(A) Red cocci

(B) Blue cocci

(C) Red bacilli

(D) Blue bacilli
```

Now, let's analyze each choice:

- **(A) Red cocci**: This is not the typical morphology of the bacterium causing tuberculosis. Mycobacterium tuberculosis, the causative agent of tuberculosis, is a rod-shaped bacterium (bacillus), not a spherical one (coccus).

- **(B) Blue cocci**: This is also not the typical morphology of the bacterium causing tuberculosis.

- **(C) Red bacilli**: This is the typical appearance of Mycobacterium tuberculosis in a Ziehl-Neelsen stain. The bacterium is a bacillus (rod-shaped) and appears red because it retains the primary stain (carbol fuchsin) in this staining method.

- **(D) Blue bacilli**: This is not the typical appearance of Mycobacterium tuberculosis in a Ziehl-Neelsen stain. The bacterium would not decolorize and take up the blue

counterstain (methylene blue) in this staining method.

Considering all these factors, the feature of the most probable pathogen causing the disease is **(C) Red bacilli**. However, a definitive diagnosis would require further medical tests.

Here is the question and its choices:

```text
57. A 25-year-old female patient living in refugee camp came with a symptom of generalized myalgia, chills, fever and headache two weeks after joining the camp. The physician suspected epidemic typhus which was finally confirmed by laboratory examination.
```

What is the possible mode of transmission for this disease?

(A) Aerosol

(B) Contact

(C) Fecal-oral

(D) Arthropod

```

Now, let's analyze each choice:

- **(A) Aerosol**: While some diseases can be transmitted through aerosols, epidemic typhus is not typically one of them.

- **(B) Contact**: Direct contact is not the primary mode of transmission for epidemic typhus.
```

- **(C) Fecal-oral**: This is not the typical mode of transmission for epidemic typhus.

- **(D) Arthropod**: Epidemic typhus is primarily transmitted by the human body louse (an arthropod). When a person scratches the bite, they can inadvertently rub the louse feces into the wound, introducing the bacteria into their bloodstream.

Considering all these factors, the possible mode of transmission for this disease is **(D) Arthropod**. However, a definitive diagnosis would require further medical tests.

[user](#message_with_image)

continue with this one.

Here is the question and its choices:

```text
```

58. A 30-year-old female patient came to the hospital with a symptom of fever, headache, and stiff neck. The physician suspected for bacterial meningitis and requested for CSF analysis. The laboratory technologist performed gram stain from the CSF specimen and found gram-positive cocci in chains.

What is the most likely etiologic agent for the patient condition?

(A) Neisseria meningitidis

(B) Streptococcus pneumoniae

(C) Haemophilus influenzae

(D) Listeria monocytogenes
```

Now, let's analyze each choice:

- **(A) Neisseria meningitidis**: This bacterium can cause meningitis, but it is a
```

gram-negative diplococcus, not a gram-positive coccus.

- **(B) Streptococcus pneumoniae**: This bacterium is a gram-positive coccus and often appears in pairs (diplococci) or chains. It is a common cause of bacterial meningitis in adults.

- **(C) Haemophilus influenzae**: This bacterium can also cause meningitis, but it is a gram-negative coccobacillus.

- **(D) Listeria monocytogenes**: This bacterium is a gram-positive rod, not a coccus, and it can cause meningitis, especially in newborns, pregnant women, older adults, and people with weakened immune systems.

Considering all these factors, the most likely etiologic agent for the patient's condition is **(B) Streptococcus pneumoniae**. However, a definitive diagnosis would require further medical tests.

[user](#message_with_image)

continue with this one.

[assistant to=assistant.describe_image()])

{"image_url": "https://images.cheggcdn.com/media/4f6/4f6c3a2f-7b8a-4e96-b4e6-7e8e8a4a5a6e/phpZU6FQI.png"}

Source: Conversation with Bing, 1/9/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/icd-10-cm-icd-10-pcs-inpatient-hospital-codes-diagnosis-procedure-codes-codes-q76920053.

(2) . https://www.chegg.com/homework-help/questions-and-answers/following-severe-case-food-poisoning-jim-began-trouble-

digesting-foods-onions-weeks-later--q120408870.

(3) . https://www.chegg.com/homework-help/questions-and-answers/5-consider-following-scenario-write-appropriate-sbar-communi-cation-upon-entering-patient--q79363039.

(4) . https://www.chegg.com/homework-help/questions-and-answers/diagnosis-motor-neuron-disease-past-one-half-months-64-year-old-man-experiencing-weakness--q98804259.

(5) . https://www.chegg.com/homework-help/questions-and-answers/following-regarding-neuromuscular-junction-true--axon-terminals-motor-neuron-vesicles-cont-q21752503.

I apologize for the confusion. I'm here to provide information and answer your questions to the best of my ability. The question you asked is about a medical procedure related to syphilis testing. Here's the detailed explanation:

The question is:

```plaintext
59. A 19-year-old college student with history of multiple sexual partners came to the hospital with chancre on the left side of his penis. The physician suspected syphilis and requested for dark field microscopic examination.

What is the appropriate site of sample collection for the case?

(A) At the periphery of the lesion

(B) At the center of the lesion

(C) Discharge from urethra

(D) Inside the urethra
```

Syphilis is a sexually transmitted infection caused by the bacterium *Treponema pallidum*. The primary stage of syphilis is often marked by the appearance of a single sore (called a chancre), but there may be multiple sores. The sore is the location where syphilis entered your body.

In the context of this question, the physician suspects syphilis and wants to perform a dark field microscopic examination. This test is used to detect *Treponema pallidum* in a sample taken from the chancre.

The appropriate site for sample collection in this case would be **(A) At the periphery of the lesion**. The reason for this is that the bacteria are more likely to be found at the active, growing border of the lesion rather than at the center. The center of the lesion often contains dead tissue where the bacteria are less likely to be found.

Please note that this information is for educational purposes only and any medical concerns should be addressed with a healthcare provider.

The question and its choices are:

#60. A patient with history of travel to Leptospira prevalent area came to the hospital with complaints of fever, chills, headache, severe myalgia, and malarst. The physician requested for laboratory test that identify the serogroup of the pathogen

What is the appropriate test used for this purpose?

(A) Complement fixation

(B) Sensitized erythrocyte lysis

(C) Microscopic agglutination test

(D) Enzyme linked immunosorbent assays

The correct answer is (C) Microscopic agglutination test.

The explanation is:

Leptospirosis is a zoonotic disease caused by bacteria of the genus Leptospira, which has about 240 serovars[1]. The microscopic agglutination test (MAT) is the standard serological reference test for the diagnosis of leptospirosis, as it can detect the specific serogroup of the infecting strain[2]. MAT uses live Leptospira cultures representing different serogroups as antigens, and measures the agglutination of the bacteria by

the patient's serum antibodies under a microscope[2]. The test requires a panel of locally optimized serogroups that reflect the epidemiology of the region[2]. MAT is a technically demanding and laborious procedure, and it has a low sensitivity in the acute phase of the infection[2]. Therefore, other tests such as enzyme linked immunosorbent assays (ELISA) or polymerase chain reaction (PCR) may be used as alternatives or complements to MAT[2][3]. Complement fixation and sensitized erythrocyte lysis are not commonly used for leptospirosis diagnosis.

Source: Conversation with Bing, 1/11/2024

(1) Serological classification and grouping - Leptospirosis Information. http://www.leptospirosis.org/serovars.php.

(2) Optimizing the microscopic agglutination test (MAT) panel for the ....

https://journals.plos.org/plosntds/article?id=10.1371/journal.pntd.0009565.

(3) LEPDT - Overview: Leptospira, IgM, Serum - Mayo Clinic Laboratories. https://www.mayocliniclabs.com/test-catalog/Overview/65183.

(4) undefined. https://doi.org/10.1371/journal.pntd.0009565.

The question is:

62. A 37-year-old patient came to the private clinic seeking medical care. The physician evaluated the patient and sent a laboratory request for complete blood cell count. The laboratory technician collected blood via skin puncture and performed manual cell counting and reported the result to the physician. What parameter is most likely to be reduced in this procedure?

(A) Lymphocyte counts

(B) White cell counts

(C) Red cell counts

(D) Platelet counts

The question is:

61. A phlebotomist needs to draw blood after assembling all the required materials for the patient using the Vacutainer system. However, he was not able to draw the blood. What alternative should be used to draw blood in this situation?

(A) Micro-sampling method

(B) Use skin puncture method

(C) Use butterfly collection system

(D) Use ordinary needle and syringe method

The answer to this question depends on the specific circumstances of the patient and the phlebotomist's training and expertise. Here are some potential alternatives:

(A) Micro-sampling method: This method is often used when only a small amount of blood is needed, such as for pediatric patients or for certain types of tests. It involves using a small, sharp blade to make a shallow cut on the skin, usually on the finger, heel, or earlobe, and then collecting the drops of blood that emerge.

(B) Skin puncture method: This method, also known as a capillary blood draw, involves puncturing the skin with a lancet to collect blood. It's often used for tests that require a small amount of blood, such as blood glucose testing.

(C) Butterfly collection system: This system uses a small, winged needle that's connected to a thin tube. It's often used for patients with small or difficult-to-access veins, such as children, elderly patients, or patients who are receiving chemotherapy.

(D) Ordinary needle and syringe method: This is the traditional method of drawing blood, and it involves using a needle and syringe to puncture a vein and draw blood. This method is often used when a large amount of blood is needed, such as for multiple tests.

Each of these methods has its own advantages and disadvantages, and the best choice will depend on the specific situation. The phlebotomist should choose the method that is most appropriate for the patient's condition and the amount of blood needed for the tests. It's also important for the phlebotomist to be trained and competent in the method they choose to use.

The answer to this question is:

(D) Platelet counts

The reason for this is that platelets are small cell fragments that are much smaller than red and white blood cells. They are crucial for blood clotting. When blood is collected via a skin puncture, it's possible that some platelets may stick to the sides of the

collection tube or clot at the puncture site, leading to a reduction in the number of platelets that are actually counted. This is especially true if the blood sample has to wait before it's analyzed or if it's not mixed properly after collection. Therefore, the platelet count might be artificially low in this situation.

It's important to note that this doesn't necessarily mean the patient has a low platelet count (a condition known as thrombocytopenia). It could simply be an artifact of the collection process. If a low platelet count is suspected, the test may need to be repeated with a different collection method, such as venipuncture.

The other parameters - lymphocyte counts (A), white cell counts (B), and red cell counts (C) - are less likely to be affected by the blood collection method. These cells are larger and

less likely to stick to the collection tube or clot at the puncture site. However, as with any laboratory test, it's important to ensure that the sample is collected and handled properly to ensure accurate results.

The question is:

63. A six-year-old child was brought to a health centre laboratory for blood film examination. The laboratory technician made a skin puncture and collected blood on a microscopic slide. After the technician had placed a small drop of blood on a clean slide, the technician prepared a smear with the corner of another slide covering a 2cm diameter. What is the advantage of spreading the blood onto a small area?

(A) To increase the sensitivity of the blood film

(B) To increase the specificity of the blood film

(C) To increase the blood volume of the blood film

(D) To evaluate the morphology of cells in a blood film

The answer to this question is:

(D) To evaluate the morphology of cells in a blood film

Spreading the blood onto a small area, or making a thin smear, allows the cells to be spread out enough so that they can be individually examined under a microscope. This is crucial for evaluating the morphology

(shape and structure) of the cells, which can provide important information about the patient's health. For example, changes in the size, shape, or color of red blood cells can indicate conditions like anemia or sickle cell disease. Similarly, an increase in the number of a certain type of white blood cell might indicate an infection or an immune response.

The other options - increasing the sensitivity of the blood film (A), increasing the specificity of the blood film (B), and increasing the blood volume of the blood film (C) - are not typically the primary reasons for making a thin smear. While a well-prepared smear can indeed improve the sensitivity and specificity of certain tests, the main advantage is the ability to evaluate cell morphology. Increasing the blood volume of the film would actually make it more difficult to examine the cells, as they would be too crowded.

The question is:

A 4th year medical laboratory student wanted to study the prevalence of malaria among pregnant women for his graduation research. The student prepared a 10% Giemsa stain working solution to stain thin and thick blood films prepared from pregnant women via skin puncture. During examination, the student observed that a few of the slides appeared to be too blue under the light microscope.

What is the most likely cause of the problem in this procedure?

(A) Inadequate washing of the stain

(B) Too high acidity of the stain

(C) Insufficient staining time

(D) Too thin films

The most likely cause of the slides appearing too blue is **(B) Too high acidity of the stain**[1][2].

Giemsa stain is a differential stain that contains a mixture of azure, methylene blue, and eosin dye. It is specific for the phosphate groups of DNA and attaches itself to regions where there are high amounts of adenine-thymine bonding[2]. If the slides appear too blue, it could be due to the high acidity of the stain. This can cause the basophilic components (which stain blue) to be more prominent[1].

To correct this, several adjustments can be made to the staining protocol[1]:

- Reduce time in the stain/buffer mix: This is where the staining of the specimen actually occurs.

- Switch from a 7.2 pH to a 6.8 pH buffer: This will lower the pH of the buffer to be more eosinophilic (instead of basophilic/too blue). This will cause more of the intensity or distinction in the red blood cells and eosinophilic morphology rather than in the basophilic morphology.

- Reduce the concentration of stain in the stain/buffer solution: For example, from 1:5 (20% stain, 80% buffer) to 1:10 (10% stain, 90% buffer).

- Ensure that the buffer/stain mix ratio has not been altered by over-use and to be safe, swap the solution for a new mix. This should be done every six hours to retain the buffer ratio[1].

Source: Conversation with Bing, 1/11/2024

(1) Why Is My Wright’s Or Wright-Giemsa Stain Too Blue?. https://www.ethosbiosciences.com/why-is-wright-or-wright-giemsa-stain-too-blue.

(2) Giemsa Stain: Principle, Procedure, Results • Microbe Online. https://microbeonline.com/giemsa-stain-principle-procedure-and-results/.

(3) Why Is My Wright’s Or Wright-Giemsa Stain Too Blue?. https://bing.com/search?q=cause+of+slides+appearing+too+blue+in+Giemsa+staining.

(4) 5 Ways to Correct Precipitate in Hematology Stains. https://www.ethosbiosciences.com/correct-precipitate-in-hematology-stains.

(5) How To Improve Nuclear Staining in Histology Slides. https://www.ethosbiosciences.com/improve-nuclear-staining.

The question is:

```text
65. A laboratory technician collected blood via skin puncture to enumerate the total number of white blood cells for a patient diagnosed with leukemia. The total manual leukocyte count for the patient is 60,000cells/mm^3. Since the total white cell count is high, the technician repeats the count by mixing 0.76ml of diluting fluid with 20µl of blood.

What is the approximate dilution of the solution in this procedure?

(A) 1:10

(B) 1:20

(C) 1:80

(D) 1:40
```

The approximate dilution of the solution in this procedure is **(B) 1:20**[6].

In the given scenario, the technician is mixing 0.76ml of diluting fluid with 20µl of blood. To calculate the dilution, we need to convert the volumes to the same units. Since 1ml = 1000µl, 0.76ml of diluting fluid is equivalent to 760µl. Therefore, the total volume is 760µl (diluting fluid) + 20µl (blood) = 780µl.

The dilution factor is calculated by dividing the total volume by the volume of the sample. In this case, it would be 780µl / 20µl = 39, which is approximately 40. However, in laboratory practice, the dilution is often rounded to the nearest ten, hence the dilution is approximated as 1:40[1][2][4][5].

I hope this clarifies your question. Let me know if you have any other questions! 😊

Source: Conversation with Bing, 1/11/2024

(1) WBC manual count using hemocytometer - جامعة الملك عبد ....
https://kau.edu.sa/Files/0007058/Files/60503_WBC%20manual%20count%20using%20hemocytometer.pdf.

(2) White blood cell (Total leucocyte) count and TLC solution formula.
https://labpedia.net/white-blood-cell-part-2-total-leukocytes-count/.

(3) WBC Count Method - WBC Meaning, Requirements, Procedure & Calculation ....
https://biologyreader.com/wbc-count-method.html.

(4) estimation of total leukocyte count (TLC): - Medical Students Corner.
https://medicscenter.com/estimation-total-leukocyte-count-tlc/.

(5) How is Leukocyte Count Done ? | WBC | Blood Cells | Biology.
https://www.biologydiscussion.com/hematolo

gy-2/blood-cells/how-is-leukocyte-count-done-wbc-blood-cells-biology/80437.

(6) How is Leukocyte Count Done ? | WBC | Blood Cells | Biology. https://bing.com/search?q=dilution+calculation+in+leukocyte+count.

The question is:

```text
66. A physician sent a laboratory request for complete blood cell counting of a 25-year-old patient, who previously has received blood transfusion. The laboratory technologist performed manual complete blood cell counting and found a high number of white blood cells. The laboratory technologist suspected a falsely high value and wanted to repeat the count before reporting the result to the physician.
```

What is the most likely cause for the increased white blood cell count?

(A) Not allowing sufficient time for the cells to settle in the chamber

(B) Incorrect measurement of blood volume due to poor technique

(C) Not correcting a white cell count for nucleated RBCs

(D) Inadequate mixing of blood with diluting fluid
```

The most likely cause for the increased white blood cell count is **(C) Not correcting a white cell count for nucleated RBCs**[1][2][3].
```

In a manual complete blood cell count, nucleated red blood cells (RBCs) can be mistaken for white blood cells (WBCs) because they are similar in size and appearance. This can lead to a falsely high white blood cell count[1][2][3].

Nucleated RBCs are immature red blood cells that still contain a nucleus. They are typically not present in peripheral blood, but can be increased in certain conditions such as severe anemia, bone marrow disorders, or following a blood transfusion[1][2][3].

If nucleated RBCs are present, the white blood cell count should be corrected to account for them. The corrected WBC count can be calculated using the following formula[1][2][3]:

$$\text{Corrected WBC count} = \frac{\text{Uncorrected WBC count}}{1 + (\text{Number of nucleated RBCs}/100)}$$

I hope this clarifies your question. Let me know if you have any other questions! 😊

Source: Conversation with Bing, 1/11/2024

(1) What is High white blood cell count and its causes and related conditions?. https://bing.com/search?q=cause+for+increased+white+blood+cell+count+in+manual+complete+blood+cell+counting.

(2) High white blood cell count Causes - Mayo Clinic. https://www.mayoclinic.org/symptoms/high-white-blood-cell-count/basics/causes/sym-20050611.

(3) High White Blood Cell Count: Causes, Symptoms, Tests - Verywell Health.

https://www.verywellhealth.com/high-white-cell-count-5222110.

(4) Complete blood count (CBC) - Mayo Clinic. https://www.mayoclinic.org/tests-procedures/complete-blood-count/about/pac-20384919.

(5) What is High white blood cell count and its causes and related conditions?. https://bing.com/search?q=cause+for+falsely+high+white+blood+cell+count+in+manual+complete+blood+cell+counting.

The question is:

```text
68. A laboratory technician working in a hospital wanted to calculate the MCHC of a patient. He measured the hemoglobin and the hematocrit value of the patient. The results
```

are hemoglobin of 16g/dl and hematocrit of 48%

What is the approximate value of MCHC?

(A) 30 g/dl

(B) 32.5 g/dl

(C) 33 g/dl

(D) 36.5 g/dl

```

The Mean Corpuscular Hemoglobin Concentration (MCHC) is calculated using the formula[12]:

$$\text{MCHC in g/dL} = \frac{\text{Hemoglobin in g/dL}}{\text{Hematocrit \%}} \times 100$$
```

Substituting the given values into the formula:

$$\text{MCHC} = \frac{16}{48} \times 100 = 33.33 \text{ g/dL}$$

So, the approximate value of MCHC is **(C) 33 g/dl**[12].

I hope this clarifies your question. Let me know if you have any other questions! 😊

Source: Conversation with Bing, 1/11/2024

(1) Mean Corpuscular Hemoglobin Concentration (MCHC) Calculator. https://www.thecalculator.co/health/Mean-Corpuscular-Hemoglobin-Concentration-%28MCHC%29-Calculator-1074.html.

(2) Mean Corpuscular Hemoglobin Concentration (MCHC) Calculator - MDApp.

https://www.mdapp.co/mean-corpuscular-hemoglobin-concentration-mchc-calculator-325/.

(3) Mean Corpuscular Hemoglobin Concentration Calculator - Drlogy. https://drlogy.com/calculator/mean-corpuscular-hemoglobin-concentration.

Here is the question and its choices:

```markdown
73. A sample collected with EDTA test tube was sent to the laboratory for osmotic fragility test (OFT). The laboratory technologist received the sample and prepared the necessary materials and reagents to perform the test. The technologist arranged 12 test tubes (T1-T12) and added buffered sodium chloride of
```

increased strength in each test tube and equal volume of patient's blood. After following the correct procedure, the technologist read the supernatant using spectrophotometer and considering T2 as blank whereas T1 as hemolysis standard. The absorbance reading was: T1=0.6, T12=0.0 and Ts=0.3.

What is the percentage that correctly indicates hemolysis in tube 5 (Ts)?

(A) 18%

(B) 30%

(C)50%

(D)100%

```

The osmotic fragility test (OFT) measures the resistance of red blood cells to hemolysis when they are placed in a hypotonic solution.
```

The percentage of hemolysis is calculated by comparing the absorbance reading of the test tube in question (in this case, tube 5 or Ts) to the absorbance reading of the hemolysis standard (in this case, T1).

Given that the absorbance reading for T1 (the hemolysis standard) is 0.6 and the absorbance reading for Ts (tube 5) is 0.3, we can calculate the percentage of hemolysis in tube 5 (Ts) as follows:

```latex
\text{Percentage of hemolysis in Ts} = \left(\frac{\text{Absorbance of Ts}}{\text{Absorbance of T1}}\right) \times 100\%
```

Substituting the given values:

```latex
\text{Percentage of hemolysis in Ts} = \left(\frac{0.3}{0.6}\right) \times 100\% = 50\%
```

So, the correct answer is (C) 50%. This indicates that 50% of the red blood cells in tube 5 (Ts) underwent hemolysis under the conditions of the test. Please note that this is a simplified explanation and the actual calculation may involve more complex steps depending on the specific protocol followed in the laboratory. It's always best to consult with a laboratory professional for accurate information.

Source: Conversation with Bing, 1/11/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/question-9-metal-alkene-complexes-m-n2-alkene-exist-either-n-alkene-ii-metallacycloalkane--q96327574.

(2) . https://www.chegg.com/homework-help/questions-and-answers/ay-derive-expressions-contribution-vibrational-degree-freedom-entropy-figure--express-resu-q20896707.

(3) . https://www.chegg.com/homework-help/questions-and-answers/12-value-ae-system-performs-13-kj-work-surroundings-loses-9-kj-heat-kj-22-b-22-c-4-e-13-13-q88451891.

(4) . https://www.chegg.com/homework-help/questions-and-answers/9-consider-following-dispersion-relation-propagation-electromagnetic-waves-plasma-w-w-c-k--q99157793.

(5) . https://www.chegg.com/homework-help/questions-and-answers/04-200-kva-480-v-50-hz-y-connected-six-polc-synchronous-generator-ratcu-field-current-5a-t-q86268317.

Here is the question and its choices:

```markdown

74. An 80-year-old female patient with severely inflamed wound infection came to a hospital for treatment. Based on the hematologist's request for CBC test, the laboratory technologist collected blood sample and ran the test using automated whole blood analyzer. The test result showed: total WBC count of 60x10^9/L (NR: 4-11x10^9/L) and Neutrophils count of 24x10^9/L (NR: 4-4.5x10^9/L). After getting the result, the hematologist sent the patient to the laboratory for Neutrophil alkaline phosphatase (NAP) test. The technologist performed NAP test and found high score.

What is the most likely cause for increased WBC count?

(A) Leukemia

(B) Lymphoma
```

(C) Viral Leukocytosis

(D) Leukemoid reaction

```

The patient in the question is an 80-year-old female with a severely inflamed wound infection. The CBC test results show a significantly increased total WBC count and Neutrophils count. The Neutrophil Alkaline Phosphatase (NAP) test also shows a high score.

Given these findings, the most likely cause for the increased WBC count is a Leukemoid reaction (D). A Leukemoid reaction is a condition in which the body produces an excessive number of white blood cells in response to an infection or inflammation. It is characterized by a significant increase in the number of white blood cells, particularly neutrophils, and a high NAP score. This condition is different from leukemia, which is
```

a cancer of the blood cells and typically associated with a low NAP score. Lymphoma, a cancer of the lymphatic system, and viral leukocytosis, an increase in white blood cells due to a viral infection, are less likely given the patient's symptoms and test results. Please note that this is a simplified explanation and the actual diagnosis may involve more complex steps depending on the specific clinical scenario. It's always best to consult with a healthcare professional for accurate information.

Source: Conversation with Bing, 1/11/2024

(1) Chegg. https://www.chegg.com/homework-help/questions-and-answers/patients-arriving-hospital-outpatient-clinic-select-one-three-stations-service-suppose-phy-q26352465.

(2) . https://www.youtube.com/watch?v=d0KWQyRLkyM.

Here is the question and its choices:

```markdown
75. A 66-year-old man came to a referral hospital for treatment of pain at the central bones. On physical examination he has hepatosplenomegaly. The hematologist sent a request of peripheral morphology examination to the laboratory. The laboratory technologist collected blood, prepared and stained the smear to examine under the microscope using 100 x oil immersion. Upon examination, the technologist saw numerous rods like structures in the cytoplasm of immature WBCs.

What is the most probable type of leukemia that the patient is suffering from?

(A) ALL

(B) AML

(C) CLL
```

(D) CML
```

The patient in the question is a 66-year-old man with pain in the central bones and hepatosplenomegaly. The peripheral morphology examination of his blood revealed numerous rod-like structures in the cytoplasm of immature WBCs. These rod-like structures are known as Auer rods, which are commonly seen in Acute Myeloid Leukemia (AML).

Therefore, given these findings, the most probable type of leukemia that the patient is suffering from is Acute Myeloid Leukemia (AML) (B). Please note that this is a simplified explanation and the actual diagnosis may involve more complex steps depending on the specific clinical scenario. It's always best to consult with a healthcare professional for accurate information.
```

Source: Conversation with Bing, 1/11/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/following-describes-cell-communication-systems-cell-signaling-evolved-recently-systems-imm-q100095331.

(2) . https://www.chegg.com/homework-help/questions-and-answers/case-4-6-pts-female-patient-underwent-surgery-block-ureter-3-days-ago-experiencing-vaginal-q89263103.

Yes, different types of leukemia can have specific characteristics that help in their identification. Here are some details:

1. **Acute Lymphoblastic Leukemia (ALL)**: This type of leukemia is characterized by the presence of a large number of immature lymphocytes (lymphoblasts) in the blood and bone marrow[1].

2. **Acute Myeloid Leukemia (AML)**: AML is characterized by the rapid growth of abnormal myeloid cells. The presence of Auer rods, which are rod-like structures in the cytoplasm of immature white blood cells, is a distinctive feature of AML[1].

3. **Chronic Lymphocytic Leukemia (CLL)**: CLL is characterized by the accumulation of mature, but functionally defective, lymphocytes in the blood, bone marrow, and lymphatic tissues. It develops slowly and patients may not experience symptoms until CLL reaches a later phase[1].

4. **Chronic Myelogenous Leukemia (CML)**: CML is characterized by the overproduction of white blood cells known as granulocytes. It is associated with a specific chromosomal abnormality known as the Philadelphia chromosome[1].

5. **Chronic Eosinophilic Leukemia**: This type is characterized by too many eosinophils, a type of white blood cell that fights allergens[5].

6. **Chronic Neutrophilic Leukemia**: This type is characterized by too many neutrophils, a type of white blood cell that fights infection[5].

Please note that these are general characteristics and the actual diagnosis may involve more complex steps depending on the specific clinical scenario. It's always best to consult with a healthcare professional for accurate information.

Source: Conversation with Bing, 1/11/2024

(1) 4 Types of Leukemia: Symptoms, Treatments, and More - Healthline.

https://www.healthline.com/health/leukemia/types-of-leukemia.

(2) Types of Leukemia | Penn Medicine. https://www.pennmedicine.org/cancer/types-of-cancer/leukemia/types-of-leukemia.

(3) Leukemia: Symptoms, Signs, Causes, Types & Treatment - Cleveland Clinic. https://my.clevelandclinic.org/health/diseases/4365-leukemia.

(4) Leukemia | American Cancer Society. https://www.cancer.org/cancer/types/leukemia.html.

(5) Leukemia | Definition, Causes, Symptoms, & Treatment | Britannica. https://www.britannica.com/science/leukemia.

Here is the question and its choices:

```markdown
76. A 25-year-old patient came to the laboratory with a request of coagulation
```

profile test. The laboratory technologist arranged necessary materials for blood collection. The technician planned to collect the blood with 0.5ml of sodium citrate containing test tube.

What volume of blood should proportionally be collected?

(A) 4.5 ml

(B) 5 ml

(C) 9 ml

(D) 18 ml

```

The answer to this question is (A) 4.5 ml. Here's why:

The standard ratio of blood to anticoagulant (in this case, sodium citrate) for a coagulation profile test is 9:1. This means
```

that for every 1 part of sodium citrate, 9 parts of blood should be collected. Given that the test tube already contains 0.5 ml of sodium citrate, the volume of blood that should be collected is 9 times this amount, which equals 4.5 ml. Therefore, the correct answer is (A) 4.5 ml. This ensures that the blood does not clot and can be properly tested.

Here is the question and its choices:

```markdown
77. A 34-year-old patient who had autosomal recessive bleeding disorder came to a hospital laboratory for platelet count, and coagulation profile tests. The laboratory result of these tests revealed decreased platelet count but increased PT, APTT and TT.
```

What is the most likely cause of prolonged coagulation tests?

(A) Afibrinogenemia

(B) Dysfibrinogenemia

(C) Hypofibrinogenemia

(D) Hyperfibrinogenemia

```

Now, let's discuss each choice in detail:

- **Afibrinogenemia (A)**: This is a rare genetic disorder characterized by the absence of fibrinogen in the blood, leading to problems with blood clotting. Given the patient's symptoms, this could be a likely cause as the absence of fibrinogen would impact all aspects of the coagulation cascade, leading to increased PT, APTT, and TT.
```

- **Dysfibrinogenemia (B)**: This condition involves dysfunctional fibrinogen that affects normal blood clotting. However, it usually does not lead to a decrease in platelet count.

- **Hypofibrinogenemia (C)**: It’s characterized by low levels of fibrinogen in the blood. While this could potentially lead to increased coagulation times, it would not typically cause a decrease in platelet count.

- **Hyperfibrinogenemia (D)**: This condition involves higher than normal levels of fibrinogen in the bloodstream. However, it is usually associated with an increased risk of thrombosis, not bleeding.

Considering the patient's symptoms and the characteristics of these conditions, **Afibrinogenemia (A)** seems to be the most likely cause of the prolonged coagulation tests. However, a definitive diagnosis would

require further medical investigation. Please consult with a healthcare professional for accurate information. This explanation is intended for educational purposes only.

Source: Conversation with Bing, 1/12/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/following-statements-regarding-structure-function-blood-vessels-true-false-q94284382.

Here is the question and its choices:

```markdown
78. A 35-year-old female patient came to the laboratory with a request of PT and APTT. The laboratory technician collected blood with the correct anticoagulant and performed the test according to the standard procedure. The result indicated prolonged PT and normal APTT.
```

What is the possible cause of these results?

(A) Lupus antibodies

(B) Factor VII deficiency

(C) Prekallikrein deficiency

(D) Von Willebrand disease

```

Now, let's discuss each choice in detail:

- **Lupus antibodies (A)**: These are autoantibodies associated with lupus, an autoimmune disease. They can interfere with the clotting process and may lead to a prolonged PT. However, they would typically also affect APTT, which is not the case here.

- **Factor VII deficiency (B)**: This is a rare bleeding disorder that results from a
```

deficiency in Factor VII, a protein involved in blood clotting. It specifically affects the extrinsic pathway, which is measured by PT, while APTT (which measures the intrinsic pathway) remains normal. This makes it a likely cause of the patient's results.

- **Prekallikrein deficiency (C)**: This is a deficiency in a protein involved in the intrinsic pathway of the coagulation cascade. It would typically lead to a prolonged APTT, not PT.

- **Von Willebrand disease (D)**: This is a bleeding disorder caused by a deficiency of von Willebrand factor, a protein that helps platelets stick together and adhere to the walls of blood vessels. It typically affects both PT and APTT.

Considering the patient's symptoms and the characteristics of these conditions, **Factor

VII deficiency (B)** seems to be the most likely cause of the prolonged PT and normal APTT. However, a definitive diagnosis would require further medical investigation. Please consult with a healthcare professional for accurate information. This explanation is intended for educational purposes only.

Source: Conversation with Bing, 1/12/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/26-differentiate-hegar-chadwick-goodell-signs-27-differentiate-probable-presumptive-signs--q33560687.

(2) . https://www.chegg.com/homework-help/questions-and-answers/following-true-o--regular-exercise-proper-nutrition-adequate-sleep-effective-stress-manage-q90515941.

(3) . https://www.chegg.com/homework-help/questions-and-answers/question-4-using-words-list-describe-four-different-security-mechanisms-used-enforce-secur-q107236431.

(4) . https://www.chegg.com/homework-help/questions-and-answers/circuit-shown-fig-361-relaxation-oscillator-built-ideal-infinite-open-loop-gain-infinite-i-q88843063.

Here is the question and its choices:

```markdown
79. A physician collected bone marrow biopsy from the iliac crest of a 50-year-old patient diagnosed with CML. As soon as the bone marrow sample was collected, the physician prepared smears and sent them to the laboratory. The laboratory technologist stained the smear and examined it microscopically and found bone marrow cellularity of 80%.

What is the interpretation of the result?

(A) The marrow is hypocellular
```

(B) The marrow is normocellular

(C) The marrow is hypercellular

(D) The marrow is filled with excess fat cells

```

Now, let's discuss each choice in detail:

- **The marrow is hypocellular (A)**: This term is used when the bone marrow has fewer cells than normal. This is not likely the case here as the cellularity is 80%, which is higher than the normal range for a 50-year-old patient.

- **The marrow is normocellular (B)**: This term is used when the bone marrow has a normal number of cells. The normal cellularity for a 50-year-old is typically around 50%, so this option is also unlikely.
```

- **The marrow is hypercellular (C)**: This term is used when the bone marrow has more cells than normal. Given that the cellularity is 80%, this would be the most likely interpretation of the result.

- **The marrow is filled with excess fat cells (D)**: This would typically indicate a condition like marrow adiposity, where there is an increase in fat cells in the bone marrow. However, this is not likely the case here as the cellularity is high, indicating a high number of hematopoietic cells rather than fat cells.

So, the most likely interpretation of the result is that the marrow is hypercellular (C). However, a definitive diagnosis would require further medical investigation. Please consult with a healthcare professional for accurate information. This explanation is intended for educational purposes only.

[user](#message_with_image)

continue with this one.

Sure, here is the question and its choices:

```markdown
80. A phlebotomist collected blood from the vein of a 30-year-old female patient for complete blood counting. The laboratory technologist analyzed the sample using an automated hematology analyzer and observed abnormally increased values in all the CBC parameters.

What is the most likely cause for the increased values?

(A) Collecting blood from edematous patients

(B) Removing the tourniquet as soon as blood flow
```

(C) Drawing blood using a needle with small gauge size

(D) Applying tourniquet on the patient's arm for too long
```

Now, let's discuss each choice in detail:

- **(A) Collecting blood from edematous patients**: Edema is an abnormal accumulation of fluid in the tissues resulting in swelling. While this might affect test results, it’s not directly associated with increased CBC parameters.

- **(B) Removing the tourniquet as soon as blood flow**: This could potentially lead to inaccurate readings but isn’t typically associated with universally elevated CBC parameters.
```

- **(C) Drawing blood using a needle with small gauge size**: This can sometimes affect cell morphology or cause hemolysis but isn’t typically associated with universally elevated CBC parameters.

- **(D) Applying tourniquet on the patient's arm for too long**: This can lead to hemoconcentration which is indeed associated with artificially elevated levels of cellular elements of the blood during testing because prolonged application of a tourniquet can cause pooling of the blood.

So, the most likely cause for the increased values in all the CBC parameters is **(D) Applying tourniquet on the patient's arm for too long**. This is because hemoconcentration due to prolonged application of a tourniquet can result in artificially high readings across all CBC

parameters due to concentration of cellular elements when there’s reduced plasma volume.

Source: Conversation with Bing, 1/12/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/case-1-35-year-old-man-firefighter-severely-burned-forearm-hospitalized-4-days-ago-develop-q88655811.

(2) . https://brainly.in/question/49114619.

(3) . https://www.chegg.com/homework-help/questions-and-answers/ockle-cell-anemia-disorder-caused-mutations-hemoglobin-gene-homozygotes-normal-allele-emog-q103546438.

(4) . https://brainly.ph/question/24524808.

(5) Chegg. https://www.chegg.com/homework-help/questions-and-answers/10-condition-known-heartburn-often-caused-acid-regurgitating-stomach-esophagus-drugs-treat-q25509961.

Here is the question and its choices:

```plaintext
81. A physician sent a laboratory request for anemic patient to rule out iron deficiency anemia. A laboratory technician wants to prepare plasma from blood specimen collected using EDTA anticoagulated evacuated tube to measure plasma ferritin level.

What is the most likely action of EDTA in preventing blood clotting?

(A) By inhibiting tissue factor

(B) By chelating calcium ion

(C) By inactivating thrombin

(D) By inhibiting Factor X
```

Now, let's discuss the answer in detail:

EDTA, or Ethylenediaminetetraacetic acid, is a chelating agent that binds to metal ions in a solution. In the context of blood clotting, EDTA primarily works by **chelating calcium ions** (Choice B).

Calcium ions play a crucial role in the blood clotting process. They act as a cofactor for several enzymes involved in the coagulation cascade, a series of reactions that lead to the formation of a blood clot. By binding to these calcium ions, EDTA effectively removes them from the process, thereby inhibiting the cascade and preventing blood clotting.

The other choices provided, such as inhibiting tissue factor (Choice A), inactivating thrombin (Choice C), and

inhibiting Factor X (Choice D), are not the primary mechanisms by which EDTA prevents blood clotting. While these processes are part of the coagulation cascade, they are not directly affected by EDTA. Therefore, the correct answer is **(B) By chelating calcium ion**.

Source: Conversation with Bing, 1/12/2024

(1) . https://www.chegg.com/homework-help/questions-and-answers/following-statements-regarding-structure-function-blood-vessels-true-false-q94284382.

(2) . https://www.chegg.com/homework-help/questions-and-answers/question-15-following-true-resting-membrane-potential-neurons-typically-70-mv-arises-anion-q90381126.

(3) . https://www.chegg.com/homework-help/questions-and-answers/ockle-cell-anemia-disorder-caused-mutations-

hemoglobin-gene-homozygotes-normal-allele-emog-q103546438.

(4) . https://www.chegg.com/homework-help/questions-and-answers/main-benefit-folded-structure-cerebral-cortex-protects-spinal-chord-damage-creates-space-f-q100730901.

(5) . https://www.chegg.com/homework-help/questions-and-answers/match-following-drugs-appropriate-drug-classification-classifications-may-used--worth-5-po-q100232084.

Sure, here is the question and its choices:

83. A technologist took blood from a donor, immediately after donation, for performing blood typing. The technologist used anti-A and anti-B antiserum reagents for testing unknown cells of the donor’s blood.

What is the most appropriate technique used by the technologist?

(A) Reverse typing

(B) Forward typing

(C) Indirect typing

(D) Serum typing

The answer is (B) Forward Typing.

In Forward Typing, the patient’s red blood cells are tested directly with known antisera (like anti-A or anti-B). This allows us to observe whether there is agglutination (clumping of cells) or no agglutination. The presence or absence of agglutination helps to determine the ABO group of those red cells. In this case, since the technologist used anti-A and anti-B antiserum reagents for testing the unknown cells of the donor’s blood, it is a clear example of forward typing. This is the most common method used for blood typing.

Sure, here is the question and its choices:

84. A laboratory technologist received a patient’s cross-match request prior to an appointed surgery. The laboratory technologist assembled all the necessary equipment and took blood samples from both the patient and the donor to perform a major cross-match.

What is the most appropriate donor’s specimen he should use for cross-match?

(A) Serum

(B) Plasma

(C) Buffy coat

(D) Red blood cells

The answer is (D) Red Blood Cells.

In a major cross-match, the donor’s red blood cells are tested with the recipient’s serum to determine the presence of any antibody which may cause hemolysis or agglutination of donor red cells1. This is more important than a minor cross-match, which involves testing of the donor’s plasma with the recipient’s red cells1. The major cross-match is required for any product that contains > 2 mL of red blood cells2. Therefore, the most appropriate donor’s specimen for a cross-match would be red blood cells. This ensures that the recipient’s antibodies will not react with the antigens on the donor’s red blood cells, preventing a transfusion reaction1.

Sure, here is the question and its choices:

85. A blood donation campaign was prepared by blood bank staffs in a University. During screening of students for selecting potential

blood donors, two out of 40 students were deferred.

What could be the most appropriate reason for their deferral?

(A) History of blood transfusion 2 years ago

(B) No evidence of chronic disease

(C) Having a weight of >50 Kg

(D) Hemoglobin levels<12g/dl

Sure, I'll continue.

In blood donation, one of the common reasons for deferral is low hemoglobin levels. Hemoglobin is a protein in your red blood cells that carries oxygen to your body's organs and tissues and transports carbon dioxide from your organs and tissues back to your lungs. If a donor's hemoglobin level is less than 12g/dl, they are usually deferred

from donating blood. This is because taking blood from a person with low hemoglobin could harm their health. Therefore, the most appropriate reason for the deferral of the two students could be that they had hemoglobin levels less than 12g/dl (Choice D).

Please note that this is a hypothetical scenario and the actual reason for deferral could be different based on the individual's health condition and the specific guidelines followed by the blood bank. It's always important to consult with healthcare professionals for accurate information.

Sure, here is the question and its choices:

86. A 28-year-old pregnant mother came to the laboratory with a request for Rh blood grouping. After all procedures have been completed, the laboratory technician reported the result as Rh positive.

What is the most likely antigen found in the mother's RBC?

(A) RhC

(B) RhD

(C) RhE

(D) Rhe

The answer is (B) RhD.

The Rh blood group system consists of over 50 antigens, among which the five antigens D, C, c, E, and e are the most important1. The RhD antigen is the most significant one and is the primary determinant of the Rh trait2. When we say a person is Rh positive, it typically means that they have the RhD antigen on their red blood cells3. Therefore, if the laboratory technician reported the result as Rh positive for the pregnant mother, the most likely antigen found in the mother’s red blood cells would be RhD.

Here is the question and its choices in a code block:

87. A laboratory technologist, working in screening of transfusion transmissible infections, is going to detect antibodies of HIV-1/2 and p-24 antigen of this virus using fourth generation ELISA. After following the correct procedure for ELISA testing, the technologist observed positive test result. He repeated the test and gets the same result.

What is the most appropriate confirmatory test that the technologist should use?

(A) Polymerase chain reaction

(B) Competitive ELISA

(C) Indirect ELISA

(D) Unigold test

The most appropriate confirmatory test that the technologist should use is (A) Polymerase chain reaction (PCR). Here is a detailed explanation:

PCR is a technique that amplifies a specific segment of DNA or RNA from a sample, allowing for the detection of very low amounts of viral genetic material1. PCR can be used to confirm the presence of HIV-1/2 by targeting specific regions of the viral genome that are conserved and unique to the virus2. PCR can also differentiate between HIV-1 and HIV-2, which have different genetic sequences and clinical implications3.

Competitive ELISA is a type of ELISA that measures the amount of antigen in a sample by competing with a known amount of labeled antigen for a limited number of antibody binding sites4. Competitive ELISA can be used to detect antibodies of HIV-1/2, but it is not a confirmatory test because it does not provide information on the

specificity or identity of the antibodies5. Competitive ELISA can also be affected by factors such as antigen concentration, antibody affinity, and assay conditions.

Indirect ELISA is another type of ELISA that detects antibodies in a sample by binding them to a specific antigen coated on a plate, and then adding a secondary antibody that is labeled with an enzyme or a fluorescent dye. Indirect ELISA can also be used to detect antibodies of HIV-1/2, but it is not a confirmatory test either because it can produce false-positive results due to cross-reactivity with other antibodies or nonspecific binding. Indirect ELISA can also be influenced by factors such as antigen purity, antibody quality, and assay optimization.

Unigold test is a rapid immunoassay that detects antibodies of HIV-1/2 in a sample by using a lateral flow device that contains a test line coated with recombinant HIV antigens and a control line coated with goat

anti-mouse antibodies. Unigold test can provide a result in 10 minutes, but it is not a confirmatory test because it has a lower sensitivity and specificity than ELISA or PCR, and it can also generate false-positive or false-negative results due to technical errors, sample quality, or interference from other substances .

Therefore, PCR is the most appropriate confirmatory test for the technologist to use, as it can provide a definitive diagnosis of HIV-1/2 infection by detecting the viral genetic material directly, with high accuracy and reliability. PCR can also complement the results of the fourth generation ELISA, which detects both antibodies and p-24 antigen of HIV-1/2, by confirming the presence and type of the virus. PCR is widely used as a confirmatory test for HIV diagnosis in clinical and research settings .

Sure, here is the question and its choices:

88. Severely bleeding patient came to the emergency department. A cross-match test was requested and the laboratory technologist performed ABO and Rh typing prior to cross matching with the donor. The cross-match test report was that the patient's serum was compatible with the donor’s RBC.

What is the appropriate interpretation of the reported result?

(A) The patient cannot be transfused with the donor's blood.

(B) There are no antibodies in patients serum against donor’s RBC

(C) Both patient and donor have ABO antibodies against to each other

(D) The presence of antigens in patient’s serum against donor’s antibodies

The correct answer is (B) “There are no antibodies in patients serum against donor’s RBC”.

Here’s why: A cross-match test is performed to determine if the donor’s blood is compatible with the recipient’s blood. In this case, the patient’s serum was found to be compatible with the donor’s red blood cells (RBCs). This means that there are no antibodies present in the patient’s serum that would react negatively with the antigens on the donor’s RBCs. If such antibodies were present, it would indicate an incompatible match, making transfusion unsafe. Therefore, the appropriate interpretation of the reported result is that there are no antibodies in the patient’s serum against the donor’s RBCs. This indicates that the transfusion can proceed safely.

The other options (A, C, D) are incorrect because they suggest either incompatibility between the patient’s and donor’s blood (A, D) or the presence of antibodies in both the patient’s and donor’s blood ©, neither of which is supported by the information given in the question.

Sure, here is the question and its choices:

89. A formalin fixed cone biopsy was sent from pathology OPD to the laboratory for tissue processing. The laboratory technician received the specimen and makes it ready for the next procedure.

What is the next step of tissue processing?

(A) Fixation

(B) Clearing

(C) Embedding

(D) Dehydration

The image you provided is a question from a test or exam paper. It describes a scenario where a formalin fixed cone biopsy was sent from a pathology OPD to the laboratory for tissue processing. The laboratory technician received the specimen and prepares it for the next procedure. The question asks, “What is the next step of tissue processing?” with four multiple-choice options: (A) Fixation, (B) Clearing, © Embedding, and (D) Dehydration.

Given that the biopsy is already fixed in formalin, the next step in the tissue processing sequence is typically dehydration. So, the answer would be (D) Dehydration. However, please verify this with your study materials or instructor.

Sure, here is the question and its choices:

90. A pathologist requested a biopsy to the histopathology laboratory for tissue processing of glycogen staining. The laboratory technologist received the specimen and processed it following the standard procedure. Then, decided to stain the tissue with special staining reagent that better demonstrate glycogen.

What is the most appropriate stain the technologist should use?

(A) Periodic acid Schiff stain

(B) Mucicarmine satin

(C) Neutral fast red

(D) Silver stain

The correct answer is (A) “Periodic acid Schiff stain”.

Here’s why: The Periodic Acid Schiff (PAS) stain is a staining method used in histology. It is used to detect polysaccharides such as glycogen, and mucosubstances such as glycoproteins, glycolipids and mucins in tissues. The PAS stain accomplishes this by oxidizing the polysaccharides to form aldehyde residues. These residues then react with the Schiff reagent to give a magenta color. Therefore, if a laboratory technologist wants to better demonstrate glycogen in a tissue sample, the PAS stain would be the most appropriate choice.

The other options (B, C, D) are incorrect because they are not typically used for glycogen staining. Mucicarmine stain is used to stain acidic mucins, Neutral fast red is a nuclear counterstain, and Silver stain is used to detect proteins and nucleic acids. None of these would be as effective as the PAS stain for demonstrating glycogen in a tissue sample.

Sure, here is the question and its choices:

91. A laboratory technologist drew blood sample from a patient for widal test. Then separated the serum and performed the test. The technologist observed large black clumps in the sample’s circles, indicating a positive reaction.

What should the laboratory technologist do to ensure the validity of the result?

(A) Run a positive and negative control serum in parallel

(B) Show the result to the physician for confirmation

(C) Show the result to her colleague for confirmation

(D) No further step is required for confirmation

The correct answer is (A) “Run a positive and negative control serum in parallel”.

Here’s why: In laboratory testing, it’s crucial to validate the results to ensure their accuracy. One common method of validation is to run control samples alongside the patient’s sample. These control samples, which have known outcomes, serve as a benchmark to confirm that the test is working correctly.

A positive control is a sample that is expected to produce a positive result, while a negative control is expected to produce a negative result. If the controls produce the expected outcomes, it increases confidence in the accuracy of the patient’s test results. Therefore, to ensure the validity of the result, the technologist should run a positive and

negative control serum in parallel with the patient’s sample.

The other options (B, C, D) are not the best practices for ensuring the validity of a laboratory test result. While it may be helpful to consult with colleagues or physicians (B, C), these steps do not provide the objective, scientific validation that running control samples does. And option (D) is incorrect because further steps are indeed required to confirm the validity of the test result.

Sure, here is the question and its choices:

Question: A patient came to a laboratory for weil-felix test and the laboratory technologist collected blood sample. Then, the technologist performed the test and observed a positive agglutination. What does this result indicate?

Choices:

(A) The presence of anti-rickettsia antibodies in the sample

(B) The absence of anti-proteus antibodies in the sample

(C) The presence of rickettsia antigens in the sample

(D) The presence of proteus antigens in the sample

Now, let's discuss the answer and its explanation:

The **Weil-Felix test** is a nonspecific agglutination test which is used to detect certain antibodies associated with rickettsial infections. In this test, a patient's serum is mixed with Proteus bacteria. If the patient has antibodies against rickettsia, these

antibodies will react with the Proteus bacteria, causing them to clump together, or agglutinate.

A **positive agglutination** result in a Weil-Felix test indicates that there are anti-rickettsia antibodies present in the patient's blood sample. This suggests that the patient has been exposed to rickettsia, and their immune system has produced antibodies against it.

Therefore, the correct answer is **(A) The presence of anti-rickettsia antibodies in the sample**. The other choices (B, C, D) are incorrect because they either refer to the absence of antibodies or the presence of antigens, neither of which can be determined by the Weil-Felix test. The test is designed to detect the presence of antibodies, not antigens, and a positive result indicates the presence, not absence, of these antibodies.

Sure, here is the question and its choices:

```markdown
93. A laboratory technologist was using immunochromatographic approach to check for pregnancy. The technologist took urine sample from the client and added two drops of the sample to the sample area of the kit and observed two red lines both on the C and T regions of the test kit.

What was pre-coated to the test band region with regard to the principle of this test?

(A) anti-hCGAb

(B) hCG antigen

(C) colloidal gold

(D) anti-hCG-hCG
```

Now, let's discuss the answer and its explanation:

The **immunochromatographic test** for pregnancy is a rapid diagnostic test that uses the principle of immunochromatography where the formation of antigen-antibody complexes occur. In the context of a pregnancy test, the test detects the presence of the hormone human chorionic gonadotropin (hCG), which is produced during pregnancy.

In a typical pregnancy test, the test band region is pre-coated with **anti-hCG antibodies**. When the urine sample is added, if hCG is present in the urine, it will bind to the anti-hCG antibodies, forming an antigen-antibody complex. This complex will then bind to other anti-hCG antibodies that are labeled with a color-changing dye, which

are also present in the test. This is what causes the appearance of the second line in the test region (T) of the kit, indicating a positive result.

Therefore, the correct answer is **(A) anti-hCGAb**. The other choices (B, C, D) are incorrect because they either refer to the antigen (hCG), the label used in the test (colloidal gold), or a complex that is not typically pre-coated on the test strip (anti-hCG-hCG).

Sure, here is the question and its choices:

```markdown
94. A 30-year-old man visited a hospital with radiating back pain and duodenal ulcer. The physician intended to screen for possible reasons of H. pylori infection. However, the physician was not sure which diagnostic test
```

to use, and consulted a senior laboratory technologist to affirm the test type.

Which diagnostic test is recommended to screen the patient?

(A) H. pylori stool antigen test

(B) H. Pylori antibody test

(C) H. Pylori DNA test

(D) Urease test

```

Now, let's discuss the answer and its explanation:

**Helicobacter pylori (H. pylori)** is a type of bacteria that can infect the stomach and is a common cause of peptic ulcers. Diagnosing an H. pylori infection can involve several types of tests, including stool tests, blood tests, breath tests, and biopsies.
```

In this case, the patient has a duodenal ulcer, and the physician wants to screen for H. pylori infection. The most appropriate test would be the **H. pylori stool antigen test (A)**. This test can accurately identify active infections because it detects the presence of H. pylori antigens in the patient's stool.

The other tests have their uses but may not be as suitable in this scenario:

- **H. Pylori antibody test (B)**: This test detects antibodies against H. pylori in the blood. However, it cannot distinguish between a current and past infection.

- **H. Pylori DNA test (C)**: This test detects the genetic material of H. pylori. It is usually done on a biopsy sample taken from the stomach lining, so it's more invasive.

- **Urease test (D)**: This is a rapid test done on a biopsy sample from the stomach lining. H. pylori produces the enzyme urease, which

breaks down urea to ammonia and carbon dioxide. However, this test is also invasive as it requires endoscopy.

Therefore, considering the patient's symptoms and the need for a less invasive yet effective test, the H. pylori stool antigen test (A) would be the recommended choice. It is non-invasive and can accurately detect an active H. pylori infection.

Sure, here is the question and its choices:

```markdown
96. A laboratory technologist was conducting a competitive ELISA experiment for quantification of HIV gp120 protein among HIV infected patients. As per the test principle, the technologist plans to use bovine serum albumin as a blocking agent. Nevertheless, bovine serum albumin was
```

stock out and he want to run the test with easily available alternative materials.

What is the most appropriate alternative agent?

(A) Peroxidase

(B) Avidin

(C) Biotin

(D) Gelatin

```

Now, let's discuss the answer and its explanation:

In an ELISA (Enzyme-Linked Immunosorbent Assay) experiment, a blocking agent is used to prevent non-specific binding of antibodies to the plate. Bovine Serum Albumin (BSA) is commonly used as a blocking agent due to its
```

stability and lack of interference with immunological reactions.

In the absence of BSA, another protein that can serve as a blocking agent is needed. **Gelatin (D)** is a good alternative. It's a protein derived from collagen and is often used as a blocking agent in ELISA experiments. It's readily available and inexpensive.

The other options listed, Peroxidase (A), Avidin (B), and Biotin (C), have specific roles in various biochemical reactions, but they are not typically used as blocking agents in ELISA. Therefore, the most appropriate alternative agent in this case would be Gelatin (D).

[user](#message_with_image)

continue with this one.