

☐ View hints ☐ View feedback

 Play All 

QUESTION 1



Drag and drop the correct answer into the space(s) provided.

A town with a population of 1000 people has an outbreak of the flu.
The growth model of the people who catch this illness is modeled by

$$f(t) = \frac{c}{1 + a e^{-bt}}$$

, where

$$\frac{c}{1 + a}$$

is the initial value (when
 $t = 0$

), c is the carrying capacity or limiting value, b is a constant determined by the growth of the virus and t is the time in days.

For this particular flu,

$$b = 0.406$$

, and there is one case at time

$$t = 0$$

.

Using this model, estimate the number of people who will have the flu after two weeks, and the rate at which that number is changing.

Knowing the limiting value of

$$c = 1000$$

, since there are only 1000 people in the town, find the value of a . Use

$$t = 0$$

,

$$f(0) = 1$$

.

$$f(0) = \frac{1000}{1+a} = 1$$

Solve for a .

$a =$

999 ✓

$$f(t) = \frac{1000}{1 + 999 e^{-0.406 t}}$$

At day 14:

$f(14) =$

$$\frac{1000}{1 + 999 e^{-0.406 (14)}} \quad \checkmark$$

$f(14) =$

227 ✓

people

The rate of growth is the derivative of the function.

$$f'(t) = \frac{d}{dt} \left(\frac{1000}{1 + 999 e^{-0.406 t}} \right)$$

$f'(t) =$

$$\frac{d}{dt} [1000 (1 + 999 e^{-0.406 t})^{-1}] \quad \checkmark$$

$f'(t) =$

$$\frac{-1000}{(1 + 999 e^{-0.406 t})^2} \frac{d}{dt} (999 e^{-0.406 t}) \quad \checkmark$$

$f'(t) =$



$$\frac{405594 e^{-0.406 t}}{(1 + 999 e^{-0.406 t})^2} \quad \checkmark$$

At day 14:

$$f'(14) = \frac{405594 e^{-0.406 (14)}}{(1 + 999 e^{-0.406 (14)})^2}$$

$$f'(14) =$$

71



people per day.

$$\frac{d}{dt} [1000 (1 + 999 e^{-0.406 t})^2]$$

$$\frac{406 e^{-0.406 t}}{(1 + 999 e^{-0.406 t})^2}$$

439

$$\frac{1000}{(1 + 999 e^{-0.406 t})^2} \frac{d}{dt} (999 e^{-0.406 t})$$

$$\frac{1000}{1 + 2a}$$

$$\frac{1000}{999 e^{-0.406 (14)}}$$

85

QUESTION 2



Drag and drop the correct answer into the space(s) provided.

The concentration of a chemical substance at time t

in a reaction can be found using the concentration formula,

$$c(t) = \frac{15}{9e^{-10t} + 3}$$

.

Using the formula, show that the derivative is always greater than 0 and that the concentration never exceeds 5.

Find the derivative of the concentration formula.

$$c'(t) =$$

$$\frac{d}{dt} \left[15(9e^{-10t} + 3)^{-1} \right]$$

Rewrite in exponential form.

$$c'(t) =$$

$$-15(9e^{-10t} + 3)^{-2} \frac{d}{dt}(9e^{-10t} + 3) \checkmark$$

$$c'(t) =$$

$$-15(9e^{-10t} + 3)^{-2}(9e^{-10t})(-10) \checkmark$$

$$c'(t) =$$

$$\frac{1350 e^{-10t}}{(9e^{-10t} + 3)^2} \checkmark$$

Find at

$$t = 0$$

.

$$c'(0) =$$

$$\frac{1350 e^{-10(0)}}{(9e^{-10(0)} + 3)^2} \checkmark$$



$$c'(0) =$$

$$9.375$$

for

$$t > 0$$

To find the limiting value of the formula, take the limit of the concentration formula:

$$\lim_{t \rightarrow \infty} c(t) = \lim_{t \rightarrow \infty} \frac{15}{9e^{-10t} + 3}$$

Substitute in

$$t = \infty$$

$$\lim_{t \rightarrow \infty} c(t) =$$

$$\lim_{t \rightarrow \infty} \frac{15}{9e^{-10(\infty)} + 3} \checkmark$$

$$\lim_{t \rightarrow \infty} c(t) =$$

$$\frac{15}{0 + 3} \checkmark$$

$$\lim_{t \rightarrow \infty} c(t) =$$

$$5 \checkmark$$

The concentration limit never exceeds 5.

$$-15(9e^{-10t} + 3)^{-2}(9e^{-10t})(10)$$

$$\lim_{t \rightarrow \infty} \frac{15}{9e^{-10(\infty)} + 5}$$

$$3.75$$

$$\frac{15}{1 + 3}$$

$$15(9e^{-10t} + 3)^{-2} \frac{d}{dt}(15e^{-10t} + 3)$$

$$\frac{1350 e^1}{(9 e^1 + 3)^2}$$

$$\frac{1350 e^t}{(9 e^{-10 t} + 3)^2}$$

QUESTION 3



Fill in the blank(s) with the correct answer.

The boiling temperature T (in celsius) of water at different pressures p (in atms) is modeled by the function

$$T(p) = 82.9 + 1.64 \ln p + 17.91 \sqrt{p}$$



Find the rate of change of the temperature when the pressure is 2 atm, rounded to two decimal places.

The rate of change of the temperature is

7.15



degrees Celsius

per atm.

QUESTION 4



Choose the correct answer from the drop-down menu.

Find the derivative of

$$f(x) = (\sin x)^x$$

.

$$y' =$$

$$(\sin x)^x (\ln(\sin x) + x \cot x) \quad \checkmark \quad \blacktriangledown$$

QUESTION 5



Choose the correct answer from the drop-down menu.

Find the derivative of

$$f(x) = (4x)^{\sqrt{x}}$$

.

$$f'(x) =$$

$$2x \ln(4x) + 4x \quad \checkmark \quad \blacktriangledown$$

QUESTION 6



Choose the correct answer from the drop-down menu.

Find the derivative of

$$y = (\sin x)^{\cos x}$$

.

$$y' =$$

$$(\sin x)^{\cos x} (-\sin x \ln(\sin x) + \cos x \cot x) \quad \checkmark \quad \blacktriangledown$$

QUESTION 7



Choose the correct answer from the drop-down menu.

A population of growing bacteria is growing modeled by the function

$$N = N_0 e^{kt}$$

, where

t

is in hours, the initial amount is

$$N_0 = 1200$$

, and the growth constant is

$$k = 0.45$$

. What is the rate of growth at

$$t = 3$$


hours.

After 3 hours, the bacteria is growing at a rate of

2083



bacteria per hour.



QUESTION 8



Choose the correct answer from the drop-down menu.

A deposit of AED 2500 is made into an account that pays interest annually. What is the relative rate of change?

The function modeling the amount of money is

$$A(t) = A_0 2^{0.5t}$$

where

$$A_0$$

is the initial amount,

$$A_0 = 2500$$

.

The relative rate of change is

35% ✓ ▼



QUESTION 9



Select the correct option.

What is the derivative of

$$y = (3x^4 + 5x)^{6x^8 + 9x}$$

?

$$\frac{dy}{dx} = (3x^4 + 5x)^{6x^8 + 9x} \left[(48x^7 + 9)\ln(3x^4 + 5x) + \left(\frac{72x^{10}}{3x^4 + 5x} \right) \right]$$



$$\frac{dy}{dx} = (3x^4 + 5x)^{6x^8 + 9x} (48x^7 + 9)\ln(3x^4 + 5x) + \left(\frac{72x^{10}}{3x^4 + 5x} \right)$$



$$\frac{dy}{dx} = (3x^4 + 5x)^{6x^8 + 9x} (48x^7 + 9)\ln(3x^4 + 5x) + (70x^{11} +$$

$u x$

$$\frac{dy}{dx} = (3x^4 + 5x)^{6x^8 + 9x} (48x^7 + 9)\ln(x) + \left(\frac{72x^{10} + 30x^7}{3x^3} \right)$$

QUESTION 10



Select the correct option.

What are the derivatives of

$$y = (3x)^x$$

?

$$\frac{dy}{dx} = (3x)^x (\ln(3x) + 3x)$$

$$\frac{dy}{dx} = (\ln(3x) + 1)$$

$$\frac{dy}{dx} = (3x)^x (\ln(3x) + 1)$$

$$\frac{dy}{dx} = (3x)(\ln(3x) + 1)$$

QUESTION 11



Select the correct option.

What are the derivatives of

$$y = x^{\cos x}$$

?



$$\frac{dy}{dx} = x^{\cos x} \left(\frac{\cos x}{x} - \ln x \cos x \right)$$



$$\frac{dy}{dx} = x^{\cos x} \left(\frac{\cos x}{x} + \ln x \sin x \right)$$



$$\frac{dy}{dx} = x^{\cos x} \left(\frac{\sin x}{x} - \ln x \sin x \right)$$

☒
$$\frac{dy}{dx} = x^{\cos x} \left(\frac{\cos x}{x} - \ln x \sin x \right)$$



QUESTION 12



Select the correct option.

What is the derivative of

$$y = (5x)^{x^2 + 3x}$$

?

☒
$$\frac{dy}{dx} = (5x)^{x^2 + 3x} (\ln(5x)(2x + 3) + x + 3)$$



$$\frac{dy}{dx} = (5x)^{x^2 + 3x} (\ln(5x)(2x + 3) + 3x)$$



$$\frac{dy}{dx} = (5x)^{x^2 + 3x} (\ln x(2x + 3) + x + 3)$$



$$\frac{dy}{dx} = (5x)^x (\ln(5x)(2x + 3) + x + 3)$$

QUESTION 13



Select the correct option.

What is the value

y'

for the function

$$y = x^{e^x}$$

?

☒ $y' = x^{e^x} e^x \left(\ln x + \frac{1}{x} \right)$



☐ $y' = x^{e^x} e^x (\ln x + x)$

☐ $y' = x^{e^x} e^x \left(\ln x + \frac{e^x}{x} \right)$

☐ $y' = x^{2e^x} \left(\ln x + \frac{1}{x} \right)$

QUESTION 14




The position of spring in motion can be modeled by the function

$$s(t) = 2e^{-t} \cos t$$

.

At what time is the velocity of the spring equal to zero? Select all that apply.



	$t = \frac{3\pi}{4}$	✓
--	----------------------	---

☐	$t = \frac{4\pi}{3}$	
--------------------------	----------------------	--

☐	$t = \frac{3\pi}{2}$	
--------------------------	----------------------	--

	$t = \frac{7\pi}{4}$	✓
--	----------------------	---

	$t = \frac{11\pi}{4}$	✓
--	-----------------------	---

☐	$t = \frac{4\pi}{11}$	
--------------------------	-----------------------	--

☐

$$t = \frac{2\pi}{3}$$

☐

$$t = \frac{4\pi}{7}$$

QUESTION 15



Find the derivative of

$$y = (7x)^{\ln x}$$

. Select all that apply.

☒

$$\frac{dy}{dx} = (7x)^{\ln x} \left(\frac{\ln(7x)}{x} + \frac{\ln x}{x} \right)$$

☐

$$\frac{dy}{dx} = (7x)^{\ln x} \left(\frac{7\ln x + \ln x}{x} \right)$$

☐

$$\frac{dy}{dx} = 7^{\ln x} x^{\ln(x-1)} (\ln(7x) + \ln x)$$

☐

$$\frac{dy}{dx} = (x)^{\ln x} \left(\frac{\ln(7x)}{x} + \frac{\ln x}{x} \right)$$

$$\frac{dy}{dx} = (7x)^{\ln x} \left(\frac{\ln(7x) + \ln x}{x} \right)$$



$$\frac{dy}{dx} = 7^{\ln x} x^{\ln x} (\ln(7x) + \ln x)$$