- May 3 has been celebrated as World Press Freedom Day every year to evaluate press freedom around the world, to defend the media from attacks on their independence and to pay tribute to journalists who have lost their lives in the exercise of their profession.

- The infant mortality rate of India according to the 2011 census was 44. It has shown a considerable decline from 129 per 1000 live births in 1971 to 110 in 1981 and from 80 in 1991 to 44 in 2011.

- The pH of water is 7.0 at about 25 deg C or about 77 deg F.
  Pure water undergoes a reversible reaction in which both H+ and OH- are generated.

- Calcium and Magnesium are major components of hard water.
  Hard water is formed when water percolates through limestone or gypsum which is high on calcium and magnesium content. Soaps do not act in hard water due to the presence of these salts but detergent is capable to give results in hard water. Water softening systems work by reducing the concentrations of minerals from the water. The total water hardness is the sum of the molar concentrations of Ca2+ and Mg2+, in mol/L or mmol/L units.

- An allosome is also referred to as sex chromosome or heterotypical chromosome. The 22 pair of chromosomes are similar in all human.
  The chromosomes of the 23rd pair are called allosomes consisting of two X chromosomes in females, and an X chromosome and a Y chromosome in males.

Click Here to Join our Telegram Cannel
Our vision is to build a learning community primely focused on those learners who has the hunger for knowledge and determination.

- Red-green color blindness is split into two different types: Whereas people affected by protan color blindness are less sensitive to red light, deuteranopia or deuteranomly (the second type of red-green color blindness) is related to sensitiveness on green light. Protans have either defective long-wavelength cones (L-cones) or the L-cones are missing at all. If they are missing it is called protanopia or sometimes red-dichromacy. Affected persons are dichromats because they have only two working cone types, short- and medium-wavelength, compared to persons with normal vision with three different cone types.

- Diastase is an enzyme group that is originally found in malt and it is produced when the germination of the seeds happens.
  Diastase enzyme is obtained from Red gram.

Click Here to Join our Telegram Cannel
Our vision is to build a learning community primely focused on those learners who has the hunger for knowledge and determination.