

Genetic
s

- *Genetics* is a branch of biology concerned with the study of gene, genetic variation, and heredity in organisms.
- *Heredity* (biological inheritance), is the passing on of one traits from parents to their offspring.

what is the relation between ?

✓ *Chromosomes*

✓ *Genes*

✓ *Alleles*


- **Chromosomes** - are thread-like structures located inside the nucleus of a cell.
 - made from DNA which is bound with histone protein

- **Gene** - is a section of DNA (therefore a section of chromosome that determines a particular feature).
For example - earlobe hang or free

- purple

- tall or

* gene is a basic physical characteristic on a chromosome.




position (locus) on a

- **Alleles** - are different “versions” of a gene

example a pea plant have purple and white allele.

■ **Homologous chromosomes** (a set of one maternal and one paternal chromosome of the same length and gene loci).



- **Dominant allele**- expressed in the homozygote and heterozygote.(represented usually by capital letter)

Example - PP

Pp

purple

purple

- appear in F₁ generation

- **Recessive allele**- are only expressed in homozygote

Example - pp

-not appear in F₁ generation

- **Co-dominant allele** -both allele are expressed in heterozygote. Both allele of a gene equally expressed.

Example - flower color of snapdragons

R-red petals

RR-red flower

r-white petals

Rr-pink flower

rr-white flower

- **Multiple allele inheritance** - these are more than two alleles of a gene.

Example - allele involved in the inheritance of blood group.


Genotype	Blood group
$I^A I^A, I^A i^O$	A
$I^B I^B, I^B i^O$	B
$I^A I^B$	AB
$i^O i^O$	O

- **Monohybrid cross**- is a genetic cross or breeding situation that relate to just one trait or feature.

The rule of inheritance was discovered by the Austrian monk Gregor Mendel .

- Mendel use the seven contrasting characteristics of pea plants in his experiments.

The contrasting characteristics are



- He crossed the trait following the same procedure for all contrasting characteristics .

Example

$PP \times Pp(\text{white})$

F1- Pp all purple

F2 - 3 purple, 1 white

- **Meiosis** – For gametes to be formed special cells in the sex organ of the organism divide by a process known as **meiosis**.


- a special type of a cell division is sexually reproducing organisms , used to produce the gametes(sex cells). The process is **Gametogenesis**.

Outcomes of meiosis


- * 4 haploid daughter cells
- * The daughter cells have only half the number of chromosome of original cells.
- * Daughter cells show genetic variation(as a result of crossing over in prophase I)

Two division of meiosis

- **Meiosis I** – chromosome that make up homologous pair are separated.



- **Meiosis II** – the chromatids that make up chromosome are separated.



- **Genetic linkage**

* if genes did not segregate, the genes are linked and inherited together because they are on the same chromosome. The two genes inherited as a single unit.

parent genotype PL/PL × pl/pl

Gametes

PL

	PL	pl
PL	PL/PL Purple, long	PL/pl Purple, long
pl	PL/pl Purple, long	pl/pl Red, round

F1 genotype PL/pl

(self-fertilised)

Gametes

male

female

PL pl

PL pl


- Crossing over in meiosis produces recombinant type of the offspring. So we get four types of offspring from a cross between two plants both heterozygous for both features. But not 9:3:3:1 ratio.
- **Cross breeding** results in hybrid vigour (heterosis) and can be used to combine desirable trait in one variety , whilst **inbreeding** is used to establish pure line.

- **Independent assortment**

- state that the inheritance of one trait is independent of the inheritance of another

Dihybrid inheritance

- the inheritance of two characteristics at the same time



- **Sex determination**

- . Gender in humans determine by x and y chromosomes.

- Male- xy

- Females- xx

- * The SRY gene found in x-chromosome determine sex in humans .

- * If SRY gene activated bi potential gonad develop to testes and the embryo is male.

- * If SRY gene is absent(or inactivated) bi potential gonad develop to ovary and the embryo is female.

- * Turner syndrome⇒ sterile female 45 chromosomes

- * Klinefelter syndrome⇒ sterile male 47 chromosomes



**results of non
disjunction of
sex
chromosomes
during meiosis**

- **sex- linked chromosome** - determined by alleles carried on the sex - chromosomes.
 - . Gene found on Y- chromosomes

Example retnitis pigmentosa

- . Gene found on X- chromosome

Example - red green color blindness
- haemophilia

- * Most sex-linked conditions are determined by recessive alleles on the x-chromosome , males are affected more frequently than female by these because they need only inherit one x-chromosome with recessive allele whilst females must inherit two.
- * Sex influenced and sex limited traits are expressed differently in two sexes. However the gene determines these trait.

- **Sex influenced trait** is more likely to occur in one sex but may sometimes occur in other sex.

Example – baldness

- **Sex limited trait** occur only in one sex and never occur in other sex.

example – gene that stimulate lactation.

DNA molecule – double helix in which the base sequence on the sense strand is complementary to the base sequence on antisense strand.

- DNA is a huge molecule made from two anti-parallel polynucleotide strand.

- The basic unit of a DNA strand is a nucleotide


- Three components of nucleotides are

 - . A phosphate group

 - . Four nitrogenous base

 - . A pentose sugar

* DNA must replicate **semi conservatively** means each new DNA molecule formed contain one strand from original DNA.



Cloning – is the process of producing genetically identical individuals of an organism either naturally or artificially.

- A clone of an organism is a group of organism that are genetically identical to each other and to organism from which they derived.
- Plants cloned using the technique of **micropropagation**.
- Animals have been cloned. The first mammal to be cloned was dolly the sheep.
- There are two types of cloning in vivo-cloning and in vitro-cloning.
- Genetic engineering – is a process in which the genome of an organism is altered, usually by having an extra gene from different organism added.
- A *genome* is an organism's complete set of DNA including all of its gene.
- A transgenic organism has had genes from a different type of organism (usually a different species) added to its genome.
- Restriction enzymes cut DNA at specific sequences, called restriction site, to leave overlapping stick ends.
- A gene inserted into a plasmid using a ligase enzyme.

- Genetically modified organisms can be used to manufacture specific products to benefit humans (for example insulin, vaccine, etc).
- Other organisms have also been genetically modified to produce increased yields.
- **Genetic fingerprinting** – is a technique of comparing the DNA of different people.
- The main stages in preparing genetic fingerprint
 - ✓ **preparation of fragments** using restriction enzyme
 - ✓ The fragments separated by **gel electrophoresis**.
 - ✓ DNA fragment transferred from the gel to nylon membrane(**blotting DNA to membrane**).
 - ✓ **Hybridization** with radio active probe made of single stranded DNA.
 - ✓ **Autoradiography** (the pattern of complementary sequence revealed using x-ray film.


- **Protein synthesis**

- ribosome is the organelle of a cell that make protein.

- the code for protein is specified by DNA.

- * Genetic code – is a triple code meaning that three bases in DNA code for amino acid in protein, only one of the strand of DNA molecules carried. The other strand is non-coding (antisense strand).

- * There are only 20 amino acids but 64 possible triplets codes.



- m RNA is similar to DNA it is built from nucleotide

How ever ,it is different from DNA in the following ways

- ✓ it is much smaller molecule.
- ✓ it is single stranded.
- ✓ the base thymine is replaced by uracil.
- ✓ the sugar in the nucleotide is ribosome.
- the triple of base in m RNA that code for amino acid are called **codons** .The mRNA codons are identical to DNA triplet except that uracil is substituted for thymine.

The main events during protein synthesis are **transcription** and **translation**.

- **Transcription** – the mechanism by which the base sequence of a section DNA representing gene is converted into complementary base sequence of mRNA occur in the nucleus.
- In the transcription of the DNA in gene RNA polymerase uses RNA nucleotide to build the single-stranded mRNA molecule that has a base sequence complementary to the strand of DNA being transcribed.
- To form mRNA by transcription only the antisense strand transcribed.
- **Translation** - the mechanism by which sequence of a base in mRNA molecule converted into sequence of amino acid in a polypeptide chain. It occurs in ribosome.
 - During translation of mRNA into polypeptide chain, tRNA molecules with **anticodon** complementary to mRNA ; their amino acids form a peptide bond and one of the tRNA molecules leaves and the process is repeated.

- The protein produced have a variety of function, they may be
structural protein - collagen keratin
enzymes - ATP synthase, DNA helicase
peptide hormones - insulin, adrenaline
anti-genes - antigen on red cells CD4
antibodies - HIV antibodies

● **Gene switch on**

- transcription factors are necessary to activate genes, these are protein that bind the promoter regions next to a gene and allow RNA polymerase to transcribe the gene.

● **Gene switch off**

- molecules of short interfering RNA (si RNA) can silence genes by degrading the mRNA transcribed from the genes.

- si RNA has the potential to treat condition such as AIDS by preventing replication of HIV and cancers by silencing genes that enhance cell division.

● **Mutation** is any spontaneous change in the genetic material of organism.

- Mutation classified into two as

1. Point mutations(gene mutations) – are spontaneous changes in a single base on DNA molecule.

different types of gene mutation are

- substitution
- addition
- deletion

- Substitution replace one base with another , because the code is degenerate, there may be no change in the amino acid coded for.
- Deletion and addition both cause a frame shift, these mutations after all this point also.
- In addition mutation an extra base added
- In a deletion mutation a base missed out during replication.

- **2 Chromosomal mutation** occur when there is any change in the arrangement or structure of chromosomes.

example of chromosomal mutation

✓ inversion

✓ deletion

non-disjunction

✓ insertion


✓ duplication

✓ chromosome


✓ translocation

types of mutation

deletion




duplication



inversion



insertion



translocation

