EDUCATIONAL ASSESSMENT AND EXAMINATIONS SERVICE (EAES)
ETHIOPIAN SECONDARY SCHOOL LEAVING CERTIFICATE EXAMINATION (ESSLCE)
Hamlie, 2016 E.C / July, 2024 G.C

SUBJECT: MATHEMATICS
SUBJECT CODE: 02

BOOKLET CODE: 110
TIME ALLOWED: 3 HOURS

GENERAL DIRECTIONS:

I. FOLLOW THE FOLLOWING RULES AND REGULATIONS IN DOING THE EXAMINATION:

- FEED THE **SUBJECT NAME, SUBJECT CODE AND BOOKLET CODE** GIVEN ABOVE ON THE SEPARATE ANSWER SHEET PROVIDED
- SELECT ONLY THE **CORRECT** ANSWER FOR EACH QUESTIONS
- USE ONLY **PENCIL** TO MARK YOUR ANSWERS
- YOUR **MARK** SHOULD BE **HEAVY** AND **DARK**
- **BLACKEN** ONLY THE LETTER OF YOUR CHOICE IN LINE WITH THE BOX ON THE SEPARATE ANSWER SHEET PROVIDED
- **ERASE** ALL UNNECESSARY MARKS COMPLETELY
- ANY FORM OF **CHEATING IS PROHIBITED** AND RESULT IN CANCELLATION OF YOUR SCORES
- YOU MUST IMMEDIATELY **STOP WORKING** WHEN TIME IS OVER
- FOLLOW **FURTHER INSTRUCTIONS** ON THE BOOKLET FOR EACH QUESTION

II. ATTENTION!

MAKE SURE THAT YOU HAVE WRITTEN ALL THE REQUIRED INFORMATION ON THE SEPARATE ANSWER SHEET CORRECTLY BEFORE YOU RETURN IT.

Join us on Telegram @qubeeacademy

DO NOT TURN THIS PAGE OVER UNTIL YOU ARE TOLD TO DO SO.

EAES WISH YOU ALL THE BEST!

CamScanner

EDUCATIONAL ASSESSMENT AND EXAMINATIONS SERVICE (EAES)
ETHIOPIAN SECONDARY SCHOOL LEAVING CERTIFICATE EXAMINATION (ESSLCE)
MATHEMATICS, Hamlie, 2016 E.C / July, 2024 G.C

Subject Code: 02 **Booklet Code: 110**
Time Allowed: 3 Hours **Number of Items: 60**

1. Let $f(x)=\frac{x+1}{x-1}$ and $g(x)=\sqrt{x+2}$ be two functions. Which one of the following is true about the combination values of f and g at $x=2$?

A. $(f.g)(2)=6$
B. $\left(\frac{f}{g}\right)(2)=\frac{2}{3}$
C. $(f+g)(2)=3$
D. $(f-g)(2)=-1$

2. Which one of the following defines an exponential function?

A. $f:\mathbb{R}\to(0,\infty)$ defined by $f(x)=b^x$, where $b>0$ and $b\neq 1$.
B. $f:\mathbb{R}\to(0,\infty)$ defined by $f(x)=x^b$, where $b>0$ and $b\neq 1$.
C. $f:(0,\infty)\to\mathbb{R}$ defined by $f(x)=b^x$, where $b>0$ and $b\neq 1$.
D. $f:(0,\infty)\to\mathbb{R}$ defined by $f(x)=x^b$, where $b>0$ and $b\neq 1$.

3. Let f be differentiable function at a number 'a' for a ∈ Domain f. In the expression $\frac{f(a+h)-f(a)}{h}$ for $h\neq 0$, as h gets closer and closer to zero from both directions, the expression becomes closer to $f'(a)$ which is equal to


A. The gradient of the normal line to the graph of f at a point (a, f(a)).
B. The gradient of the vertical line to the graph of f at a point (a, f(a)).
C. The gradient of the secant line to the graph of f at a point (a, f(a)).
D. The gradient of the tangent line to the graph of f at a point (a, f(a)).

Join us on Telegram @qubeeacademy

CamScanner

Subject Code: 02 **Booklet Code: 110**

4. In the following figure, O is the center of the circle and the line segments $\overline{AB}$ and $\overline{AC}$ are intersecting the circle at D and E respectively.
If $m(\angle ACD) = 30°$ and $m(\angle BEC) = 50°$, then which one of the following is equal to $m(\angle BAC)$?



A. 70° C. 20°
B. 80° D. 30°

5. Let p and q be propositions. Which one of the following propositions is a tautology?
A. $(p \Rightarrow \neg q) \Leftrightarrow (p \wedge q)$ C. $(p \wedge q) \Rightarrow (p \vee q)$
B. $(\neg p \Rightarrow q) \Rightarrow (p \wedge q)$ D. $(p \vee q) \Rightarrow (p \wedge q)$

6. If 180 individuals can complete a job in 15 days by working 10 hours a day, how many individuals are needed to complete the job in 10 days if they work 12 hours a day?
A. 200 C. 225
B. 144 D. 220


If the measure of each interior angle of a regular polygon is 135°, then which one of the following is the number of lines of symmetry of the polygon?
A. 7 C. 9
B. 6 D. 8

Join us on Telegram @qubeeacademy

CamScanner

Subject Code: 02 **Booklet Code: 110**

8. Consider the following graph of a given exponential function f(x).



Which one of the following functions represents the graph shown above?

A. $f(x) = (2)^{x-2}$ C. $f(x) = (2)^{x} - 2$

B. $f(x) = \left(\frac{1}{2}\right)^{x-2}$ D. $f(x) = \left(\frac{1}{2}\right)^{x} - 2$

9. What is the perimeter of a regular hexagon of radius 6cm?

A. $18\sqrt{3}$ cm C. 36 cm

B. 18 cm D. $36\sqrt{3}$ cm

10. Which one of the following is a quantitative data?

A. Measuring the heights of each students in the given class.
B. Determining the tastes of foods in a restaurant.
C. Identifying the social status of individuals.
D. Checking the quality of a new product.

11. Which one of the following is true about the integral of a function f(x) with respect to a variable x?

A. $\int f'(x)dx$ is the set of all anti-derivatives of f.
B. $\int f'(x)dx$ is the set of all derivatives of f.
C. $\int f(x)dx$ is the set of all anti-derivatives of f.
D. $\int f(x)dx$ is the set of all derivatives of f.

join us on telegram @ujueeacademy

...WER SHEET ...L THE REQUIRED CORRECTLY


...URN THIS PAGE OVER UNTIL YOU ARE TOLD TO DO SO

EAES WISH...

CamScanner

12. Which one of the following is **NOT** a valid reason for saving?
    A. For betting in football
    B. For vacation
    C. To maximize interest
    D. For the time of retirement

13. The shaded region in the following figure is the graph of a relation R.



Which one of the following is the domain and range of R respectively?

A. $(-\infty, -\frac{1}{2}]$ and R

B. R and $\left(-\infty, -\frac{1}{2}\right)$

C. R and $\left(-\infty, -\frac{1}{2}\right]$

D. $\left(-\infty, -\frac{1}{2}\right)$ and R

Join us on Telegram @qubeeacademy

CamScanner

14. Which one of the following statements is the polynomial division theorem?

A. Let f is a non-constant polynomial with integer coefficients. If the rational number $\frac{c}{d}$ in its lowest terms is a root of f, then c is a factor of the constant term of f and d is a factor of the leading coefficient.

B. If f(x) and d(x) are polynomials such that $d(x) \neq 0$ and degree(d(x)) ≤ degree(f(x)), then there exist unique polynomials q(x) and r(x) such that $f(x) = d(x)q(x)+r(x)$, where $r(x) = 0$ or degree of r(x) < degree of d(x).

C. If f is a non-constant polynomial and c is a real number, then the remainder when f(x) divided by $x - c$ is f(c).

D. If f is a non-constant polynomial and c is a real number, then $x - c$ is a factor of f(x) if and only if $f(c) = 0$.

15. Which one of the following is equal to $\cot(1755^\circ)$?

A. $\sqrt{3}$ C. 1

B. $-\sqrt{3}$ D. -1

16. Let p and q be proposition with truth values true and false respectively. Which one of the following statements is **NOT** correct?

A. $p \Rightarrow q$ is false. C. $p \wedge q$ is false.

B. $p \Leftrightarrow q$ is true. D. $p \vee q$ is true.

17 Which one of the following is equal to the sum $\sum_{n=1}^{5} 2n^2$?

A. $2+8+16+32+64$ C. $2+4+6+8+10$

B. $2+4+8+16+32$ D. $2+8+18+32+50$

Join us on Telegram @qubeeacademy

CamScanner

18. A company produces shirts of black and white colors from polyester and cotton. The company has 6000 units of polyester and 3000 units of cotton in its store. A black shirt requires 5 units of polyester and 4 units of cotton where as a white shirt requires 3 units of polyester and 3 units of cotton. If the factory gets a profit of Birr 120 from each black shirt and Birr 200 from each white shirt, what is the maximum profit of the company?

A. Birr 144,000
B. Birr 120,000
C. Birr 344,000
D. Birr 200,000

19. Let S, R and T be any three sets. Which one of the following is **NOT** true about the union of the sets?

A. $(S \cup R) \cup T = T \cup (R \cup S)$
B. $S \cup S = S$
C. $S \cup R = R \cup S$
D. $S \cup T = T$

20. Which one of the following is equal to the polynomial function $f(x) = x^2 + 4x + 4$?

A. $k(x) = (x-2)(x+2)$
B. $\ell(x) = (x^2 + 10x + 3) + (-6x + 1)$
C. $g(x) = \dfrac{x^4 + 4x^2 + 4}{x^2 + 2}$
D. $h(x) = (x^2 + 6x + 8) - (2x - 4)$

Join us on Telegram @qubeacademy

CamScanner

21. Which one of the following is true about the function $f(x) = \tan x$?

A. f is a periodic function with period $\frac{\pi}{2}$.

B. The graph of f intersects y-axis at $\left(\frac{\pi}{2}, 0\right)$.

C. The domain of f is $R \setminus \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.

D. The range of f is the set of real numbers, R.

22. Which one of the following is an equation of the parabola with vertex V(0, 0) and focus F(2, 0)?

A. $x^2 = 8y$

B. $x^2 = \frac{1}{8}y$

C. $y^2 = 8x$

D. $y^2 = \frac{1}{8}x$

23. Which one of the following is equal to the sum $\sum_{k=1}^{7} 729(3^{1-k})$?

A. 1088

B. 1079

C. 1093

D. 1092

24. What is the maximum value of $z = 2x + 3y$ subjected to the constraints

$$\begin{cases} x + y \le 6 \\ 2x - y \le 6 \\ x \ge 0 \\ y \ge 0 \end{cases}$$?

A. 18

B. 22

C. 12

D. 14


Join us on Telegram @qubeeacademy

CamScanner

25. Which one of the following numbers is a common multiple of 324 and 1152?

A. 13824 C. 5184
B. 20736 D. 7776

26. Consider a circle with center O and the line segments $\overline{AB}$, $\overline{AC}$ and $\overline{BC}$ are chords of the circle as shown in the figure below. If $m(\angle ABC) = 60°$ and $m(\angle ACB) = 40°$, then what is $m(\angle BOC)$?



A. 120° C. 100°
B. 160° D. 80°

27. Which one of the following is the solution of the equation $(243)^{2x-3} = \frac{(81)^{-3x+2}}{3}$ in the set of real numbers?

A. 2 C. −1
B. −2 D. 1

28. What is the total surface area of a regular square pyramid with slant height 8cm and base area 25 cm^2?

A. 80 cm^2 C. 105 cm^2
B. 65 cm^2 D. 90 cm^2

Join us on Telegram @qubeeacademy

CamScanner

Subject Code: 02 **Booklet Code: 110**

29. The following table presents the scores of Grade 11 students of a certain school in a Mathematics Examination.

Score	31-40	41-50	51-60	61-70	71-80	81-90	91-100
Number of Students	10	9	15	12	10	4	2

What is the median of the data given in the above table?

A. 58.5 C. 50.5
B. 60.5 D. 55.5

30. What is the area of the region bounded by the graph of $f(x) = x^3$ and the x-axis on the interval $[-4, 2]$?

A. 120 Square units C. 60 Square units
B. 136 Square units D. 68 Square units

31. If $a = 2 - 3\sqrt{5}$ and $b = 3 + 2\sqrt{5}$, then which one of the following is true?

A. $ab = 36$ C. $a + b = 5$
B. $11a \div b = -36 + 13\sqrt{5}$ D. $a - b = -1 - \sqrt{5}$

32. A sector of a circle of diameter 8cm is formed by a central angle of 45°. What is the area of the sector?

A. 8π cm^2 C. π cm^2
B. 2π cm^2 D. $(2\pi+8)$ cm^2

Join us on Telegram @qubeeacademy

CamScanner

33. Which one of the following intervals is the solution set of the inequality $\left|3-\frac{1}{2}x\right|<5$?

A. $(-16, 4)$
B. $(-4, 16)$
C. $(-8, 2)$
D. $(-2, 8)$

34. The altitude of a frustum of a regular square pyramid is 9 cm. What is the volume of the frustum if the side length of the upper base is 6 cm and the side length of the lower base is 10 cm?

A. $108\ cm^3$
B. $324\ cm^3$
C. $588\ cm^3$
D. $196\ cm^3$

35. Which one of the following numbers is the coefficient of x^6y^4 in the expansion of $(x+y)^{10}$?

A. 4860
B. 1260
C. 210
D. 240

36. Which one of the following is true about the Range and Interquartile Range of a given data?

A. Both are considering only the middle 50% of the data.
B. Both are measures of variation of the data.
C. Both are affected by extreme values in the data.
D. Both are measures of the variability of each item in the data.

Join us on Telegram @qubeeacademy

CamScanner

Subject Code: 02 **Booklet Code: 110**

37. Which one of the following is equal to the expression $\frac{(125)^{\frac{2}{3}}(81)^{-\frac{3}{4}}}{(27)^{-\frac{2}{3}}(625)^{\frac{3}{4}}}$?

A. $\frac{1}{15}$ C. $\frac{3}{5}$


B. 15 D. $\frac{5}{3}$

38. If the length of the sides of a triangle are 8cm, 6cm and 4 cm , then which one of the following is the area of the triangle?

A. $3\sqrt{15}\,cm^2$ C. $24\ cm^2$

B. $1.5\sqrt{15}\,cm^2$ D. $12\ cm^2$

39. The following figure is the graph of a quadratic function $f(x) = ax^2 + bx + c$ in the set of real numbers.



Which one of the following is the solution set of the inequality $f(x) \leq 0$?

A. $\{x : x \leq 0 \text{ and } x \geq -4\}$ C. $\{x : x \leq 2 \text{ or } x \geq 6\}$

B. $\{x : x \leq -4 \text{ or } x \geq 0\}$ D. $\{x : x \geq 2 \text{ and } x \leq 6\}$

Join us on Telegram @qubeeacademy

CamScanner

40. Which one of the following is an onto function from R onto R?

A. $f(x)=2x-3x^2$

B. $f(x)=5x^3+1$

C. $f(x)=|x-1|$

D. $f(x)=\sqrt{x^2+1}$

41. Let M be a matrix such that $M=\begin{pmatrix}9 & 24 & 6\\ -12 & 18 & 15\\ 0 & 27 & 3\end{pmatrix}$. Which one of the following matrices is equal to $\frac{2}{3}M$?

A. $\begin{pmatrix}6 & 16 & 2\\ -8 & 12 & 10\\ 0 & 18 & 2\end{pmatrix}$

B. $\begin{pmatrix}6 & 16 & 4\\ -8 & 12 & 10\\ 0 & 18 & -2\end{pmatrix}$

C. $\begin{pmatrix}3 & 8 & 2\\ -4 & 6 & 5\\ 0 & 9 & 1\end{pmatrix}$

D. $\begin{pmatrix}6 & 8 & 4\\ -8 & 6 & 10\\ 0 & 9 & 2\end{pmatrix}$

42. Consider that a data set given below:

10, 8, 10, 2, 12, 6, 12, 18, 3, 7, 10, 8, 8

Which one of the following is the coefficient of range of the data?

A. $\frac{5}{4}$

B. 20

C. 16

D. $\frac{4}{5}$

43. Let r_1 and r_2 be the roots of a quadratic equation $ax^2+bx+c=0$, such that $r_1+r_2=-2.5$ and $r_1r_2=1.5$, for a, b and c are real numbers and $a\neq 0$. Which one of the following can be the values of a, b and c respectively?

A. 1, 6 and 5

B. 2, 3 and 5

C. 1, 5, and 6

D. 2, 5 and 3

Join us on Telegram @qubeeacademy

CamScanner

Subject Code: 02 **Booklet Code: 110**

44. The following data gives the scores of 10 students in a Mathematics examination: 6, 10, 8, 5, 9, 8, 5, 10, 10, 9. Which one of the following is the standard deviation of the given data?

A. 3.6
B. $\sqrt{3.6}$
C. 5.8
D. $\sqrt{5.8}$

45. Which of the following pairs of lines with the given equations are parallel?

A. $\ell_1 : x - y + 3 = 0$ and $\ell_2 : -x + y - 9 = 0$
B. $\ell_1 : 4x - y = 2$ and $\ell_2 : -x + 4y - 3 = 0$
C. $\ell_1 : 2x + 3y = 5$ and $\ell_2 : 3x + 2y = 4$
D. $\ell_1 : x - 2y + 1 = 0$ and $\ell_2 : x + 2y - 1 = 0$

46. Which one of the following is the range of the function $f(x) = \frac{2x-1}{x+3}$?

A. $R \setminus \{-3\}$
B. $R \setminus \{2\}$
C. $R \setminus \left\{-\frac{1}{3}\right\}$
D. $R \setminus \left\{\frac{1}{2}\right\}$

47. Let $z = -a + 2i$ and $w = 5 - bi$ be two complex numbers, where a and b a real numbers. If $z = w$, then which one of the following pairs of numbers a equal to a and b respectively?

A. 5 and −2
B. −5 and −2
C. 5 and 2
D. −5 and 2

Join us on Telegram @qubeeacademy

CamScanner

48. The following table presents the number of traffic accidents in a certain city per year for five consecutive years from 2019 to 2023.

Year	2019	2020	2021	2022	2023
Number of Accidents	3500	2750	2850	3500	3750

What is the average number of accidents per year for the given five years?

A. 3250
B. 3270
C. 3200
D. 3210

49. Which one of the following is the solution set of the equation $(\sqrt{16})^{2x+1} = (\sqrt[3]{4096})^{3x-5}$?

A. $\left\{\frac{9}{4}\right\}$
B. $\left\{\frac{9}{2}\right\}$
C. $\left\{\frac{11}{2}\right\}$
D. $\left\{\frac{11}{4}\right\}$

50. Which one of the following is a primary data?

A. The data obtained from a chart prepared by Ministry of Health regarding the spread of HIV/AIDS.
B. The data obtained by reading a journal regarding the opinion of the society about holidays.
C. The data obtained by measuring the weights of students in a certain school.
D. The data obtained from a document prepared by the MOE regarding the achievements of students in National Examination.

Join us on Telegram @qubeeacademy

CamScanner

51. Which one of the following ordered pairs of numbers gives the coordinates of the point that divides a line segment with end points A(1, 1) and B(−6, 8) in the ratio of 3:4?

A. (2, −4)
B. (−2, −4)
C. (2, 4)
D. (−2, 4)

52. Which one of the following pairs of functions is an inverse of each other in their respective domains?

A. $f(x)=\sqrt{x+3}$ and $g(x)=x^2-9$

B. $f(x)=x^5$ and $g(x)=\frac{1}{x^5}$

C. $f(x)=x^3-1$ and $g(x)=\sqrt[3]{x+1}$

D. $f(x)=\frac{x+1}{x-1}$ and $g(x)=\frac{x-1}{x+1}$

53. Which one of the following numbers is the determinant of the matrix $M=\begin{pmatrix}1 & 2 & 3\\ 4 & 5 & 6\\ 7 & 8 & 10\end{pmatrix}$?

A. −3
B. 6
C. −6
D. 3

54. Which one of the following is a simple random sampling?

A. First the population is divided into homogenous groups and a sample is selected from each homogenous group.
B. First divide a city into ten groups and choose the secondary school principals from the randomly selected 3 groups.
C. Selecting five students from grade 12 students of the same school randomly.
D. Selecting a group randomly from groups formed by dividing the population into groups.

Join us on Telegram @qubeeacademy

CamScanner

55. Let A and B be two sets that contain 15 and 17 elements respectively. If $A \cap B$ contains 10 elements, what is the number of elements in $A \cup B$?

A. 22
B. 12
C. 42
D. 32

56. Which one of the following is the position vector of the vector with initial point A (1, −1) and terminal point B (2, −3)?

A. $3i+4j$
B. $i-4j$
C. $3i-4j$.
D. $i-2j$

57. Let θ be an angle in standard position and P(−3, 4) be a point on the terminal side of θ. Which one of the following is true about the corresponding trigonometric functions?

A. $\cos\theta = \frac{-3}{5}$
B. $\sin\theta = \frac{-4}{5}$
C. $\sin\theta = \frac{3}{5}$
D. $\cos\theta = \frac{4}{5}$

58. Which one of the following conic section is defined by "the set of points in a plane such that the sum of the distances from two given points is constant"?

A. Ellipse
B. Hyperbola
C. Parabola
D. Circle


59. If the fifth and the tenth terms of a geometric sequence are 640 and 20 respectively, then which of the following numbers are the first term and the common ratio of the given geometric sequence respectively?

A. 1024 and 2
B. 1024 and $\frac{1}{2}$
C. 10240 and 2
D. 10240 and $\frac{1}{2}$

Join us on Telegram @qubeeacademy

CamScanner

60. Given that a graph of system of linear inequalities as shown below.



Which one of the following system of linear inequalities is represented by the shaded region in the graph?

A. $\begin{cases} 2x+y \leq 4 \\ 3x-y \leq 6 \end{cases}$

B. $\begin{cases} 2x+y \geq 4 \\ 3x-y \leq 6 \end{cases}$

C. $\begin{cases} 2x+y \geq 4 \\ 3x-y \geq 6 \end{cases}$

D. $\begin{cases} 2x+y \leq 4 \\ 3x-y \geq 6 \end{cases}$

THE END

Join us on Telegram @qubeeacademy

CamScanner