NANDA improved nursing diagnosis

- Imbalanced nutrition
- Risk for imbalanced nutrition
- Risk for infection
- Risk for imbalanced body temperature
- Hypothermia
- Hyperthermia
- Ineffective thermoregulation
- Autonomic dysreflexia
- Constipation
- Perceived constipation
- Diarrhea
- Bowel incontinence
- Risk for constipation
- Impaired urinary elimination
- Stress urinary incontinence
- Reflex urinary incontinence
- Urinary retention
- Ineffective tissue perfusion (specify type renal, cerebral, cardio-pulmonary, gastrointestinal, peripheral)
- Risk for imbalanced fluid volume
- Decreased cardiac output
- Impaired gas exchange
- Ineffective breathing battern
- Risk for injury
- Risk for falls
- Risk for suffocation
- Risk for poisoning
- Risk for trouma
- Risk for aspiration
- Risk for disuse syndrome
- Latex allergy response
- Risk for latex allergy response
- Ineffective protection
- Impaired tissue integrity
- Impaired oral mucous membrane
- Impaired skin integrity
- Risk for impaired skin integrity
- Impaired dentition

- Disturbed energy field

Pattern two: communicating

Impaired verbal communication

Pattern three: relating

- impaired social interaction
- social isolation
- risk for loneliness
- impaired parenting
- risk for impaired parenting
- sexual dysfunction
- interrupted family process
- caregiver role strain
- dysfunctional family process: alcoholism
- parental role conflict
- ineffective sexuality pattern

Pattern four: evaluating

- spiritual distress
- risk for spiritual distress
- readiness for enhanced spiritual well-being

Pattern five: choosing

- ineffective coping
- impaired adjustment
- defensive denial
- defensive coping
- disabled family coping
- compromised family coping
- readiness for enhanced community coping
- ineffective community coping
- ineffective therapeutic regimen management
- noncompliance (specify)
- effective therapeutic regimen management
- decisional conflict

- health seeking behavior

Pattern six: moving

- impaired physical mopility
- risk for peripheral neurovascular dysfuncion
- risk for perioperative-positioning injury
- impaired walking
- impaired wheelchair mobility
- impaired transfer ability
- impaired bed mobility
- activity tolerance
- fatigue
- risk for activity intolerance
- disturbed sleep pattern
- sleep deprevition
- deficient diversional activity
- impaired home maintainance
- ineffective health maintainance
- delayed surgical recovery
- adult failure to thrive
- feeding self-care deficit
- impaired swallowing
- ineffective breast feeding
- interrupted breast feeding
- effective breast feeding
- ineffective infant feeding pattern
- bathing/hygiene self-care deficit
- toileting self-care deficit
- delayed growth and development
- risk for delayed development
- relocation stress syndrome
- risk for relocation stress syndrome
- risk for disorganized infant behavior
- disorganized infant behavior
- readiness for organized infant behavior