

Ethiopian University Entrance Examination (EUEE) Mathematics for
Natural Science Ginbot 2010/June 2018

BOOKLET CODE:25

SUBJECT CODE:02

Number Of Items:65

Time Allowed:3hours

DIRECTIONS: for each of the following problems. Choose the best answer from the given alternative and carefully **blacken** the letter of your best choice on the separate answer sheet provided.

1. Which one of the following is true about signum, absolute value and greatest integer functions?

- (A) $\operatorname{sgn}(x) = \pm |x|$, for all $x \in \mathbb{R}$.
 (B) $|x| = x \operatorname{sgn}(x)$, for all $x \in \mathbb{R}$.
 (C) $\operatorname{sgn}(x) < [x]$, for all $x \leq 0$.
 (D) $\operatorname{sgn}(x) \leq [x]$, for all $x \geq 0$.

$|x| = x \operatorname{sgn} x$ for all real number x $\operatorname{sgn} x \leq |x|$ for all $x \geq 0$

Ans. B and D

2. What is the equation of the line that passes through (1,1) and is parallel to line $3y - x = 1$?

- (A) $3x - y = 2$
 (B) $x + 3y = 4$
 (C) $3y - x + 2 = 0$
 (D) $x - 3y + 2 = 0$

$3y - x = 1 \Rightarrow 3y = x + 1$ equation of a line that pass through (1,1) and parallel to $9 = \frac{1}{3}x + 1$

Is $\frac{y-1}{x-1} = \frac{1}{3} \Rightarrow 3y - 3 = x - 3 + 1 = 0$
 $3y - x - 2 = 0$

3. Which one of the following is an equation of the circle whose endpoints of a diameter are (0,-2) and (2, 2)?

(A) $x^2 + y^2 - 2x - 4 = 0$

(B) $X^2 + y^2 = 4$

(C) $(x-1)^2 + y^2 = 4$

(D) $x^2 + y^2 - 2y - 4 = 0$

The end points of diameter are (0,-2) and (2,-2) by mid point formula center is (1,0) and radius is $\sqrt{5}$
 $\Rightarrow$ Equation of circle is $x^2 + y^2 - 2x - 4 = 0$

Ans. A

4. What are the greatest lower bound and least upper bound of the sequence $\{(-1)^n(1 + \frac{1}{n})\}$, respectively?

(A) -2 and 2

(B) -2 and $\frac{3}{2}$

(C) $-\frac{3}{2}$ and 2

(D) -2 and $-\frac{3}{2}$

$$\text{if } x = 1 \left\{(-1)^n(1 + \frac{1}{n})\right\} \text{ is } -2$$

$$\text{if } x = 2, \left\{(-1)^n(1 + \frac{1}{n})\right\} \text{ is } \frac{3}{2}$$

$$\text{if } x = 3, \left\{(-1)^n(1 + \frac{1}{n})\right\} \text{ is } -\frac{4}{3}$$

$$\text{if } x = 4 \left\{(-1)^n(1 + \frac{1}{n})\right\} \text{ is } \frac{5}{4}$$

$\Rightarrow$ GLb is -2 and L.U.b is $\frac{3}{2}$

Ans. B

5. What is the maximum value of the function $f(x) = x^4 - 2x^2$ on $[-2, 1]$?

(A) 8

(B) 12

(C) 24

(D) 40

$$\begin{aligned}
 f(x) &= x^4 - 2x^2 \\
 f'(x) &= 4x^3 - 4x \\
 f'(x) = 0 &\Leftrightarrow 4x^3 - 4x = 0 \\
 4x(x^2 - 1) &= 0 \\
 4x = 0 \text{ or } x &= \pm 1
 \end{aligned}$$

Critical numbers are -2,0,1,-1

$$f(-2) = 8 \quad f(1) = -1$$

$$f(0) = 0 \quad f(-1) = 1$$

8 is maximum value at (2)

Ans. A

6. If the truth value of $(p \wedge \neg p) \Leftrightarrow [(q \vee \neg q) \Rightarrow r]$ is True, then which one of the following must be True?

(A) p

(B) q

(C) $\neg q$

(D) $\neg r$

$$(p \wedge \neg p) \Leftrightarrow [(q \vee \neg q) \Rightarrow r] \text{ is true}$$

$$\Leftrightarrow p \wedge \neg p \text{ is false since } p \text{ or } \neg p \text{ is true}$$

$$\Leftrightarrow (q \vee \neg q) \text{ is true since } (p \wedge \neg p) \text{ is false and by property of implication}$$

$$\Leftrightarrow (q \vee \neg q) \text{ is true since } q \text{ or } \neg q \text{ is true}$$

$$\Leftrightarrow \neg r \text{ must be true to make the proposition true}$$

Ans. D

7. Let A and B be two events. Suppose that the probability that neither event occur is $3/8$. What is the probability that at least one of the events occur?

(A) $5/8$

(B) $1/4$

(C) $1/8$

(D) $3/4$

$$P(A \cup B)' = \frac{3}{8}$$

$$\begin{aligned} P(A \cup B) &= 1 - p(A \cup B)' \\ &= 1 - \frac{3}{8} \\ &= \frac{5}{8} \end{aligned}$$

Ans. A

8. What is the value of $\sum_{i=0}^n \left(\frac{1}{n-1} - \frac{1}{n} \right)$?

(A) $\frac{17}{20}$

(B) $\frac{21}{20}$

(C) $\frac{19}{20}$

(D) $\frac{23}{20}$

$$\begin{aligned} \sum_{n=1}^{20} \left(\frac{1}{n-1} - \frac{1}{n} \right) &= \left\{ \left(1 - \frac{1}{2} \right) + \left(\frac{1}{2} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{4} \right) + \dots + 19 - \frac{1}{20} \right\} \\ &= \left\{ 1 - \cancel{\frac{1}{2}} + \cancel{\frac{1}{2}} - \cancel{\frac{1}{3}} + \cancel{\frac{1}{3}} - \cancel{\frac{1}{4}} + \cancel{\frac{1}{4}} - \dots - \frac{1}{20} \right\} \\ &= \left\{ 1 - \frac{1}{20} \right\} = \frac{19}{20} \end{aligned}$$

Ans. C

9. Which one of the following is a convergent sequence?

(A) $\left\{ \frac{(-1)^n}{2} \right\}$

(B) $\left\{ \frac{1}{n} + \sin(n) \right\}$

(C) $\left\{ \frac{1-3^n}{2^n} \right\}$

(D) $\left\{ \frac{1-2^n}{3^n} \right\}$

$$\begin{aligned} \text{if } n = 1, \left\{ \frac{1+2^n}{3^n} \right\} \text{ is } 1 \\ \text{Take } \left\{ \frac{1+2^n}{3^n} \right\} \text{ if } n = 2, \left\{ \frac{1+2^n}{3^n} \right\} \text{ is } \frac{5}{9} \\ \text{if } n = 3, \left\{ \frac{1+2^n}{3^n} \right\} \text{ is } \frac{9}{27} \\ \text{if } n = 4, \left\{ \frac{1+2^n}{3^n} \right\} \text{ is } \frac{17}{18} \end{aligned}$$

If it converges to zero the others are divergent.

Ans. D

10. If a function f is differentiable at a , then what is the value of

$$\lim_{x \rightarrow a} f(x)$$

(A) $f'(a)$

(B) $f(a)$

(C) $f'(a) - f(a)$

(D) 0

To differentiate the function at a means find $f'(a)$. the derivative of $f(x)$ at a is also expressed as

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

Ans. A

11. If $f'(x) = e^{x-1} + 3x^2 - \frac{1}{x}$ and $f(1)=5$, what is $f(x)$?

(A) $f(x) = e^{x-1} + x^2 - \ln x + 3$

(B) $f(x) = e^{x-1} - x^3 - \ln x + 5$

(C) $f(x) = e^{x-1} + 3x^2 - \frac{1}{x} + 5$

$$f(x) = e^{x-1} + 3x^2 - \frac{1}{x}, f(1) = 5$$

$$\int f'(x) = f(x) = e^{x-1} + x^3 - \ln x + c$$

$$f(1)=5 \Rightarrow e^x + x^3 - \ln x + c = 5$$

$$\Rightarrow 1+1-0+C=5$$

$$\Rightarrow C = \underline{3}$$

$\therefore$ the equation of $f(x)$ is $e^x + x^3 - \ln x + 3$

Ans. A

12.If A is a square matrix of order 3 and $\det(A)=5$, then *what is the value of $\det(A.\text{adj}(A))$?*

- (A) 3
- (B) 5+
- (C) 25
- (D) 125

A is square matrix with order 3×3
 $\det(A \text{ and } A) = \det(A)(A)(\det \text{adj } A)^{n-1}$
 $= \det(A)(A.\det(\text{adj } A)^{3-1})$
 $= \det(A)(A.\det(\text{adj}(A))^2)$
 $= 5 \times 5^2 = \underline{125}$

Ans. D

13. A salesman sold items x_1, x_2 and x_3 , with different rates of commission as shown in the table below.

Month	Sales of units			Total commission (in Birr)
	x_1	x_2	x_3	
February	90	100	20	800
March	130	50	40	900
April	60	50	30	850

What are the rates of commission on items x_1, x_2 and x_3 , respectively.

- (A) 4,2 and 11
- (B) 4,11 and 2
- (C) 2,4 and 11
- (D) 11,2 and 4

$$\begin{aligned} 90c + 100x_2 + 20x_3 &= 800 \\ \text{Soln from the above table } \int 130x_1 + 50x_2 + 40x_3 &= 90 \\ 60x_1 + 100x_2 + 30x_3 &= 850 \end{aligned}$$

$$\text{Solve by Cramer's rule } D = \begin{vmatrix} 9 & 10 & 2 \\ 13 & 5 & 4 \\ 6 & 10 & 3 \end{vmatrix} = -175 \quad D_x = \begin{vmatrix} 80 & 10 & 2 \\ 90 & 5 & 4 \\ 85 & 10 & 3 \end{vmatrix} = -350 \quad D_y = \begin{vmatrix} 9 & 80 & 2 \\ 13 & 90 & 4 \\ 6 & 85 & 3 \end{vmatrix} = -700$$

$$D_z = \begin{vmatrix} 9 & 10 & 80 \\ 13 & 5 & 90 \\ 6 & 10 & 80 \end{vmatrix} = -1925$$

$$\Rightarrow x = \frac{D_x}{D} = \frac{-350}{-175} = 2$$

$$y = \frac{D_y}{D} = \frac{-700}{-175} = 4$$

$$z = \frac{D_z}{D} = \frac{-1925}{-175} = 11$$

The rate of commission x_1 , x_2 and $x_3 = x, y$ and $z = 2, 4$ and 11 respectively

Ans. C

14. If $z = -3 + 4i$ and $w = 1 + 2i$, then what is the value of $\frac{2z}{w} + \bar{w}$?

(A) $2+3i$

(B) $3+2i$

(C) $3+5i$

(D) $3-2i$

$$z = -3+4i \quad w = 1+2i \text{ then}$$

$$\frac{2z}{w} + \bar{w} = \frac{2(-3+4i)}{1+2i} + 1-2i$$

$$= \frac{-6+8i}{1+2i} \times \frac{1-2i}{1-2i} + 1-2i$$

$$= \frac{-6+12i+8i+16}{1+4} + 1-2i$$

$$= \frac{10+20i}{5} + 1-2i$$

$$= 2+4i+1-2i$$

$$= \underline{\underline{3+2i}}$$

Ans. B

15. The population of a certain city is increasing at a rate of 3% per year. If the population was 100,000 in 2012 E.C., then what will be the population in 2020 E.C.?

(Given: $(1.03)^9 = 1.30$, $(1.03)^{10} = 1.34$, $(1.3)^9 = 10.60$, $(1.3)^{10} = 13.78$)

- (A) 130,000
- (B) 134,000
- (C) 1,060,000
- (D) 1,378,000

Given $P = 100,000$

$R = 0.03$

$T = 10$ year

$$\begin{aligned} A &= P(1+r)^t \Rightarrow A = 100,000 (1+0.03)^{10} \\ &= 100,000 (1.03)^{10} \\ &= 134,000 \end{aligned}$$

Ans. B

16. What is the slope of the tangent line to the graph of $f(x) = 3e^x + \sin x + 2$ at $(0, 5)$?

- (A) 2
- (B) 3
- (C) 4
- (D) 5

$F(x) = 3e^x + \sin x + 2$ at $(0, 5)$

$F'(x) = 3e^x + \cos x$

$F'(0) = \text{slope of tangent line} = 3e^0 + \cos 0$
 $= 3 + 1 = \underline{4}$

Ans. C

17. What is the value of $\int_1^2 \frac{x+4}{x(x+2)} dx$

- (A) 2
- (B) $\ln 2$
- (C) $\ln 3$
- (D) $\ln 4 - \ln 3$

$\int \frac{x+4}{x(x+2)} dx$ by partial fraction

$$A + B = 1$$

$$2 + B = 1$$

$$B = -1$$

$$\frac{x+4}{x(x+2)} = \frac{A}{x} + \frac{B}{x+2}$$

$$\frac{x+4}{x(x+2)} = \frac{A(x+2)+Bx}{x(x+2)}$$

$$\frac{A}{x} + \frac{B}{x+2} = \frac{2}{x} - \frac{1}{x+2}$$

$$x+4 = Ax+2A+Bx$$

$$x+4 = (A+B)x + 2A$$

$$\Rightarrow A+B = 1$$

$$2A = 4 \Rightarrow A = 2$$

$$\Rightarrow 2(1) - 1(4) - (0 - (1)(3))$$

$$= 1(3)$$

$$\int \frac{2}{x} - \frac{1}{x+2} dx = 2 \ln|x| - \ln|x+2| + C$$

Ans. C

18. What is the area of the region enclosed by the graph of $x^2 = x + 1$ and $y^2 = -x + 1$?


(A) $\frac{8}{3}$ sq. units

(B) $\frac{4}{3}$ sq. units

(C) $\frac{3}{8}$ sq. units

(D) $\frac{3}{4}$ sq. units

The area of a region enclosed by the graphs of $y^2 = x+1$ and $y^2 = -x+1$



The region is symmetric with respect to y-axis and we can find the area in terms of y

$$A = 2 \int_0^1 (-y^2+1) - (y^2-1) dy$$

$$A = 2 \int_0^1 (-y^2+1-(y^2-1)) dy$$

$$A = 2 \int_0^1 (-2y^2+2) dy$$

$$= 2 \left(-\frac{2y^3}{3} + 2y \right) \Big|_0^1$$

$$= 2 \left(-\frac{2}{3} + 2 \right)$$

$$A = \frac{8}{3} \text{ sq. Unit}$$

Ans

19. What is the partial fraction decomposition of $\frac{x^2 + x + 1}{(x+2)(x^2+1)}$?

(A) $\frac{5}{3(x+2)} + \frac{2x+1}{3(x^2+2)}$

(B) $\frac{3}{5(x+2)} + \frac{2x+1}{5(x^2+1)}$

(C) $\frac{2}{5(x+2)} + \frac{3x+1}{5(x^2+1)}$

(D) $\frac{2}{3(x+2)} + \frac{2x+1}{3(x^2+1)}$

Partial fraction of $\frac{x^2 + x + 1}{(x+2)(x^2+1)}$ let $\frac{A}{x+2} + \frac{Bx+C}{x^2+1}$

$$\frac{x^2 + x + 1}{(x+2)(x^2+1)} + \frac{A(x^2+1) + (Bx+C)(x+2)}{(x+2)(x^2+1)}$$

$$x^2 + x + 1 = Ax^2 + A + Bx^2 + 2Bx + Cx + 2C$$

$$x^2 + x + 1 = (A+B)x^2 + (2B+C)x + A + 2C$$

$$A + B = 1 \quad \text{if } x = -2$$

$$2B + C = 1 \quad 4 - 2 + 1 = 5A$$

$$A + 2C = 2 \quad 3 = 5A$$

$$A + B = 1$$

$$\frac{3}{5} + B = 1$$

$$B = 1 - \frac{3}{5}$$

$$B = \frac{2}{5}$$

$$2B + C = 1$$

$$\frac{4}{5} + C = 1$$

$$C = \frac{1}{5}$$

$$\begin{aligned} \frac{A}{x+2} + \frac{Bx+C}{x^2+1} &= \frac{3}{5(x+2)} + \frac{\frac{2}{5}x + \frac{1}{5}}{x^2+1} \\ &= \frac{3}{5(x+2)} + \frac{2x+1}{5(x^2+1)} \end{aligned}$$

Ans. B

20. What is the area of the triangle (in sq. units) formed by the lines joining the vertex of the parabola $x^2 = 36y$ to the end points of the latus rectum?

(A) 126

(B) 162


(C) 216

(D) 261

$$x^2 = -369$$

$$-4P = -36$$

$$P = 9$$



It is isosceles triangle with base = length of latus rectum of parabola. As shown in the figure
length of latus rectum (base)

$$= |4P| = |-36| = 36$$

$$A = \frac{1}{2} b, h$$

$$A = \frac{1}{2} \times 36 \times 9 \text{ (where } h = p \text{)}$$

$$A = 162 \text{ sq unit}$$

Ans. B

21. The time needed to type a sample of 8 business letters in an office is 7, 8, 6, 8, 9, 7, 5, 6 minutes. What are the mean ($\bar{x}$) and the standard deviation (s) of the data in minute?

(A) $\bar{x} = 7, s = \sqrt{2}$

(B) $\bar{x} = 7, s = \sqrt{1.5}$

(C) $\bar{x} = 8, s = \sqrt{2}$

(D) $\bar{x} = 8, s = \sqrt{1.5}$

The data is 7, 8, 6, 8, 9, 7, 5, 6,

$$\bar{x} = \frac{7+8+6+8+9+7+5+6}{8} = \frac{56}{8} = 7$$

$$s^2 = \frac{(7-7)^2 + (8-7)^2 + (6-7)^2 + (8-7)^2 + (9-7)^2 + (7-7)^2 + (5-7)^2 + (6-7)^2}{8}$$
$$= \frac{1+1+1+1+4+1+1+1}{8} = \frac{8}{8} = 1$$

$$s = \sqrt{1} = 1$$

Ans. B

22. Let A be a 3×3 invertible matrix and B be any 3×3 matrix. If $|A| = a$ and, $|B| = b$, then which one of the following is **NOT** true?

- (A) $|A^T A| = a^2$
- (B) $|A^T A| = a^2$
- (C) $|kA| = k^2 |A|$, for any $k \in \mathbb{R}$
- (D) $|A^{-1} B| = ab$

Two matrices A and B are 3×3 matrices then $|A^T A| = |A^T| |A| = a \cdot a = a^2$. $|KA| = K^2 |A|$ since if is 3×3 matrix but $|\bar{A}B| \neq ab$

Ans. D

23. Let $z = \left(\frac{1-i}{1+i}\right)^{18}$. Then what is the value of ?

- (A) i
- (B) -1
- (C) $-i$
- (D) $1-i$

$$z = \left(\frac{1-i}{1+i}\right)^{18} \Rightarrow z = \left(\frac{1-i}{1+i}\right)^{2 \cdot 9}$$

$$\Rightarrow z = \left(\frac{-2i}{2i}\right)^9 = 1^9 = \underline{-1}$$

Ans. B

24. If $f(x) = \frac{|x|}{x}$ and $g(x) = \frac{x+2}{x^3-4x}$, then what is the value of

$\lim_{x \rightarrow -2} (f(x) + g(x))$?

- (A) $\frac{-9}{8}$
- (B) $\frac{-7}{8}$
- (C) $\frac{9}{8}$
- (D) ∞

$$F(x) = \frac{|x|}{x} \text{ and } g(x) = \frac{x+2}{x^3-4x}$$

$$\begin{aligned}\Rightarrow \lim (f(x) + g(x)) &= \lim_{x \rightarrow -2} f(x) + \lim_{x \rightarrow -2} \frac{x+2}{x^3-4x} \\ &= \lim_{x \rightarrow -2} \frac{|x|}{x} + \lim_{x \rightarrow -2} \frac{x+2}{x^3-4x} \\ &= -1 + \lim_{x \rightarrow -2} \frac{x \neq 2}{x(x \neq 2)(x-2)} \\ &= -1 + \lim_{x \rightarrow -2} \frac{1}{x(x-2)} \\ &= -1 + \frac{1}{8} = \frac{-7}{8}\end{aligned}$$

Ans. B

25. What is the value of $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin^2\left(\frac{x}{2}\right)$?

(A) $\frac{1}{4}$

(B) $\frac{1}{2}$

(C) 2

(D) 4

(E)

$$\lim_{x \rightarrow 0} \frac{1}{x^2} \sin^2\left(\frac{x}{2}\right) = \lim_{x \rightarrow 0} \frac{\sin^2 \frac{x}{2}}{x^2}$$

$$= \lim_{x \rightarrow 0} \frac{\sin^2 \frac{x}{2} \cdot \frac{1}{x^2}}{\frac{1}{x^2}}$$

$$= \frac{1}{4} \lim_{x \rightarrow 0} \frac{\sin^2 \frac{(x)^2}{2}}{\frac{(x)^2}{2}} = \frac{1}{4} \cdot 1 = \frac{1}{4}$$

Ans. A

26. Which one of the following is NOT true about the function $f(x) = 3x^4 - 4x^3$?

(A) (0,0) is point of inflection of f .

(B) 0 and 1 are critical number of f

(C) f is decreasing on $(-\infty, 1)$ and increasing on $(1, \infty)$

(D) f is concave up on $(-\infty, 0)$ and concave down on $(0, \infty)$

$$F(x) = 3x^4 - 4x^3$$

$$F'(x) = 12x^3 - 12x^2 \Rightarrow f'(x) = 0 \Rightarrow 12x(x^2 - 1) = 0$$

$$F''(x) = 36x^2 - 24x$$

$x=0, x=-1, x=1$ are critical numbers

$$F''(x) = 0 \Rightarrow 36x^2 - 24x = 0$$

$$6x(6x-4) = 0$$

$x=0, x=\frac{2}{3}$ are inflection points is true the others are not true

Ans. A

27. What is the value of $\int x \sqrt{1-x^2} dx$?

(A) $\frac{1}{2} \sqrt{1-x^2} + C$

(B) $\frac{-1}{2} (1-x^2)^{3/2} + C$

(C) $(1-x^2)^{3/2} + C$

(D) $-2x\sqrt{1-x^2} + C$

$$\begin{aligned} & \int x \sqrt{1-x^2} \\ \Rightarrow & \int x \sqrt{u} \cdot \frac{du}{-2x} \\ \Rightarrow & -\frac{1}{2} \cdot \frac{u^{3/2}}{3/2} + c \end{aligned}$$

$$\begin{aligned} \Rightarrow & -\frac{1}{2} \int u^{1/2} du \quad \Rightarrow \frac{1}{2} \cdot \frac{u^{3/2}}{3/2} + c \\ & = -\frac{1}{3} (1-x^2)^{3/2} + c \end{aligned}$$

$$\begin{aligned} \text{let } u &= 1-x^2 \\ du &= -2x dx \\ dx &= \frac{du}{-2x} \end{aligned}$$

Ans. B

28. What is the value of $\int_1^9 e^{\frac{\sqrt{x}}{\sqrt{x}}} dx$?

(A) $e^3 - e$

(B) $\frac{e^3}{3} - e$

(C) $2(e^3 - e)$

(D) $e(e^2 - \frac{1}{3})$

$$\int_1^9 \frac{e^{\sqrt{x}}}{\sqrt{x}} dx = \quad \text{Let } u = \sqrt{x} \quad \begin{aligned} du &= \frac{1}{2\sqrt{x}} dx \\ dx &= 2\sqrt{x} du \end{aligned}$$

$$\int_1^9 \frac{e^u}{u} 2u du = \int_1^9 2e^u du$$

$$= 2(e^u) \Big|_1^9$$

$$= 2(e^{\sqrt{x}}) \Big|_1^9$$


$$= 2(\underline{e^3 - e})$$

Ans. C

29. A running a race- course noted that the sum of the distance from the two flag posts form him is always 10 meters and the distance between the flag posts is 8 meters. What is the equation of the path traced by the man?

- (A) $\frac{x^2}{9} + \frac{y^2}{25} = 1$
- (B) $\frac{x^2}{64} + \frac{y^2}{100} = 1$
- (C) $\frac{x^2}{100} + \frac{y^2}{64} = 1$
- (D) $\frac{x^2}{25} + \frac{y^2}{9} = 1$

the man running in shep of ellipse. the sum distance form the two flags means $2a = 10, a = 5$



the distance $\frac{b}{n}$ flages means $\frac{a}{c} \frac{b}{n}$ foci = $\frac{2c}{C} = \frac{8}{4}$

$$b^2 = a^2 - c^2$$

$$b^2 = 25 - 9 \Rightarrow b = 3$$

The equation path is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$\Rightarrow \frac{x^2}{25} + \frac{y^2}{9} = 1$$

Ans. D

30. Suppose the following are the premises of an argument.

He is healthy and he is not angry.

He is angry or his plan fails.

His plan does not fail if he does not travel abroad. Type equation here.

Given that an the premises are true, which one of the following be conclusion that makes the argument valid?

- (A) He travels abroad.
- (B) His plan fails and he is angry.
- (C) His plan does not fail.
- (D) His plan fails and he does not travel abroad.

Let p : he is healthy

q : He is angry

r : His plan fails

S : He travel abroad

The premises are $P \wedge q, q \vee r, \neg S \Leftrightarrow \neg r$ the conclusion that the argument true is he travel abroad (r)

Ans. B

31. The age distribution of students in a certain class is given below:

Age	10-14	15-19	20-24	25-29
No. of students	2	10	6	7

What is the modal value of the distribution?

- (A) 17.38
- (B) 17.83
- (C) 18.37
- (D) 18.73

Age	10-14	15-19	20-24	25-29
No stu	2	10	6	7

The medal class is the largest number (class 2)

$$\begin{aligned}
 mo &= LB + W \left(\frac{d1}{d1 + d2} \right) & \text{where } d1 &= 10 - 2 = 8 \\
 & & d2 &= 10 - 6 = 4 \\
 & & w &= 5 \\
 &= 14.5 + 5 \left(\frac{8}{8 + 4} \right) \\
 &= 14.5 + 3.33 \\
 mo &= 17.83
 \end{aligned}$$

Ans. B

32. Let $A = \{1,2,3,4,5,6,7\}$, $B = \{8,9,10\}$. If one of the numbers is deleted randomly from each of these sets, what is the probability that all the three deleted numbers are even or all are multiples of 3?

- (A) $\frac{1}{9}$
 (B) $\frac{2}{21}$
 (C) $\frac{8}{63}$
 (D) $\frac{4}{5}$

Son let $P(E_1)$ - even and $p(E_2)$ - multiple of 3

In Set A

$$P(E_1) = \frac{3}{7}$$

$$P(E_2) = \frac{2}{3}$$

$$P(E_1 n E_2) = 0$$

$$P(E_1 n E_1 n E_1) = \frac{3}{7} \times \frac{1}{3} \times \frac{2}{3} = \frac{2}{21}$$

$$P(E_2 n E_2 n E_2) = \frac{2}{3} \times \frac{1}{3} \times \frac{1}{3} = \frac{2}{27}$$

$$P(E_3 n E_3 n E_3) = \frac{1}{7} \times 0 \times 0 = 0$$

$$P(\text{even or multiple of 3}) = P(\text{even}) + p(\text{multi 3})$$

$$\frac{2}{7} + \frac{2}{27} = \frac{8}{63}$$

Set B

$$P(E_1) = \frac{1}{3}$$

$$P(E_2) = \frac{1}{3}$$

$$P(E_1 n E_2) = 0$$

In set C

$$P(E_1) = \frac{2}{3}$$

$$P(E_2) = \frac{1}{3}$$

$$P(E_1 n E_2) = 0$$

Ans. C

33. Let A be a 3x3 matrix and $|A| = -2$. Then what is the value of $|adj(A)|$?

- (A) 4
 (B) -2
 (C) $-1/2$
 (D) -8

Let A be 3x3 matrix and $|A| = -2$

$$|adj(A)| = |A|^{n-1}$$

$$|adj(A)| = (-2)^{n-1} \quad n = 3$$

$$= 4$$

Ans. A

34. If the sum of the three consecutive terms of an arithmetic progression $\{A_n\}$, with $A_n > 0$ for n , is 9 and the sum of their squares is 35, then what is the sum S_n of the first n terms?

- (A) $n^2 + 1$
 (B) $n^2 - 1$
 (C) $2n^2 + 1$
 (D) n^2

(An) be writhe metric sequence with $A_n > 0$ $a_1 + a_2 + a_3 = 9 \Rightarrow 5_3 = \frac{3}{2} (a_1 + a_3) = 9 \Rightarrow a_1 + a_3 = 6$
 $a_1^2 + a_2^2 + a_3^2 = 35 \Rightarrow a_1^2 + a_3^2 = 26$
 $\Rightarrow a_1 + a_3 + a_2 = 9 \Rightarrow 6 + a_2 = 9 \Rightarrow a_2 = 3 \Rightarrow (a_1 + a_3)^2 = a_1^2 + 2a_1a_3 + a_3^2 = 36$
 $a_1^2 + a_3^2 = 26 \Rightarrow 26 + 2a_1a_3 = 36 \Rightarrow 2a_1a_3 = 10$
 $\Rightarrow a_1 + 3 + \frac{5}{a_1} = 9 \Rightarrow a + \frac{5}{a} = 6 \Rightarrow a^2 - 6a + 5 = 0$
 $a_1 = \frac{6 \pm \sqrt{36 - 4 \times 5}}{2} = \frac{6 \pm 4}{2} = 5 \text{ or } 1$
 $a_3 = \frac{5}{5} = 1 \text{ or } a_3 = \frac{5}{1} = 5$
 If $a_1 = 1, d = 2$ then $a_n = a_1 + (n - 1)d$
 if $a_n = 1 + (n - 1)2 = 2n - 1$
 $5n = \frac{n}{2} (1 + 2n - 1) = n^2$
 the sum of the 1st terms is $5n = n^2$

Ans. D

35. Let $\{a_n\}$ be a sequence with $a_1 = a, a_2 = f(a_1) = f(a), a_3 = f(a_2) = f(f(a)), \dots, a_{n+1} = f(a_n)$, where f is a continuous function? If $\lim_{n \rightarrow \infty} a_n = 5$, what is $f(5)$?

- (A) 5^n
 (B) 5
 (C) 5^{n+1}
 (D) 1

Soln $\{a_n\}$ be sequence with
 $a_1 = a, a_2 = f(a_1) = f(a)$
 $a_3 = f(a_2) = f(f(a))$
 $a_{n+1} = f(a_n)$ where f is continuous function
 $\lim_{n \rightarrow \infty} a_n = 5, f(5) = \underline{5}$

Ans. B

36. If $f(x) = \frac{1}{3}x^3 + cx^2 + ax + 5$ has a local minimum value at $x=1$, then which one of the following is true about the possible value of a and c ?

(A) $a = 3, c = -2$

(B) $a = -2c - 1, c < -1$

(C) $a = -2c - 1, c > -1$

(D) $a = -2c - 1, c$ any real number.

$$F(x) = \frac{1}{3}x^3 + Cx^2 + ax + 5$$

$$F'(x) = x^2 + 2cx + a$$

$$F'(1) = 0 \Rightarrow 1 + 2c + a = 0$$

$$A = -2c - 1$$

If $c > -1$, $f(x)$ has minimum value at $x = 1$

Ans. C

37. What is the value of $\int \frac{1}{x}(\ln x + x^2 e^{-x}) dx$?

(A) $\frac{1}{2} \ln^2 x - (x + 1)e^{-x} + C$

(B) $\frac{1}{2} \ln^2 x + (2 - x)e^{-x} + C$

(C) $\frac{1}{2} x^2 \ln x + (2 - x)e^{-x} + C$

(D) $\frac{1}{2} x^2 \ln x - (x - 1)e^{-x} + C$

$$\int \frac{1}{x}(\ln x + x^2 e^{-x}) dx = \int \left(\frac{\ln x}{x} + x e^{-x} \right) dx$$

$$= \int \left(\frac{\ln x}{x} + x e^{-x} \right) dx \text{ by parts}$$

$$\begin{aligned} \text{let } u = \ln x & \quad \text{let } u = x, du = dx \\ = du = \frac{1}{x} dx & \quad dv = e^{-x} \quad v = -e^{-x} \\ dx = x du & \end{aligned}$$

$$= \int \frac{u}{x} x du + uv - \int v dv$$

$$= \int u du + x(-e^{-x} du)$$

$$= \frac{u^2}{2} + c_1 - x e^{-x} - e^{-x} + C_2$$

$$= \frac{(\ln x)^2}{2} - e^{-x}(x + 1) + C$$

Ans. A

38. Let $f(x) = \frac{3x+1}{x-2}$, then what is the range of $f(x)$?

(A) $\mathbb{R} \setminus \{2\}$

(B) $\mathbb{R}$

(C) $\mathbb{R} \setminus \left\{-\frac{1}{3}\right\}$

(D) $\mathbb{R} \setminus \{3\}$

$$F(x) = \frac{3x+1}{x-2} \text{ Domain} = \mathbb{R} \setminus \{2\} \text{ and this graph has horizontal asymptote at}$$

$$y = 3 \text{ and the graph never crosses this horizontal asymptote}$$

$$\Rightarrow \text{Range} = \mathbb{R} \setminus \{3\}$$

Ans. D

39. Which one of the following is true about the graph of $(x) \frac{2x^3+2x^2+3x}{x^2+x}$?

(A) The graph has y-intercept at (0,3)

(B) The vertical asymptotes of the graph are at $x=0$ and $x=-1$ and its oblique asymptote is $y=2x$.

(C) The graph has at least one x-intercept.

(D) The vertical asymptotes of the graph are at $x=0$ and $x=-1$ but it has no horizontal asymptote.

$$F(x) = \frac{2x^3+2x^2+3x}{x^2+x} = \frac{x(2x^2+2x+3)}{x(x+1)} \\ = \frac{2x^2+2x+3}{x+1}$$

The graph has vertical asymptote at $x = -1$ and $x=0$ is a hole. This graph also has oblique asymptote we can find by long division

$$\begin{array}{r} 2x \\ x^2 \overline{) 2x^3 + 2x^2 + 3x} \\ \underline{2x^3 + 2x^2 + 3x} \end{array}$$

$\Rightarrow y = 2x$ is oblique the others are not correct

Ans. B

40. A private college has 1000 students. 60 % of these students are males. 45% of these students pay their payment by credit card including 175 females. What is the probability that the student is a male or a credit card user?

(A) 0.675

(B) 0.225

(C) 0.775

(D) 0.325

Let m be male students and c be students that uses card

$$P(m) = \frac{600}{100}, p(c) = \frac{450}{100}, p(mnc) = \frac{275}{100} (450 - 175)$$

$$P(muc) = p(m) + p(c) - p(mnc)$$

$$= \frac{600}{100} + \frac{450}{100} - \frac{275}{100} = 0.775$$

Ans. C

41. If $z = (1 + \sqrt{3}i)(1 + i)$, then which one of the following is the polar representation of z ?

(A) $z = 2\sqrt{2} \left(\cos(105^\circ) + i \sin(105^\circ) \right)$

(B) $z = 2\sqrt{2} \left(\cos(15^\circ) + i \sin(15^\circ) \right)$

(C) $z = 4 \left(\cos(105^\circ) + i \sin(105^\circ) \right)$

(D) $z = 4 \left(\cos(75^\circ) + i \sin(75^\circ) \right)$

$$Z = (1 + \sqrt{3}i)(1 + i) \Rightarrow \begin{aligned} z &= (1 + i + \sqrt{3}i - \sqrt{3}) \\ z &= 1 - \sqrt{3} + (1 + \sqrt{3})i \end{aligned}$$

$$R = \sqrt{(1 - \sqrt{3})^2 + (1 + \sqrt{3})^2}$$

$$R = \sqrt{1 - 2\sqrt{3} + 3 + 1 + 2\sqrt{3} + 3}$$

$$R = \sqrt{8} = 2\sqrt{2}$$

$$\theta = \tan^{-1} \left(\frac{1 + \sqrt{3}}{1 - \sqrt{3}} \right)$$

θ is on the 2nd quadrant and $\theta = 150^\circ$

$Z = 2\sqrt{2} (\cos(150^\circ) + i \sin(150^\circ))$ is polar representation

Ans. A

42. What is the sum of the series $\sum_{n=1}^{\infty} 3^n 4^{-n}$?

(A) ∞

(B) $3/16$

(C) 4

(D) 3

$$\sum_{n=1}^{\infty} 3^n \cdot 4^{-n}$$

if $n = 1$, $\sum_{n=1}^{\infty} 3^n \cdot 4^{-n}$ is $\frac{3}{4}$

if $n = 2$, $\sum_{n=1}^{\infty} 3^n \cdot 4^{-n} = \frac{9}{16}$

if $n = 3$, $\sum_{n=1}^{\infty} 3^n \cdot 4^{-n} = \frac{27}{64}$

It is geometric progression with $G_1 = \frac{3}{4}$ and $r = \frac{3}{4}$

$$\Rightarrow \sum_{n=1}^{\infty} 3^n \cdot 4^{-n} = S_{\infty} = \frac{G_1}{1-r} = \frac{\frac{3}{4}}{1-\frac{3}{4}} = 4$$

Ans. C

43. Which one of the following is equal to $\frac{d}{dx} \log_2 \sqrt{6x}$?

(A) $\frac{3x}{2 \ln(2)}$

(B) $\frac{1}{2x \ln(2)}$

(C) $\frac{3}{2x \ln(2)}$

(D) $\frac{1}{6x \ln(2)}$

$$\begin{aligned} &= \frac{1}{\sqrt{6x \ln 2}}, (\sqrt{6}) \\ \frac{d}{dx} (\log_2 \sqrt{6x}) &= \frac{1}{\sqrt{6x \ln 2}} \cdot \frac{1}{\sqrt{6x}} \cdot 6 \\ &= \frac{6}{2x \sqrt{6x \ln 2}} = \frac{1}{2x \ln 2} \end{aligned}$$

Ans. B

44. Let $f(x) = \ln(x\sqrt{x})$. Then what is $f'(x)$ equal to ?

(A) $2x/3$

(B) $\sqrt{x}/2$

(C) $3/2$

$$\begin{aligned} F(x) &= \ln(x\sqrt{x}) \Rightarrow f'(x) = \frac{1}{x\sqrt{x}} \cdot \left(x \cdot \frac{3}{2}\right) dx \\ &= \frac{1}{x\sqrt{x}} \cdot \frac{3}{2} x^{\frac{1}{2}} \end{aligned}$$

45. What is the maximum possible area of a rectangle in square units with diagonal of length 16 units ?

(A) 128

(B) 64

(C) 48

(D) 256

Soln area = $x y$

$$x^2 + Y^2 = (16)^2 = 256$$

$$Y = \sqrt{256 - x^2}$$

$$\text{Area: } A(x) = x\sqrt{256 - x^2} = \sqrt{256x^2 - x^4}$$

$$\text{Let } g(x) = 256x^2 - x^4 \text{ and } g'(x) = 512x - 4x^3$$


$$g'(x) = 0 \Rightarrow 512x - 4x^3 = 0$$

$$4x(128 - x^2) = 0$$

$$x = 0 \text{ or } x\sqrt{128} = 8\sqrt{8^2}$$

$$\text{The maximum area} = (8\sqrt{2})(8\sqrt{2}) = 128 =$$

Ans. A



46. Let $U = \mathbb{N}$ (the set of natural numbers) be the universe. Which one of the following proposition is true?

(A) $(\exists x)(x + = x)$

(B) $(\forall x)(\exists y)(x - y = x)$

(C) $(\forall x)(\exists y)(y < x)$

(D) $(\forall x)(\exists y)(x \div y = y \div x)$

Let $u = \mathbb{N}$ The true proposition is

$$(\forall x)(\exists y)(x \div y = y \div x)$$

$$(\forall x)(\exists y)\left(\frac{x}{y} = \frac{y}{x}\right)$$

47. What are the values of a and b so that the function $f(x) = \begin{cases} x+1 & x < 1 \\ ax+b, & 1 \leq x < 2 \\ 3x, & x \geq 2 \end{cases}$

is continuous every where ?

(A) $a = 4, b = 2$

(B) $a = 4, b = -2$

(C) $a = -4, b = -2$

(D) $a = -4, b = 2$

$$F(x) = \begin{cases} x+1, & x < 1 \\ ax+b, & 1 \leq x < 2 \\ 3x, & x \geq 2 \end{cases}$$

F(x) is continuous at $x = 1$

$$\Rightarrow \lim_{x \rightarrow 1^-} (x+1) = f(1) \Rightarrow \lim_{x \rightarrow 1} x+1 = ax+b$$

$$2 = a+b$$

F(x) is continuous on $x = 2$

$$\Rightarrow \lim_{x \rightarrow 2^-} (a+b) = f(2) \Rightarrow \lim_{x \rightarrow 2} ax+b = 3x$$

$$2a+b = 6 \dots 2$$

Solve 1 and simulation eagle

$$\begin{cases} a+b=2 \\ 2a+b=6 \\ -a=-4 \end{cases} \Rightarrow \begin{cases} a=4 \\ a+b=2 \\ 4+b=2 \end{cases}$$

$$b = -2$$

F(x) is continuous when $a = 4$ and $b = -2$

Ans. B

48. A cylindrical tank whose inner diameter is 2m contains $4\pi \text{ m}^3$ oil. If the oil is discharged from the tank at the rate of $\frac{2\pi}{3} \text{ m}^3/\text{min}$, then how long (in min) does it take for the tank to be empty?

(A) $\frac{4}{3}$

(B) 4

- (C) 6
(D) 12

$\sqrt{v} = Abh = \pi v^2$ since it is cylindrical tank given diameter = 2m and radius = 1m volume = 4π

$$\frac{dv}{dt} = \frac{2\pi}{2} m^3/min$$

$$V = \pi r^2 h$$

$$4\pi m^3 = \pi (1)^2 m 2h$$

$$H = 4m = \text{distance where } 5 \text{ vt}$$

$$\frac{dv}{dt} = \sqrt{v} = \frac{2}{3} m/min$$

$$\Rightarrow t = \frac{5}{v}$$

$$= \frac{4m}{\frac{2}{3} m/min} = 6 \text{ min} = t$$

Ans. C

49. Let $f(x) = x - x^2$ and $g(x) = \frac{1}{x}$. Then what is $\left(f\left(\frac{1}{x}\right)\right)$ equals to

- (A) $\frac{x^2}{x-1}$
(B) $x - x^2$
(C) $\frac{1}{x^2 - x}$
(D) $\frac{x-1}{x^2}$

$$F(x) = x - x^2, g(x) = \frac{1}{x}$$

$$= 9\left(\frac{1}{x} - \frac{1}{x^2}\right)$$

$$9\left(f\left(\frac{1}{x}\right)\right) = 9 \frac{x-1}{x^2}$$

$$= \frac{1}{\frac{x-1}{x^2}} = \frac{x^2}{x-1}$$

Ans. D

50. variance of 20 observations is 5. If each observation is multiplied by 2, the what is the variance of the resulting observations?

- (A) 5
(B) 10
(C) 40

(D) 20

If x is the variance of x, y, z and t then the variance $2x, 2y, 2z$ and $2t$ is $2^2 x$

$\Rightarrow$ The new variance $= 2^2 x = 20$

Ans. D

51. If $f(x) = k \ln x + e^{\sin x}$ and $f''(\pi) = -1$, then what is the value of k ?

(A) π

(B) $2\pi^2$

(C) π^2

$$F(x) = k \ln x + e^{\sin x}$$

$$F'(x) = \frac{k}{x} + \cos x e^{\sin x}$$

$$F''(x) = \frac{k}{x^2} - \sin x e^{\sin x} + \cos x e^{\sin x} \cos x$$

$$= \frac{k}{x^2} - \sin x e^{\sin x} + \cos^2 x e^{\sin x}$$

$$\frac{-k}{\pi^2} - 0 + 1 = -1$$

$$F''(\pi) = -1 \quad = \frac{-k}{\pi^2} = -2$$

$$= k = 2\pi^2$$

Ans. C


52. If θ is a fourth quadrant angle and $\sec \theta = \sqrt{2}$ then what is $\csc \theta$ equal to?

(A) $1/\sqrt{2}$

(B) $-1/\sqrt{2}$

(C) $-\sqrt{2}$

(D) $\sqrt{2}$



$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \sqrt{2}$$

$$\Rightarrow \sec \theta = \frac{\text{hyp}}{\text{adj}} = \sqrt{2} \text{ but it is}$$

Quadrant 4 ($\sec \theta = \text{negative}$) $\sec \theta = -\sqrt{2}$

Ans. C

53. Consider the following assertion:

$p + 2^n$ is an odd number for any prime p and any $n \in \mathbb{N}$.

Which one of the following is correct about a prove or disprove of the assertion?

(A) There is a counter example that disproves the assertion.

(B) The assertion can be proved by direct method; because p is odd and $2^n = 2(2^{n-1})$ is even imply that $p + 2^n$ is odd since the sum of odd and even is odd.

(C) The assertion can be proved by the indirect method; because if $n \notin \mathbb{N}$, then $2^n \notin \mathbb{N}$ and hence $p + 2^n$ is not odd.

(D) The assertion can be disproved by the method of contradiction.

Solve. $p + 2^n$ is an add number

$p + 2^n$ is add numbe for any prime p and only $n \in \mathbb{N}$ The asser tian can be proved by indirect proo of becouse there is a counter example that disprove the assertion $p = 2$ Is only even proem Number $p + 2^n$ is even.

Ans. C

54. What is the standard equation of the line passing through the point (2,3) and parallel to the

line given by $\begin{cases} x = 1 + 2\lambda \\ y = -2 - \lambda \end{cases}, \lambda \in \mathbb{R}$?

(A) $\frac{x-2}{-1} = \frac{y-3}{2}$

(B) $\frac{x-1}{1} = \frac{y-3}{-2}$

(C) $\frac{x-1}{2} = \frac{y-3}{-2}$

(D) $\frac{x-2}{2} = \frac{y-3}{-1}$

Solve The standard equation

(2, 3) parallel to $\begin{cases} x = 1 + 2\lambda \\ y = -2 - \lambda \end{cases}$

Use water equation of the line $x = x_0 + \lambda v$

$$\lambda = \frac{x-1}{2} = \frac{y+2}{-1} \text{ and parallel vator } v = (1_2 - 1)$$

$$r = r_0 + \lambda v \Leftrightarrow r = (x, y) = (3, 3) + \lambda (2, -1)$$

$$\therefore x = 2 + 2\lambda \text{ and } 9 = 3 - \lambda$$

$$\lambda = \frac{x-2}{2} \text{ and } \lambda = \frac{y-3}{-1}$$

Ans. D


55. If in ΔABC , $AB = 3$, $BC = 4$ and $m(\angle B) = 60^\circ$, then what are the length of AC and the cosine of $\angle A$, respectively?

(A) $\sqrt{13}$ and $\frac{6}{5\sqrt{13}}$

(B) $\sqrt{13}$ and $\frac{1}{\sqrt{13}}$

(C) $\sqrt{13}$ and $\frac{-1}{\sqrt{13}}$

(D) $\sqrt{13}$ and $\frac{-6}{5\sqrt{13}}$



Using cosine law

$$\begin{aligned} x^2 &= 3^2 + 4^2 - 2(3 \cdot 4)\cos 60^\circ \\ &= 9 + 16 - 24(\cos 60^\circ) \\ &= 25 - 12 \left(\frac{1}{2}\right) \end{aligned}$$

$$x^2 = 25 - 12 = 13$$

$$x = \sqrt{13}$$

And using sin law

$$\begin{aligned} \frac{4}{\sin A} &= \frac{\sqrt{13}}{\sin 60^\circ} \\ \frac{4}{\sin A} &= \frac{\sqrt{13}}{\frac{\sqrt{3}}{2}} \end{aligned}$$

Ans. B

$$\sqrt{13} \sin A = \left(\frac{\sqrt{3}}{2}\right)$$

$$\sin A = \frac{2\sqrt{3}}{\sqrt{13}}$$

$$\text{but } \sin 2\theta + \cos 2\theta$$

$$\cos 2\theta = -\sin 2\theta$$

$$= 1 - \left(\frac{2\sqrt{3}}{\sqrt{13}}\right)$$

$$= 1 - \frac{12}{13} = \frac{1}{13}$$

$$\cos = \frac{1}{\sqrt{13}}$$

56. Let $p(n)$ be an open proposition on the set of natural numbers ($\mathbb{N}$). Which one of the following is a correct application of the principle of mathematical induction?

(A) If $P(1)$ is true; for $n = 1$; and if both $P(n)$ and $P(n + 1)$ are true for certain $n \in \mathbb{N}$, then $P(n)$ is true for all $n \in \mathbb{N}$.

(B) If $P(10)$ is true; and if $P(n)$ is true for implies that $P(n + 1)$ is true, then $P(n)$ is true for all $n \in \mathbb{N}$.

(C) If $P(10)$ is true; and assuming $P(n)$ is true for any $n > 10$ if it follows that $P(n + 1)$ is true, then $P(n)$ is true for all $n \geq 10$.

(D) If $P(1)$ is true; and $P(n) \Rightarrow P(n + 1)$ is true for any $n \in \mathbb{N}$, then $P(n)$ is true for all $n \in \mathbb{N}$.

If $P(1)$ is true for $n = 1$

If $P(n)$ is true for $n \in \mathbb{N}$ and $P(n + 1)$ is true for $n + 1 \in \mathbb{N}$. Then $P(n)$ is true for $n \in \mathbb{N}$

Ans. A

57. If the image of the line $2x - 3y = 7$ under a translation is $2x - 3y = 0$, which one of the following is a translation vector of the translation line?

(A) $u = (-2, 1)$

(B) $u = (1, -2)$

(C) $u = (-1, 2)$

(D) $u = (2, -1)$

$$2x - 3y = 7 \xrightarrow{v(a,b)} 2x - 3y = 0$$

$$\Leftrightarrow P(x, y) \xrightarrow{v(a,b)} (x', y')$$

$$X + a = x' \Leftrightarrow x = (x' - a)$$

$$Y + b = y' \Leftrightarrow y = (y' - b)$$

$$\therefore 2(x - a) - 3(y - b) = 7 \rightarrow 2x - 3y = 0$$

$$2x - 2a - 3y + 3b = 7 \rightarrow 2x - 3y = 0$$

$$2x - 3y = 7 + 2a - 3b$$

$$\Leftrightarrow \frac{7 + 2a - 3b = 0}{2a - 3b = -7} \Leftrightarrow a = -\frac{7 + 3b}{2}$$

Ans. A

58. Let $P = (1, a, a)$ and $Q = (a - 1, 1, 1)$ be points in space and the distance between P and Q is 3.

Then what is the value (S) of a ? (M)

(A) $a = -3, a = \frac{1}{3}$

(B) $a = 1, a = -9$

(C) $a = -1, a = 9$

(D) $a = -3, a = -\frac{1}{3}$

$$\begin{aligned}
 D &= \sqrt{(a-1)-1)^2 + a-1^2 + a-1)^2} \\
 d^2 &= (a-2)^2 + a^2 - 2a + 1 + a^2 - 2a + 1 \\
 \Rightarrow a^2 - 4a + 4 + a^2 - 2a + 1 + a^2 - 2a + 1 &= 9 \\
 \Rightarrow 3a^2 - 8a - 3 &= 0 \\
 \Rightarrow 3a^2 - 9a + a - 3 &= 0 \\
 \Rightarrow 3a(a-3) + 1(a-3) &= 3 \\
 3a+1=0 \text{ or } a-3=0 \\
 A = -\frac{1}{3} \text{ or } a = 3
 \end{aligned}$$

Ans. D

59. What is the image of the circle $x^2 + y^2 - 4x - 6y + 12 = 0$ when it is reflected with respect to the line $y = -x$?

- (A) $(x-2)^2 + (y-3)^2 = 1$
 (B) $(x+3)^2 + (y+2)^2 = 1$
 (C) $(x+2)^2 + (y+3)^2 = 1$
 (D) $(x-3)^2 + (y-2)^2 = 1$

Soln. The image of circle $x^2 - 4x + y^2 - 6y + 12 = 0$

$$x^2 - 4x + 4 + y^2 - 6y + a = 4 + 9 + 12$$

The center of circle is (1,3)

$$(2,3) \xrightarrow{\text{reflected } y=-x} (-3,-2)$$

$$\therefore \text{image } (x+3)^2 + (y+2)^2 = 1$$

Ans. B

60. If the dot products of a vector $\vec{A}$ with the vectors $i - j + k, 2i + j - 3k$ and $i + j + k$ are 4, 0 and 2, respectively, what is $\vec{A}$?

- (A) $\vec{A} = (2, 1, 1)$
 (B) $\vec{A} = (-2, 1, -1)$
 (C) $\vec{A} = (2, -1, 1)$
 (D) $\vec{A} = (-2, -1, 1)$

ASoln. let $u = i-j+ki$ $u = 2i+j-3k$ $w = i+5+k$

Let $A = xi+yi+2k$

$$A \cdot u = x-y+z=4 \dots 1$$

$$A \cdot u = u \Rightarrow 2x + y - 3z = 0 \dots 2$$

$$A \cdot w = 2 \Rightarrow x + y + z = 2 \dots 3$$

From 1 and 2 and 1 and 3 respectively

$$\begin{cases} 3x - 2k = 4 \Rightarrow 5x = 10 \\ 2x + 2z = 6 \Rightarrow x = 2 \end{cases}$$

$$Z = 3 - x \Rightarrow z = 3 - 2 = 1$$

$$Y = x + z - 4$$

$$Y = 2 + 1 - 4 = -1$$

Vector A is $(x, y, z) = (2, -1, 1)$

Ans. C

61. In order to measure the height a of a tower, suppose a surveyor takes two sightings from a transit 1 meter high which are positioned d meters apart on the same ground level as in the figure below.


If the first measured angle of elevation is α and the second is β (see, the figure), then what is the height of the tower (in meter) in terms of α, β and d ?

(A) $\left(\frac{\tan \alpha \tan \beta}{\tan \alpha - \tan \beta} \right) d + 1$

(B) $\frac{\tan \alpha \tan \beta}{(\tan \alpha - \tan \beta) d} + 1$

(C) $\left(\frac{\tan \alpha - \tan \beta}{\tan \alpha \tan \beta} \right) d + 1$

(D) $\left(\frac{\tan \alpha \tan \beta}{\tan \alpha + \tan \beta} \right) d + 1$



Let $h = h_1 + 1m$

Let the height of tower x be d/c b/n the end survey or sighting

$$\tan \alpha = \frac{h}{x-d} \Rightarrow h_1 = (x-d) \tan \alpha \dots 1$$

$$\tan \beta = \frac{h}{x} \Rightarrow h_1 = x \tan \beta \dots 2$$

from 1 and 2, $x \tan \beta = (x-d) \tan \alpha$

$$\Rightarrow x \tan \beta + d \tan \alpha = x \tan \alpha$$

$$x(\tan \alpha - \tan \beta) = d \tan \alpha \Rightarrow x = \frac{d \tan \alpha}{\tan \alpha - \tan \beta}$$

$$H = h_1 + 1m = \frac{d \tan \alpha}{\tan \alpha - \tan \beta} \tan \beta + 1$$

$$H = \left(\frac{\tan \alpha - \tan \beta}{\tan \alpha - \tan \beta} \right) d + 1$$

Ans. A

62. Let l be the line given by the vector equation $(x, y) = (-2, 1) + \lambda(1, 1), \lambda \in \mathbb{R}$. which one of the following is the equation of the image of l after being translated by the vector $u = (2, -1)$ followed by a rotation through 45° about the origin?

(A) $y = -2\sqrt{2}x$

(B) $y = \sqrt{2}$

(C) $y = \sqrt{2}x$

(D) $x = 0$

Soln. vector equation $(x, y) = (-2, 1) + \lambda(1, 1), \lambda \in \mathbb{R}$

$$x + 2 = y \Rightarrow y = x + 2$$

$Y = x + 2 \xrightarrow{\text{translated by } (2, -1)} y = x$ take $(1, 1)$ be a

Point on $y = x$, then $(1, 1) \xrightarrow{\theta = 45^\circ}$

$$X' = x \cos \theta - y \sin \theta$$

$$X' = 1 \cos 45^\circ - 1 \sin 45^\circ$$

$$Y' = x \sin \theta + y \cos \theta$$

$$Y' = 1 \sin 45^\circ + 1 \cos 45^\circ = \frac{2\sqrt{2}}{2} = \sqrt{2}$$

$\therefore$ The image is $y = \sqrt{2}$

Ans. B

63. What is the period (P) and the range of $f(x) = 5 \sin\left(\frac{1}{3}x + 2\right) + 3$?

(A) $P = 6\pi$, range $[-5, 5]$

(B) $P = 6\pi$, range $[-2, 8]$

(C) $P = \frac{2\pi}{3}$, range $[-5, 5]$

(D) $P = \frac{2\pi}{3}$, range $[-2, 8]$

Soln. $f(x) = 5 \sin\left(\frac{1}{3}x + 2\right) + 3$

$$\text{Period} = \frac{2\pi}{1/3} = 6\pi$$


$$\text{Range} = [-a + c, a + c] = \{-5 + 3, 5 + 3\} = (-2, 8)$$

The period and range of f is 6π and $[-2, 8]$ respectively

Ans. B

64. Let $A(a, 2, 5)$, for $a > 0$, be a point on the sphere $x^2 + y^2 + z^2 + 6z = 0$ and C be the center of the sphere. If $P(k, 2, 4)$ is a point in space such that $\overrightarrow{PA}$ is perpendicular to $\overrightarrow{CA}$, what is the cosine of the angle between $\overrightarrow{PA}$ and $\overrightarrow{PC}$?

- (A) $\frac{5}{\sqrt{35}}$
 (B) $\frac{7}{\sqrt{35}}$
 (C) $\frac{5}{\sqrt{70}}$
 (D) $\frac{7}{\sqrt{70}}$



Soln.

$A = (a, 2, 5)$ for $a > 0$ be point on the sphere $x^2 + y^2 + z^2 - 6z = 0$

$$\Rightarrow x^2 + y^2 + z^2 - 6z + 9 = 9 \Rightarrow x^2 + y^2 + (z - 3)^2 = 9 \text{ and } r = 3$$

C be center of the sphere i.e $(0, 0, 3)$

$P(K, 2, 4)$ is a point the sphere

$$\overrightarrow{PA} \perp \overrightarrow{CA}, \overrightarrow{CA} = (1, 2, 3)$$

$$r = \sqrt{a^2 + 2^2 + 2^2} = 3$$

$$a^2 + 4 + 4 = 9 \Rightarrow a^2 = 1 \Rightarrow a = 1$$

$$\overrightarrow{PA} = (1 - k, 0, 1)$$

$$\overrightarrow{PA} \cdot \overrightarrow{CA} = 0 \Rightarrow 1 - k + 0 + 2 = 0 \Rightarrow k = 3$$

$$\cos \theta = \frac{\overrightarrow{PA} \cdot \overrightarrow{CP}}{|\overrightarrow{PA}| |\overrightarrow{CP}|}$$

$$\overrightarrow{PC} = (-3, -2, -1) \quad |\overrightarrow{CP}| = \sqrt{3^2 + 2^2 + 1^2} = \sqrt{14}$$

$$\overrightarrow{PA} = (-2, 0, 1) \quad |\overrightarrow{PA}| = \sqrt{4 + 0 + 1} = \sqrt{5}$$

$$\cos \theta = \frac{\overrightarrow{PA} \cdot \overrightarrow{CP}}{|\overrightarrow{PA}| |\overrightarrow{CP}|} = \frac{6 + 0 + (-1)}{\sqrt{5} \sqrt{14}} = \frac{5}{\sqrt{5} \sqrt{14}} = \frac{5}{\sqrt{70}}$$

Ans. C

65. What is the value of $\lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{2}{n} \left[\left(\frac{2k}{n} \right)^2 + 5 \left(\frac{2k}{n} \right) \right]$

(A) 14

(B) 10

(C) 4

(D) 18

Soln. $\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{2}{n} \left[\left(\frac{2k}{n} \right)^3 + 5 \left(\frac{2k}{n} \right) \right]$

$$\lim_{n \rightarrow \infty} \frac{16}{n^4} \sum_{k=1}^n k^3 + \frac{20}{n^2} \sum_{k=1}^n k$$

$$\lim_{n \rightarrow \infty} \frac{16}{n^4} \left(\frac{n^2(n+1)^2}{4} \right) + \lim_{n \rightarrow \infty} \frac{20(n(n+1))}{2n^2}$$

$$\frac{16}{4} \lim_{n \rightarrow \infty} \left(\frac{n^4 + 2n^3 + n^2}{n^4} \right) + \frac{20}{2} \lim_{n \rightarrow \infty} \frac{n^2 + n}{n^2} = 4 + 10 = 14$$

Ans. A

