

Isticmaalka daawada

**Bugaan waxaa ku baran dontaa habka daawada lo
isticmaalo**

***Abdishahid Abdikadir Mohamed
(Dr Shaahid)***

Drug class Helmentic

Albendazole tablet /syrup Daawadan waxay ku timaadaa kaniini iyo sharoobo

Albendazol. Waxaa loo yaqanaa **Albendazolium**

Indication or principle uses.

Waxaa lagu daweeeyaa gooryanada kala duwan sida

- Giardiasis
- Trichuriasis
- Ascariasis
- Filariasis
- Pinworm disease
- Taenia sainata
- Taenia solium
- Parasite infection
- Tapworm

How to drug work.. Sidee dawadaan ushaqeeysaa

Dawadaan waxay faragalin kala duwan kusameysaa iyoo hab awood leh kusoo saaresa dayactireysana sokorta kana sameyneyso awood

Side effect: dhibaata ay keeni karto

- Unugyada difaaca jirka oo hoos u dhaca
- ,xinjirow ga diiga oo hoos u dhaca
- cudurada beerka ayey ku keeni kartaa hadii inbadan la qaato,
- cunto xumi ayaa lagu arki karaa ,
- cuncun iyo finan maqaarkaa

Dosage =Daawada qiyaas ahaan la siin karo

Caruurta ka weyn 6. lixbilood iyo dadka waa weyn waxaa la siinayaa 1 xabo kaniin ah 400mg halmar

Caruurta lixbilood ka weyn uu miisaankoodu ka yaryahay 10kg waxaa la siinayaa 200mg halmar

Precaution :taxadarka

Waa in la qaataa talooyinka dhaqtarku ku siinaayo

Over dose daawada oo qiyaasteeda layska badiyo

Majiro ilaa hada wax dhibaato ah oo la diiwaangaliyay

Drug intrection = daawada aan lala qaadan karin

In ta bada waxaan lala qaadan karin daawada loo yaqaano

- Amethazone
- Praziquantel
- Theophylline

Food intrection= cuntada saamaynta ku leh

Ma jiraan cunto saameeyn ku leh

Disease intrection= ma jiraa cudur lid ku ah

- Cirrhosis,
- cudurada beerka ku dhaca ayna keento isticmaalka aashitada sida cabitaanada

Contraindication lama siinkaro

Lama siin karo hooyada uurka leh sadexda bilood ee ugu horaysa ,uurkeeda

Laakiin hadii hooyadu nuujinayso cunug weyqaadan kartaa

Hasiin cunuga ka yar 6 bilood

Membendazole

Composition= daawadu waxa ay ka samaysan tahay

Benzoyl benzimidazole and carbamate

Treaputic drug = daawooyinka uu ka mid yahay

Anti helmentic

Available= maxaa la heli karaa daawadan

Waxaa laga heli karaa kaniini iyo sharoobo

Indication or principle uses. Isticmaalka daawada

dawadaan waxaa loo isticmaalaa qaar kamid ah goor yaanada

worm infaction. Sida

- ascariasis
- pinworm diseases
- hook worm infaction
- hydatid disease
- and giardia

Side effect =dhibaatada ay keento

- Matag vomitin
- Madax xanuun headache

How to drug work =side daawadaani u shaqaysaa

Mebendizole waxay dhex gashaa inay soo saarto hab ay ku jirto ka hortag gooryaanka ee laga soo nuugay sonkorta

Daawada qiyaasta laga qaato = Dosage

daawadan waxaa la siinkaraa caruurta ka weyn 2 sanno iyo dadka waaweyn 100mg mudo sadex maalmood ah daawadaan ku celi bukaanka 2 ilaa 3 usbuuc

Drug interactions= ma jiraan dawooyin ay isdiidayaana

waxaan lala qaadan Karin daawwadan bamazepine

food interaction ma jiraan cunto aan lagu qaadan Karin daawada

lamayaqaano cunto dhibta

disease intraction= ma jiraan cuduro saamaynku leh daawadan

hadii aad qabto cudurada ay ka midyihiin cudurka beerka ama crohn,s disease ama ulcerative colitis

Drug class malaria Chloroquine

Brand name: Aralen ,aralen HCl

Aralen phosphate tablet 500mg /syrup/injection

Daawadan waxay ku timaadaa kaniini iyo sharoobo

Indication or principal uses. Istimaalka

Waxaa lagu dawaayaa loogana hortagaa malariyada oo ay keento plasmodium sidoo kale waxaa lagu dawayaa **amebiasis parasite** aadka u adag iyo lafa xanuunka

Common side effect. Dhibaatooyinka ugu caansan

- | | | |
|---|-----------------------|------------------|
|  | Dhibaato murqaha ah.. | muscle problems |
|  | Cuntada oo xiranta... | loss of appetite |
|  | Shuban | diarrhea |
|  | Nabro maqaarka ah | skin rash |

Serious side effects... dhibatooyinka ugu qatarsan

- Aragti la, aan whth out vision
- Qatar kutimaada murqaha muscle damage
- Unugyada dhiiga oo hoos udhaca.. low blood cells

How to drug work.. sidee dawadaan ushaqeysaa

Daawadan chloroquine waa sun loo isticmaalo cudurka malariada

Dietary advice = cuntada haboon

Qaado cunto ama caano si loo yareeyo dhibaataada caloosha

Precaution :taxadarka

Waa in la qaataa talooyinka dhaqtarku ku siinaayo

Over dose= daawada oo qiyaasteeda layska badiyo

- ❖ Madax xanuun
- ❖ Dawaqaad iyo wareer
- ❖ Aragtida oo isbedesha
- ❖ Isbedelku yimaada garaacista wadnaha
- ❖ Cadaadiska dhiiga oo hoos u dhaca

Drug intrection= daawada aan lala qaadan karin

Magnesium salts and anti acid cemitadine

Food intrection= cuntada saamaynta ku leh

Ma jiraan cunto saameeyn ku leh

Disease intrection= ma jiraa cudur lid ku ah

- Cudurada beerka
- cudurada neerfayaasha

Contraindication= lama siinkaro

Lama siin karo hooyada uurka leh sadexda bilood ee ugu horaysa ,uurkeeda

Laakiin hadii hooyadu nuujinayso cunug weyqaadan kartaa

Metronidzole.

Brand name . flagyl

Waxay kutimadaa

	Kaniinyaal	tablets
	Sharoopo	syrup
	Faleembo	infucion

Indication or principle uses. Waxa loo istimaalo

Dawadaan waxaa loo is timalaan gooryamada kala duwan sida

- Amebisis
- Giardiasis
- Trichomonisis

Sidookala metronidzole waxaa loo isticmali kara xanuunada ku dhaca kadi mareenka iyo xanuunada ey keenaan **bacterial vaginosis**

How to drug work. Side dawadaan ushaqeysaa

Dawadaan waxay burburisaa difaaca gooryanada waxayna ku kuuleysataa inay suuxiso ama aydisho gooryanada

Dosege. Qiyaasta dawadaan laga qaadan karo

- ❖ dadka waawey waxaa lasiyaa 500mg kanini 16 xabo 1x2
- ❖ faleembo infusion 12 xabo 1x2
- ❖ caruurta yayar shoropo 200mg 1x2 dawo siinta caruuta waxay ku xiran tahat da,da iyo miisaanka

Side effect, dhibaatooyinka uu yeelan karo

- matag nausea
- cuntada oo ka xirmata loss of appetite
- madax xanuun . headaches

looma isticmaali karo b

over dose. Dawadaan oo qiyaastayda la iska badiyo

waxaa kadhalaan kara dhibaatooyin sida

- lalabo
- matag

food intrection. Cunto saameyn kuleh

lalama isticmaalikar cuntooyinka aashitada leh sida

- muufada ama qamiirka

Disease intrection . cudur lid ku ah

Waxaa lid ku ah cudurada kala ah

-  neerfaha
-  kilya xanuunka
-  beer xanuun
-  ka

Drug class malaria

HydroxyChloroquine sulfate

Brand name: magaca kiimikada waa

Plaquenil Apothecon tablet 200mg

Daawadan waxay ku timaadaa kaniini iyo sharoobo waxayna ka kooban tahay

Daawada malariada iyo daawada caabuqa ,iyo daawada lafoxanuunka

Indication or principal uses... isticmaalka

Waxaa lagu daweyaa loogana hortagaa malariyada oo ay keento **plasmodium** waxaan sidookale lagu daween karaa romatiisimada aadka udaran

Stomach upset

Side effect =dhibaataada ay keeni karto

- lalabo nausea
- calool xanuun stomach crmps
- cuntada oo xiranta loss of appetite
- shuban diarrhea
- daciif nimo dizziness
- madax xanuun headche

How to drug work ... side dawadaan ushaqeysa

Daawadan hydroxychloroquine waa daawo loo isticmaalo cudurka malariada iyo lafo xanuunka waxay hoos u dhigayaan kiimikada taas oo sababaysa xanuuno

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Caruurta yar yar waxaa ku haboon in la siiyo sharoobooyinka hadii ay waxkatariwaayaan ayaa looisticmaali karaa cirbad iyadoo loo fiirinaayo qiyaasta ku haboon hydroxychloroquine

Laakiin dadka waaweyn waxaa la siinayaa kaniini ama cirbad chloroquine tablet 500mg maalintii halmar ama 150mg injection iyo wixii kale ee la heli karo

Hydroxychloroquine 200mg tablet

Age	0	2 Months	1 year	5 year	15 year	Adult
-----	---	----------	--------	--------	---------	-------

weight		4 Kg	8 kg	15 Kg	35 kg	
100mg base table day 1 day 2		$\frac{1}{2}$	1tablet	2.1/2	6	
Day 3		$\frac{1}{4}$	$\frac{1}{2}$	1	3	
150 mg table base Day 1 and day 2		$\frac{1}{8}$	$\frac{1}{2}$	1.1/2	4	
Day 3		$\frac{1}{8}$	$\frac{1}{4}$	$\frac{3}{4}$	2	

Dietary advice cuntada haboon

Qaado cunto ama caano si loo yareeyo dhibaataada caloosha

Precaution :taxadarka

Waa in la qaataa talooyinka dhaqtarku ku siinaayo

Over dose daawada oo qiyaasteeda layska badiyo

- ❖ Madax xanuun
- ❖ dawaqaad

Drug intrection daawada aan lala qaadan karin

- Magnesium salts
- Anta acid
- cemitadine or digoxin

Disease intrection ma jiraa cudur lid ku ah

- Dhiig yaraan
- Dhiig bax
- Unugyada cadcad oo hoos u dhaca

Mefloquine

Brand name: magaca kiimikeed

lariam , tablet

Daawadan waxay ku timaadaa kaniini iyo sharoobo waxayna ka kooban tahay

Daawada malariada iyo daawada tayo aad u leh cudurka duumada

Indication or principal uses = isticmaalka

Waxaa lagu daaweeyaa malariada ama duumada ku soo boodka ah oo ay keentay plasmodium falciprium iyo plasmodium vivax isdoo kale waxaa loo isticmaalaa kahortaga daawada malaria ama markay jirka adkaysi ku yeelato daawada chloroquine

How to drug work

Mefloquine waa daawo loo isticmaalo malariada taas oo sun ahaan u disha protaza

Side effect. Dhibatooyinka ay leeyihiin

- ❖ Matag vomiting
- ❖ Shuban diarrhea
- ❖ Madax xanuun headaches
- ❖ Nabro rash

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Caruurta yar yar waxaa ku haboon in la siiyo sharoobooyinka hadii ay waxkatariwaayaan ayaa loo isticmaali karaa cirbad iyadoo loo fiirinaayo qiyaasta ku haboon hydroxychloroquine

Laakiin dadka waaweyn waxaa la siinayaa kaniini ama cirbad chloroquine tablet 500mg maalintii halmar ama 150mg injection iyo wixii kale ee la heli karo

Hydroxychloroquine 250mg tablet

Age	0	2 Months	1 year	5 year	15 year	Adult
weight		4 Kg	8 kg	15 Kg	35 kg	
100mg base table day 1 day 2		$\frac{1}{2}$	1tablet	2.1/2	6	
Day 3		$\frac{1}{4}$	$\frac{1}{2}$	1	3	
150 mg table base Day 1 and day 2		$\frac{1}{8}$	$\frac{1}{2}$	1.1/2	4	
Day 3		$\frac{1}{8}$	$\frac{1}{4}$	$\frac{3}{4}$	2	

Dietary advice cuntada haboon

Daawadan ha qaadan adigoo baahan ugu yaraan biyo badan u isticmaal

Precaution :taxadarka

Waa in la qaataa talooyinka dhaqtarku ku siinaayo

Daawadan hala isticmalin wax alaale iyo wixii
muqaadaraad ah iyo alcolo ah

Over dose daawada oo qiyaasteeda layska badiyo

Waxay isku dhibaato yihiin waxyeelada daawada side
effect

Drug intrection daawada aan lala qaadan karin

Quinine ,choloroquine,calcium canal blockers,anti
histamine iyo andti convulsants

Disease intrection cuduro lid ku ma jiraan

Cudurada wadnaha,kuwa indhaha ,maskax xanuunka iyo
shaqada beerk

Permethrin

Brand name: elimite,Nix

permethrin ointment or cream

Daawadan waxay ku timadaa

-  Cream
-  solution

Indication or principal uses = isticmaalka

Waxaa loo isticmaalaa injirta madaxa gasha iyo xaaladaha kadhaha isnaadama ku dhaca jirka qeebihiida kala duwan dawadaan waxaa laga isticmaalaa qeebaha sare ee jirka

topical uses

How to drug work

Daawadan waxay suuxisaa injirta kadibna wey dishaa wax dhibaataana uma laha bini aadanka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa lagu daaweeyaa injirta marka la dhaqo madaxa ka dib waxaana loo marinayaa hal marka toban daqiiqo ka dib waa laga dhaqayaa todobo maalin ka dib ayaa lagu calinayaa

Side effect.. dhibatada ay leedahay

- Cuncun itching
- Barar swelling

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Tobon daqiiqo

Duration of action xiliga uu soconkaro

Tobon maalmood

Dietary advice cuntada haboon

Daawadan waa inaad qaada taa adigoo baahan

Prolonged use hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadato waa in la ogaadaa in daawadii ay baabi isay injirtii timaha madaxa

Precaution :taxadarka

Daawadan u isticmaal caruurta yar yar malahan dhibaato

Over dose daawada oo qiyaasteeda layska badiyo

Majirto bukaano la diwaan galiyey oo ay ka soo gaartay dhibaato

Drug intrection daawada aan lala qaadan karin

Daawadan hala isticmaalin cream

kale ilaa ay iyadu ka dhamaanayso

Food intrection cuntada saamaynta ku leh

Ma jiraan cunto saameeyn ku leh

Disease intrection cuduro lid ku ma jiraan

Hadii ay jiraan cuduro maqaarka ah midkasta iskii u daaweey

Piperazine

Brand name: piperazine

Daawadan waxay ku timaadaa

kaniini Tablets

Indication or principal uses Tilmaanta

Waxaa loo isticmaalaa gooryaano badan oo kala duwan sida

- 1 ascariasis
- 2 Enterobius vermicularis
- 3 worm infaction
- 4 Roun worm

How to drug work

Daawadan waxay suuxisaa gooryaanka kadibna waxay ka soo saartaa saxarada

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Waxaa lagu daaweeyaa gooryaano kala duwan waxaana la siinayaa 3. 5g labo maalin

Ku celi hal isbuuc ,caruurta sii 75mg ku celi 2 isbuuc ka dib

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Lama yaqaano

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Ma jiraan cunto dhibaato ku ah

Prolonged use hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadatao waa in baaris lagu sameeyo

Precaution :taxadarka

Daawadan u isticmaal caruurta yar yar malahan dhibaato

Dawadaan hooyada uurka leh way isticmaali kartaa

Over dose= daawada oo qiyaasteeda layska badiyo

Majirto bukaano la diwaan galiyey oo ay ka soo gaartay dhibaato

Drug intrection= daawada aan lala qaadan karin

Daawadan hala isticmaalin phenothiazines

Food intrection= cuntada saamaynta ku leh

Ma jiraan cunto saameeyn ku leh

Disease intrection =cuduro lid ku ma jiraan

cvcvcv bhadii aad qabto qalal wuxuu kuu keenikaraa dibaato

Praziquantel

Brand name: biltricide

Antihelmentic tablet 600mg

Daawadan waxay ku timaadaa kaniini

Indication or principal uses = waxa loo isticmaalo

Waxaa loo isticmaalaa gooryaano badan oo kala duwan sida

Worm infection

- 1 Schistosomiasis
- 2 Clonorciasis
- 3 Tap worm infection
- 4 Hydaid disease

Dhamaan gooryaanadan waa in lagu daaweeyaa praziuantel waana goor yanada ugu xooga badan

How to drug work:

Daawadan waxay kala dhig dhigtaa murqaha gooryaanadan iyo difaacooda kadibna waxay ku guulaysataa inay dibada u soo saarto

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa la siiyaa dadka waaweyn iyo caruurta yar yar waxaana loo qaataa 2.2ibs 1kg 1x3

Side affect.. dhibatada uu leeyahay

- ✚ Calool xanuun abdominal pain
- ✚ Matag vomiting
- ✚ Madax xanuun headache
- ✚ Xasaasiyad allergic
- ✚ Haddii uu uur jiro lama isticmaali karo

Onset of effect= xiliga uu bilaawdo dhibaatiisa

Lama yaqaano

Duration of action = xiliga uu soconkaro

Lama yaqaano

Dietary advice =cuntada haboon

Daawadan waad qaadan kartaa cuntada kadib ama inta aad ku gudajirto cabitaanada laakiin hajajabin daawadan

Prolonged use =hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadata waa in baaris lagu sameeyo

Precaution :taxadarka

Daawadan u isticmaal sida dhaqtarku kuu amray

Over dose= daawada oo qiyaasteeda layska badiyo

Majirto bukaano la diwaan galiyey oo ay ka soo gaartay dhibaato

Drug intrection= daawada aan lala qaadan karin

Daawadan hala isticmaalin corticosteroids,keteconazole

Food intrection =cuntada saamaynta ku leh

Ma jiraan cunto saameeyn ku leh

Disease intrection= cuduro lid ku ma jiraan

hadii aad qabto cudurada beerka

Drug class anti protozoal

Metronidazole

Brand name: magaca kiimikeed waa bactericide

Protozoal tablet 400mg, 250mg, 500mg syr 200mg

Daawadan waxay ku timaadaa kaniini sharoobo siboosto kaabsool

Side effect dhibaataada ay keeni karto

Daawadan qatar teedo waxay ku xiran tahay qaybaha hoose ee ah common iyo less common

Common= dhibaato caan ah oo ay keensato daawadan

- ❖ Lalabo iyo matag
- ❖ Cunto xumo, calool xanuun
- ❖ Shunaban marmar ah iyo saro adeyg
- ❖ Finan iyo cuncun, barar

Less common= dhibaato yar ayaa lagu arki karaa sida

- Finan maqaarka ah
- cuncun

Indication or principal uses= isticmaalka

Waxaa loo isticmaalaa gooryaano badan oo kala duwan sida

- 1 Amebic dysenteric infection diarrhea
- 2 Gairdia lambia
- 3 Iyo kuwo kale

Dhamaan gooryaanadan waa in lagu daaweeyaa praziuantel

How to drug work = sida daawadu u shaqeeyso

Daawadan waxay kala dhig dhigtaa murqaha gooryaanadan iyo difaacooda kadibna waxay ku guulaysataa inay dibada u soo saarto iyadoo suuxinaysa kadibna jawaabta ugu danbaysa weydishaa

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa la siiyaa dadka waaweyn waxaa la siiyaa kanii 500mg ama 100ml infusion iyo caruurta yar yar waxaana loo qaataa sharoobo waxayna ku xiran tahay cudurada loo isticmaalaayo hadi ay tahay protozoa or bacteria

Onset of effect xiliga uu bilaawdo dhibaatiisa

Lama yaqaano

Duration of action = xiliga uu soconkaro

Daawadan waxay soconkartaa 14maalmood

Dietary advice = cuntada haboon

Daawadan waad qaadan kartaa cuntada kahor ilaa 10/20 daqiiqo

Prolonged use= hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadata waa in baaris lagu sameeyo

Precaution :taxadarka

Daawadan u isticmaal sida dhaqtarku kuu amray

Over dose= daawada oo qiyaasteeda layska badiyo

- Lalabo
- matag
- qalal

Drug intrection= daawada aan lala qaadan karin

Daawadan hala isticmaalin asprine,iyo kuwo kale

Food intrection =cuntada saamaynta ku leh

Cuntooyinka ay aashitada ku badantahay waxaa lama ogola waa in dib looga dhigaa ilaa mudo 1saac ah

Disease intrection= cuduro lid ku ma jiraan

cudurada beerka cudurada neerfayaasha nadiifinta kalyaha ,cusbada badan

infection Bacterial

Drug class: penicillin antibiotic

Amoxicillin

Brand name: amoxil,larotid, moxlin,polymox,trimox,wymox

**Protozoal tablet 500mg inject,250mg cop, 500mg tablet
1gr inject liquid drops**

Daawadan waxay ku timaadaa kaniini sharoobo kaabsool

Side effect dhibaataada ay keeni karto

Kalagoosyada oo xanuuna , calool xanuu daran ,qandho ,saxaro dhiig iyo calool majiir iyo faruuryaha oo barara iyo neefta oo dhibta carabka cunaha ,wajiga oo barara cuncun iyo finan jirka ka soo baxa

Common dibaato caan ah oo ay keensato daawadan

Finan ,shuban dhexdhexaad ah ,lalabo iyo matag madax xanuun , iyo vaginaha oo dheecaan ka yimaado cuncunana afka oo xanuuna cadaadana

Less common dhibaato yar ayaa lagu arki karaa sida

Tabar daro daal ,kaadi badan,qarqaryo

Indication or principal uses Tilmaanta

Waxaa loo isticmaalaa jeermiska bacteriada oo ay ka mid yihiin

- 1 Dhago xanuunka
- 2 Cunah iyo sanku
- 3 Xubinta taranka
- 4 Hawomareenka
- 5 Jabtada,qaliinka kahor iyo kadiba ,caabuqa ku dhaca wadnaha
- 6 Gaaska boog calooleedka iyo kuwo kale

How to drug work sida daawadu u shaqeeyso

Daawadaani waxay xanibtaa bacteriada darbiga unugeeda mana awoodeeyso inay isbadiso ama ay faafto

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa la siiyaa caruurta yar yar iyo dadka waaweyn

Dadka waaweyn **500mg sided saac kasta kadib**

Caruurta yar yar 250 mg sided saac kasta kadib iyadoo loo fiirinaayo dada cunuga iyo qiyaasta daawada

Onset of effect xiliga uu bilaawdo dhibaatiisa

Labo saac gudahood

Duration of action xiliga uu soconkaro

Daawadan waxay soconkartaa 8saac

Dietary advice cuntada haboon

Waxaa fiican baahida in lagu qaato laakiin cuntada kadib waa la qaadan karaa waxay hoos u dhigaysaa cuncunka caloosha oo ay yareeyneysaa

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato

Stopping the drug

Daawadan waxaad joojinkartaa mark ay kaa dhamaato laakiin hadii aad ku fiicnaato iyadoo daawadii wax ka dhiman yihiin dhamaystir

Prolonged use hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadatao waxaa kordhaaya dhibaataadeeda ee inta laguugu talo galay hadiifin

Precaution :taxadarka

Ma jirto dhibaato la ogyahay oo laga filaayo

Over dose daawada oo qiyaasteeda layska badiyo

Lalabo iyo matag ,shuban , murqo daal ,qalal

Drug intrection daawada aan lala qaadan karin

Daawadan hala isticmaalin pulfnpyrazone
,cholestyramine

Food intrection cuntada saamaynta ku leh

Lama yaqaano cunto dhibaysa

Disease intrection cuduro lid ku ma jiraan

cudurka alleriga ,neefta,wadno xanuunka,

Bacterial infection

Drug class: penicillin antibiotic
combination Amoxicillin/potassium
clavulanate

Brand name: augmentin

**Protozoal tablet ,250mg cop, 500mg tablet 1000mg
125mg syrup**

Daawadan waxay ku timaadaa kaniini

Side effect= dhibaataada ay keeni karto

- Calool xanuun daran
- ,kala goysyada oo xanuun
- saxaro dhiig
- xasaasiyad daran
- barar degdeg ah,
- hawada oo yaraata
- cuncun maqaarka ah iyo finan,
- indhaha iyo maqaarka oo guduuta

Common =dibaato caan ah oo ay keensato daawadan

Finan ,shuban dhexdhexaad ah ,lalabo iyo matag madax xanuun , iyo vaginaha oo dheecaan ka yimaado cuncunana afka oo xanuuna cadaadana

Less common= dhibaato yar ayaa lagu arki karaa sida

Tabar daro, daal ,

Indication or principal uses =Tilmaanta

Waxaa loo isticmaalaa jeermiska bacteriada oo ay ka mid yihiin

- 1 Dhago xanuunka
- 2 Cunah iyo sanku
- 3 Xubinta taranka
- 4 Hawomareenka
- 5 Jabtada,qaliinka kahor iyo kadiba ,caabuqa ku dhaca wadnaha
- 6 Gaaska boog calooleedka iyo kuwo kale
- 7 Sanboor
- 8 Iyo jeermisyada ay keenaan bacteraya laakiin kuma daaweyn karno kuwa ay keenaan fungus,viruska iyo noolayaasha kale

How to drug work = sida daawadu u shaqeeyso

Daawadani waxay xanibaysaa darbiga unuga bacteriyada iyadoo u diidayso inuu isbadiyo waxaana intabadan loo

isticmaalaa marka ay adkaysi u yeelato daawooyin kale
maadaama ay ku jirto clavulanate

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa la siiyaa caruurta yar yar iyo dadka
waaweyn

Dadka waaweyn **500mg sided saac kasta kadib 1000mg**

**Caruurta yar yar 250 mg sided saac kasta kadib iyadoo loo
fiirinaayo dada cunuga iyo qiyaasta daawada ama sharoobo
420mg**

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Hal ilaa Labo saac gudahood

Duration of action xiliga uu soconkaro

Daawadan waxay soconkartaa 6=8saac

Dietary advice cuntada haboon

Waxaa fiican baahida in lagu qaato laakiin cuntada kadib
waa la qaadan karaa waxay hoos u dhigaysaa cuncunka
caloosha oo ay yareeyneysaa

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc
ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato

Stoping the drug

Daawadan waxaad joojinkartaa mark ay kaa dhamaato laakiin hadii aad ku fiicnaato iyadoo daawadii wax ka dhiman yihiin dhamaystir

Prolonged use hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadata waxaa kordhaaya dhibaataadeeda ee inta laguugu talo galay hadhiifin waxaana dhacaysa in jirku la qabsado oo waxkatariweyso

Precaution :taxadarka

Ma jirto dhibaato la ogyahay oo laga filaayo

Over dose daawada oo qiyaasteeda layska badiyo

Lalabo iyo matag ,shuban Daran ,alal

Drug intrection daawada aan lala qaadan karin

Daawadan hala isticmaalin erythromycin tetracycline or contraceptive

Food intrection cuntada saamaynta ku leh

Lama yaqaano cunto dhibaysa

Disease intrection cuduro lid ku ma jiraan

cudurka alleriga ,neefta,wadno xanuunka,gaaska

Azitromycine

Brand name: zitromax

Daawadan waxay ku timaadaa kaniini kaabsool injection sharoopo,

Indication or principal uses Tilmaanta

Waxaa loo isticmaalaa jeermiska bacteriada oo ay ka mid yihiin

- | | | |
|---|------------------|----------------------|
| 1 | Oof wareen | pnemonia |
| 2 | Caabuqa Dhagaha | ear infection |
| 3 | Shuban sumoobid | traveler diarrhea |
| 4 | Caabuqa caloosha | intestinal infection |

Waxaa kaloo lagu daweenkaraa cudurada la, isku gudbiyo galmoodka **sexual transmitted infection** sida

-  Chlamydia
-  Gonorrhea infection

Sidoo kale waxaa loo isticmaali karaa malaria

How to drug work sida daawadu u shaqeeyso

Daawadani waxay ka hortagtaa unugyada bacteriyada ama jeermiska ee laga soo saarayo brotinka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadani waxaa la siiyaa 125mg 250mg 500mg 2 waqti
ama 1000mg daily ama 1200mg weekly

Dadka waaweyn 500mg sided saac kasta kadib 1000mg

Daawadan hadii ay tahay caruur tayaryar raac warqada ay wadato daawadan

Side effect dhibaatooyinka ay keeni karto

- Shuban diarrhea
- Saxarada oo lawaayo loose stools
- Calool xanuun abdominal pain
- Matag vomiting
- Daciifnimo dizziness

Onset of effect= xiliga uu bilaawdo dhibaataadiisa

Lama yaqaan daawadan bilaawgeeda

Duration of action = xiliga uu soconkaro

Lama yaqaan xiliga uu soconayo

Dietary advice = cuntada haboon

Waxaa fiican baahida in lagu qaato laakiin cuntada kadib waa la qaadan karaa waxay hoos u dhigaysaa cuncunka caloosha oo ay yareeyneysaa waxaa fiican in cuntada laga dabamariyo 2 saac

Prolonged use= hadii daawada oo waqti dheer la qaato

Hadii aad daawadan waqtidheer qaadato waxaa kordhaaya dhibaataadeeda ee inta laguugu talo galay hadhiifin waxaana dhacaysa in jirku la qabsado oo waxkatariweyso

Precaution :taxadarka

Ka taxadar dhibaataada ay keeni karto oo u isticmaal sida ay tahay

Over dose= daawada oo qiyaasteeda layska badiyo

Ma jirto daawadan dhibaato la diiwaan galiyey oo la ogyahay

Drug intrection daawada aan lala qaadan karin

Warfarin ,phnytoin

Food intrection cuntada saamaynta ku leh

Daawadan qaado caloosha oo baahan

Disease intrection cuduro lid ku ma jiraan
cudurada beerka

Cefixime

Brand name: suprax

Daawadan waxay ku timaadaa kaniini sharoobo iyo cirbad

Indicatoin or principal uses waxa loo isticmaalo

Dawadaan waxaa loo isticmalaa xanuunada kaladuwan sida

	Dhaga xanuun	otitis media
	Cunaha	strep throat
	Qaar jeexa	pneumonia
	Kaadi mareenka	urinary tract infection
	Jabtada	gonorrhea

Side effect dhibatooyinka ay leeyihiin

Common side effect

- Shuban diarrhea
- Lalabo nausea
- Calool xanuun abdominal pain
- Matag vomiting
- Nabro maqaarka ah skin
- Kal goosy xanuun Joint pain
- Cuncun itching

How to drug work sida daawadu u shaqeeyso

Daawadani waxay ka hortagtaa unugyada bacteriyada ama jeermiska ee laga soo saarayo brotinka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawada waxaa la siiyaa 100mg 200mg ama 400mg labo waxti maalintii

Onset of effect= xiliga uu bilaawdo dhibaataadiisa

2ilaa 4saacadood

Duration of action= xiliga uu soconkaro

6=18 saacadood

Dietary advice cuntada haboon

Daawadan waxaa la qaataa cuntada ka dib

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

Prolonged use= hadii daawada oo waqti dheer la qaato

Daawadani waxay keeni kartaa dhibaato hadii isticmaalkeeda la badiyo

Precaution :taxadarka

Dadka ka weyn 60 uga taxadar

Over dose daawada oo qiyaasteeda layska badiyo

Hadii dasawadan waxbadan laga qaato waxay keeni kartaa qalal,calool xanuun shuban iyo matag

Drug intrection daawada aan lala qaadan karin

Heparin ,

Food intrection cuntada saamaynta ku leh

Lama yaqaano

Disease intrection cuduro lid ku ma jiraan

kalyo xanuunka iyo jeermiska colitis

Cephalexine

Brand name: cefanex,keflex,keftab

Cefixim tablet or capsul , 500mg cop,250 oral suspension

Daawadan waxay ku timaadaa kaniini kaabsool iyo sharoobo

Indication or principal uses= waxa loo isticmaalo dawadaan

Dawadaan waxaa lagu daweeeyaa qaar kamid ah bagtariyada waxaan lagu dilaa labada nuuc ee **gram.positive iyo gram.**

Negative bacteria

- 1 Xanuunada kaadi mareenka urinary tract infection
- 2 Infagshanka lafaha bone infaction
- 3 Infagshanka dhagaha ear infaction
- 4 Infagshanka dhuunta respiratory infaction
- 5 Infagshanka kaadi heysta bladder infaction

How to drug work =sida daawadu u shaqeeyso

Daawadani waxay ka hortagtaa unugyada bacteriyada iyadoo samaysanaysa darbi difaac

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawada waxaa la siiyaa 250mg ama 500mg labo waxti maalintii

Side effect. Dhobatooyinka ay layihiin dawadaan

-  Calool xanuun abdominal pain
-  Kala goosyo xanuun joint pain
-  Cucun saxaha ah vaginal itching
-  Daal tiredness

Onset of effect= xiliga uu bilaawdo dhibaataadiisa

1 saacadood

Duration of action = xiliga uu soconkaro

Lama yaqaano

Dietary advice = cuntada haboon

Daawadan waxaa la qaataa iyadoo wax lacunay ama ay caloosha marantahay

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadani waxay keeni kartaa dhibaato hadii isticmaalkeeda la badiyo

Precaution :taxadarka

Dadka ka weyn 60 uga taxadar

Over dose= daawada oo qiyaasteeda layska badiyo

Hadii daawadan waxbadan laga qaato waxay keeni kartaa qalal,calool xanuun shuban iyo matag

Drug intrection daawada aan lala qaadan karin

Food intrection cuntada saamaynta ku leh

Lama yaqaano

Disease intrection cuduro lid ku ma jiraan

kalyo xanuunka iyo jeermiska colitis

Ceprofexine

Brand name: clioxan

Cefixim tablet or capsul , 500mg cop,250 oral suspension

Indication or principal uses =isticmaalka dawadaan

Dawadaan waxaa loo isticmalaa waxa yaabaha ay keento bagtariyada infagshan sida

- 1 Infagshan calooleed abdominal infaction

- 2 Infagshanka lafaha joint infaction
- 3 Infagshanka kudhaca dhuunta respiratory tract infaction
- 4 Infagshanka maqaarka skin infaction
- 5 Qandhada tiifowga typhoid fever
- 6 Kaadi mareenka urinary tract infaction

How to drug work = sida daawadu u shaqeeyso

Daawadan waxay dhexgashaa enzymka bacteriada aadka u koriya

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawada waxaa la siiyaa 250mg ama 500mg labo waxti maalintii mudo 1 isbuuc ah

Side effect dibatooyinka dawadaan

- Madax xanuun heache
- Barar gacmahaga iyo cagta swelling in your hands and feet
- Dhibaato indhahaga propems with your eyes

Onset of effect= xiliga uu bilaawdo dhibaatadiisa

Lama yaqaan

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice= cuntada haboon

Daawadan waxaa la qaataa iyadoo wax lacunay ama ay calooshu marantahay dhib na ma keento

Prolonged use =hadii daawada oo waqti dheer la qaato

Daawadan waxaa la dheerayn karaa marka dhaqtar kuu soo qoro oo kali ah

Precaution :taxadarka

Dadka ka weyn 60 uga taxadar

Over dose= daawada oo qiyaasteeda layska badiyo

Wax la og yahay oo ilaa hada la diiwaan galiyey ma jiraan

Drug intrection= daawada aan lala qaadan karin

Hadii labo dhaqtar daawooyin kala duwan aad ka soo qaadata iyadoon lagaa koontaroolin

Food intrection cuntada saamaynta ku leh

Lama yaqaano

Disease intrection cuduro lid ku ma jiraan

hadii aad qabto cuncun oo aad ka qaadayso daawadan cefproflaxine

Doxycycline

Brand name: apo.doxy,doryx,,monodox,vibramycin

doxycycline capsul , 100mg cop

Daawadan waxay ku timaadaa kabsul

Indication or principal uses= isticmaalka

Waxaa loo isticmaalaa in lagu daweeyo infagshanka ey sababto bacteria and cartain parasites dabcan goor yanada waxaana loo isticmaalaa

- | | | |
|---|-------------|---------------------|
| 1 | Oofwareen | Pnemonia |
| 2 | Finanka | acne |
| 3 | | Chlamydia infection |
| 4 | Daacuun | cholera |
| 5 | Isfiilitada | syphilis |

How to drug work= sida daawadu u shaqeeyso

Daawadani waxay dishaa bacteria and protozou waxay hoos u dhigtaa soosaarida borotiinka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawada waxaa la siiyaa 100mg labo waqti ama 200mg

Side effect dhibatooyinka uu la yahay

	Shuban	diarrhea
	Lalabo	nausea
	Matag	vomitin

Onset of effect= xiliga uu bilaawdo dhibaatiisa

Shan maalmood kadib

Duration of action = xiliga uu soconkaro

Maalmo ayaa la isticmaalikaraa

Dietary advice= cuntada haboon

Daawadan waxaa la qaataa iyadoo wax lacunay
waxaadna la isticmaalaysaa galas biyo ah oo buuxa

Prolonged use= hadii daawada oo waqti dheer la qaato

Daawadani waxay keeni kartaa dhibaato hadii
isticmaalkeeda la badiyo

Over dose= daawada oo qiyaasteeda layska badiyo

Lalabo iyo matag ,shuban waxliqista oo ku adkaata

Drug intrection daawada aan lala qaadan karin

Warfrin

Food intrection= cuntada saamaynta ku leh

Hala siin caanaha ee ka danbeysii ilaa halsaac ama kabadan

Disease intrection= cuduro lid ku ma jiraan

kalyo xanuunka iyo beerka xanuun

Erythromycin

Brand name: E.mycin,EES,Ery,Erythrocin

Erythromycin. 125mg 250mg 1000mg

Daawadan waxay ku timaadaa kabsul ,kaniini,sharoobo iyo cirbad

Indication or principal uses .. isticmaalka

Waxaa loo isticmaalaa in lagudaweeyo qaar kamid ah bactariyada iyo waxyaabaha ay sababaa

- 1 Infagshanka hawo mareenka respiratory tract infection
- 2 Infagshanka maqaarka skin infection
- 3 Xanuunada kaadi mareenka pelvic inflammatory disease
- 4 Chlamydia infection

How to drug work sida daawadu u shaqeeyso

Daawadan waxay ka hortagtaa jeermiska bacteriada unugyadeeda iyo soo saarida proteinka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawada waxaa la siiyaa 250mg labo waqti ama 1000mg ama 500mg lix saac kasta inta ay daawadu kuu dhamaanayso

Side effect dhibaatooyinka uu leeyahay

- Calool abdominal cramps
- Matagag vomitin
- Shuban diarrhea
- Xasaasiyad allergic
- Dhibaato beerka ah liver problems

Dadka xasaasiyada ka qaada daawada **penicillin** erythromycin waxay utahay badbaado

Onset of effect xiliga uu bilaawdo dhibaatadiisa

Si dhaqso ayey u bilaabata infection

Laakiin hadii afka laga qaadanaayo lama yaqaano

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Daawadan waxaa la qaataa iyadoo iyadoo calooshu maran tahay ugu yaraan hal saac ama cuntada ka dib 2 saac

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadani waxay keeni kartaa dhibaato hadii isticmaalkeeda la badiyo

Precaution :taxadarka

Daawadani waxay keeni kartaa in ay maqalk wax ka badasho okey uurka

=pregnancy daawadani waxay keeni kartaa inay dhaawacdo beerka ee inta aadan qaadan la tasho dhaqtarka ku shaqada leh

Over dose daawada oo qiyaasteeda layska badiyo

Lalabo iyo matag ,shuban wareer ,calool xanuun , maqalka oo yaraada

Drug intrection daawada aan lala qaadan karin

phenothiazines

Food intrection cuntada saamaynta ku leh

Lama yaqaano cuntada saamayta ku leh

Disease intrection cuduro lid ku ma jiraan

wadno xanuun ,kalyo xanuun ,cudurada beerka

Penicillin G

Brand name: bicillin,crystasilline 300 pentids,

Benzylpenicillin waxaa loo yaqaanaa **penicillin G** waxaana la hindisay 1929 waxaana hidisay [Alexander flaming](#)

Penicillin injection 1gr or 500mg

Daawadan waxay ku timaadaa cirbad

Indication or principal uses .. isticmaalka dawadaan

Waxaa loo isticmalaa qaar kamid ah bacterial infection waxaana lagu daweeeyaa qaar kamid ah bacterial infection , sida

- Oofwareen pneumonia
- Quman step throed
- Isfiilitada syphilis
- Goowracato diphtheria
- Goojiska tetanus
- Masgax garaad meningitis

How to drug work sida daawadu u shaqeeyso

Daawadan waxay xanibtaa unugyada bacteriada iyo darbiyadeeda si aysan isu faafin

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawada waxaa la siiyaa 500000unite ama 200000 unite
1,200,000 unite ama 1000000 unite labo waqti lix saac kasta
inta ay daawadu kuu dhamaanayso

Side effect dhibaatooyinka ayleeyihiin

- ❖ Xasaasiyad allergic reactions
- ❖ Shuban diarrhea

Onset of effect xiliga uu bilaawdo dhibaata diisa

Si dhaqso ah halooga siiyo xididka

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Daawadan waxaa la qaataa iyadoo calooshu maran tahay ugu yaraan hal saac ama cuntada ka dib 2 saac

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadani waxay keeni kartaa dhibaato hadii isticmaalkeeda la badiyo marka waa in la baaraa qofka markay daawadu u dhamaato

Precaution :taxadarka

Daawadan malahan wax report oo la diiwaan galiyey

=pregnancy daawadani umala waxdhibaato ah hooyada xiliga ay uurkeeda

Over dose daawada oo qiyaasteeda layska badiyo

Lalabo iyo matag ,shuban wareer ,

Drug intrection daawada aan lala qaadan karin

Cholestyramine

Food intrection cuntada saamaynta ku leh

Lama yaqaano cuntada saamaynta ku leh

Disease intrection cuduro lid ku ma jiraan

cuncun , neef,dhiigbax sida hemophilia ,wadno xanuun,cudurada caloosha sida gaaska

Tetracycline hydrochloride

Brand name: sumycin

Penicillin capsul , 250mg or 500mg

Daawadan waxay ku timaadaa kabsul ,

Indication or principal uses .. isticmaalka

Dawadaan waa **antibiotic** oo loo isticmaalo in lagu daweeyo qaar kamid xanuunada kala duwan sida

-  Finanka acne
-  Daacuunka cholera
-  Buruseelaha brucellosis
-  Plague
-  Is fiilitada syphilis

How to drug work sida daawadu u shaqeeyso

Daawadan waxay dilaysaa oona hoos u dhigaysaa soosaarida jeermisyada iyo protenka ay u baahan yihiin

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa la siiyaa dadka qaba cudurada aan soo sheegnay waana 250mg ama 500mg ilaa 4 waqti ayaa la siinkaraa

Side effect.. dhibaatooyinka ayleeyihiin dawadaan

- Matag vomiting
- Shuban diarrhea
- Nabro rash
- Cuntada oo xiranta loss of appetite
- Waxay dhibaato uleedahay ikaha hadii lasiiyo caruurta kayar 8sano

Onset of effect xiliga uu bilaawdo dhibaatiisa

Lama yaqaano

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Daawadan waxaa wanaagsan caloosha oo maran in la qaato halgalaas oo biyo ahna lagu cabo

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan hadii layka badiyo waxay keeni kartaa inay jirka la qabsato

Precaution :taxadarka

Daawadan malahan wax report oo la diiwaan galiyey

=pregnancy daawadani looma isticmaali karo hooyo uurleh

Over dose... daawada oo qiyaasteeda layska badiyo

Lalabo iyo matag ,shuban liqitaanka oo ku adkaada ,

Drug intrection daawada aan lala qaadan karin

Daawadan hala qaadan calcium supplements ,choline

Food intrection cuntada saamaynta ku leh

Caanaha ha la isticmaalin

Disease intrection cuduro lid ku ma jiraan

cudurada kalyaha ,iyo beerka

Fungus

Drug class: topical anti fungul

Bethametazone

Brand name: lotrisone

Daawadan waxay ku timaadaa

boomaato cream

kaiini tablets

cirbad injection

hawo inhalation

Indication or principal uses Tilmaanta

Dawadaan waxaaa loo isticmalaa qaar kamid ah cudurada ay kamid yihiin.

- ❖ Ramatiisimada rheumatoid arthritis
 - ❖ Cudurada maqaarka skin diseases
 - ❖ Xasaasiyada allergic
 - ❖ Neefta
 - ❖ Cansarka cancers such as leuemia
- 1 ku daawee fungus infectioun kiisa

How to drug work sida daawadu u shaqeeyso

Daawadaan waxa ka hortagtaa jeermiska fungaska waxayna u diidaa in la soosaaro protienka taas oo dhex gasha lana falgasha jeermiskan daawadani waxay wax ka qabataa fungaska iyo xanuuno kale

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo mariyaa labo waqti subax iyo habeen daawadan waxaa ku darsan anti inflammentory sida steroid isticmaalka dawadaan waxay ku xirantahay nuuca daawada iyo da,da qofka

Side effect.. dhibata ay leedahay dawadaan

-  murqaha oo dabca muscle weakness
-  xasasiyad kala duwan allergic
-  dawadaan hadii waqti dheer la isticmaalo waxay keenneysaa dhibaato aad Qatar u ah waxaana kamid ah
-  in ay tintu kori waayso
-  maqaarka jirka oo isbadala

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Sii bukaanka cunto ku filan

Precaution :taxadarka

Daawadani waxay u daran tahay dadka dada ah

=pregnancy daawadani ma malahan dhibaati ka diiwaab galiyey

Over dose daawada oo qiyaasteeda layska badiyo

Malahan case la diiwaan galiyey oo dhibaato ,

Drug intrection daawada aan lala qaadan karin

Daawadani ma lahan daawo aan lala qaadan karin ee la diiwaan galiyey

Food intrection cuntada saamaynta ku leh

Ma jiro cunto ka hor imaanaysa daawadani

Disease intrection cuduro lid ku ma jiraan

cudurada cuncunka ama allerjiga

Fungus

Drug class: topical anti fungul

Fluconazole

Brand name: diflucan

Fluconazole : dawadaan waxay kutimadaa

Kaniiniyaal tablets 50mg , 100mg, 150mg

Boomaato cream

Cirbado injection

Faleembo infusion

Indication or principal uses .. isticmaalka

Dawadaan waxaa loo isticmaalaa qaar kamid ah waxayaabaha uu sabao fangay infagshan sida waxyaabaha maqaarka kazoo baxa iyo cudurakale

1. cabeebka afka ama saxaha dumarka candidiasis
2. maqaarka kore dermatophytosis

How to drug work sida daawadu u shaqeeyso

Daawadaan waxa ka hortagtaa jeermiska fungaska iyo waliba daaweynta

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo isticmaalaa dadka waaweyn iyo caruurta dhexe 200=400mg injection vaginaha funguska ku dhaca ayaa loo isticmaalaa 150mg suspenton

Side effect.. dhibatooyinka ay leeyihiin dawadaan

- ✓ Matag vomiting
- ✓ Shuban diarrhea
- ✓ Nabro rash
- ✓ Dhiicaanka beerka oo hoos udaco increased liver inzyms

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Lama yaqaano goorta ay bilaabanayo

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Biyo badan ku cab daawadan hadii ay sharoobo tahayna aad u rug

Precaution :taxadarka

Daawadani waxay u daran tahay dadka qaba kalyo xanuunka,

sidoo kale ha isticmaalin alcolada iyo waxyaabaha aashitada leh

=pregnancy daawadani ma malahan dhibaato la diiwaan galiyey

Over dose daawada oo qiyaasteeda layska badiyo

Malahan case la diiwaan galiyey oo dhibaato ,

Drug intrection daawada aan lala qaadan karin

Daawadan lalama isticmalikaro rifabutin or phenytion

Food intrection cuntada saamaynta ku leh

Ma jiro cunto ka hor imaanaysa daawadani

Disease intrection cuduro lid ku ma jiraan

cudurada beerka kuwa kalyaha

Fungus

Drug class: topical anti fungul

Griseofulvin

Brand name: fulvicin,grivulvin

Griseofulvin :copsul and table

Daawadan waxay ku timaadaa kaniini iyo kaabsool

Side effect dhibaataada ay keeni karto

Finan maqaarka ah iyo cuncun iyo wareer cuncun maqaarka ah

Common dibaato caan ah oo ay keensato daawadan

Madax xanuun

Less comman dhibaato yar ayaa lagu arki karaa sida

Calool xanuun lalabo iyo matag daal iyo wareer shuban

Indication or principal uses Tilmaanta

- 1 ku daawee fungus infectioun kala duwan ay ku jiraan ring worm tinea barae ,tinea capatis tinea corporis iyo tinea cruris ,nail fungus

How to drug work sida daawadu u shaqeeyso

Daawadaan waxa ka hortagtaa jeermiska fungaska waxayna soo saartaa waxyaabaha nool ee u baahan in la qashinsaaro

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo isticmaalaa dadka waaweyn iyo caruurta dhexe 250=400mg injection vaginaha funguska ku dhaca ayaa loo isticmaalaa 150mg suspenton

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Lama yaqaano goorta ay bilaabanayo

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Daawadan cuntada ka hor iyo kadib waad qaadan kartaa u isticmaal biyo badan

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato una qaadataa sidii lagu faray

Stoping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan lagama badin karo usbuucyo hadii aad ka badiso waxay keeni kartaa beerxanuun iyo kalyo xanuun

Precaution :taxadarka

Iska ilaali meelaha boorka badan wasaqda badan ama aadida warshadaha soo saara wasaqda inta aad isticmaalayso, sidoo kale alcoloda ,ha u isticmaalin hooyada uurleeyda ah

Over dose daawada oo qiyaasteeda layska badiyo

Malahan case la diiwaan galiyey oo dhibaato ,

Drug intrection daawada aan lala qaadan karin

Hala isticmaalin daawooyinka uurka joojiya contraceptives

Food intrection cuntada saamaynta ku leh

Ma jiro cunto ka hor imaanaysa daawadani

Disease intrection cuduro lid ku ma jiraan
cudurada beerka

Fungus

Drug class: topical anti fungul

Ketoconazole topical

Brand name: nizoral

Ketoconazole :cream

Daawadan waxay ku timaadaa

boomaato cream

kaniini tablets

shambo Shampoo

Indication or principal uses Tilmaanta

Waxaa lagu daweeeyaa qaar kamid ah anti fangal infection
sida infagshanka kudhaca maqaarka

- qawadka tinea
- cabeebka candidiasis
- agoolka dandruff
- seborrhoeisis dermatitis

How to drug work sida daawadu u shaqeeyso

Daawadaan waxa ka hortagtaa jeermiska fungaska
waxayna soo saartaa waxyaabaha nool ee u baahan in la
qashinsaaro

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo isticmaalaa maqaarka waxaana loo
mariyaa 2jeermaalintii mudo 2/6 isbuuc ah

Side effect.. dhibaatooyinka ay leeyihiin

-  lalabo nousea
-  madax xanuun headache
-  dhibaato beerka ah liver problem

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Daawadan waxay bilaawdaa sii dilida jeermiska funguska

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Wad cuntada iyo cabitaanka siina kor dhi cabitaanka

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc
ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato una qaadataa sidii lagu faray

Stopping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato intii dhaqtarku kuu soo qoray

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan lagama badin karo 4 usbuucyo

Precaution: taxadarka

Hala isticmaalin alcol iyo waxyaabaha kale ee aashitada leh

Hana u ismaalin hooyada uurka leh iyo caruurta la nuujinaayo

Over dose daawada oo qiyaasteeda layska badiyo

Malahan case la diiwaan galiyey oo dhibaato ,

Drug intrection daawada aan lala qaadan karin

Majiro daawo la diiwaan galiyey oo aan lala qaadan karin

Food intrection cuntada saamaynta ku leh

Ma jiro cunto ka hor imaanaysa daawadani

Disease intrection cuduro lid ku ma jiraan
cudurka allerjiga

Fungus

Drug class: topical anti fungul

Nystatin

Brand name: mycostatin ,nystex 500.000 unite

Ketoconazole :cream /vaginal tablets cream or ointment

Daawadan waxay ku timaadaa kaniiniyaal hoos lagashanayo
iyo boomaato

Side effect dhibaataada ay keeni karto

Malahan wax hibaato ah oo la xiriirta daawadan

Common dibaato caan ah oo ay keensato daawadan

Malahan wax hibaato ah oo la xiriirta daawadan

Less comman dhibaato yar ayaa lagu arki karaa sida

Lalabo ,matag iyo shuban maqaarka iyo vaginaha oo cuncuna
daaweynta ugu horaysa

Indication or principal uses Tilmaanta

- 1 ku daawee fungus infectioun sida maqaarka afka iyo vagina

How to drug work sida daawadu u shaqeeyso

Daawadaan waxa ka hortagtaa jeermiska fungaska waxayna soo saartaa waxyaabaha nool ee u baahan in la qashinsaaro

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo isticmaalaa maqaarka waxaana loo mariyaa 2/4jeer maalintii mudo 2 isbuuc ah 500.000unite

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Daawadan ma lahan waqti cayiman oo la diiwaangaliyey oo ay bilaabanayso

Duration of action xiliga uu soconkaro

Lama yaqaano

Dietary advice cuntada haboon

Wad cuntada iyo cabitaanka siina kor dhi cabitaanka

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato
una qaadataa sidii lagu faray

Stopping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato
intii dhaqtarku kuu soo qoray

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan lagama badin karo 1/3 usbuucyo

Precaution: taxadarka

Ka ilaali meelaha wasaqada badan ama hawadeedu
wasaqaysan tahay

Hala isticmaalin alcol iyo waxyaabaha kale ee aashitada
leh

Hana u ismaalin hooyada uurka leh iyo caruurta la
nuujinaayo

Over dose daawada oo qiyaasteeda layska badiyo

Lalabo matag iyo shuban ,

Drug intrection daawada aan lala qaadan karin

Majiro daawo la diiwaan galiyey oo aan lala qaadan
karin

Food intrection cuntada saamaynta ku leh

Ma jiro cunto ka hor imaanaysa daawadani

Disease intrection cuduro lid ku ma jiraan

markasta la tasho dhaqtarkaaga

Herpes simplex

Drug class: topical anti viral

Acyclovir

Brand name: zovirax 200mg

Acyclovir : tablet injection ointment liquid

Daawadan waxay ku timaadaa kaniiniyaal boomaato iyo duriin

Side effect dhibaataada ay keeni karto

Malahan wax hibaato ah oo la xiriirta daawadan

Common dibaato caan ah oo ay keensato daawadan

Lalabo matag finan xanuun iyo gubasho cuncun meesha lamariyo

Less comman dhibaato yar ayaa lagu arki karaa sida

Shuban calool xanuun ,wareer, jareyn kaadi badan

Indication or principal uses Tilmaanta

- 1 ku daawee viruska ee ah genital herpes and chicken box

How to drug work sida daawadu u shaqeeyso

Daawadan waxay la howl gashaa enzyme ubaahan in labadalo viruska ku jira unuga gudaha dna si looga hostago jeermiska

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo isticmaalaa maqaarka waxaana loo mariyaa 2/4jeer maalintii mudo 2 isbuuc ah 1200mg 4/8 hours ama 4000 4saacadood ama ilaa todobo saac

Onset of effect xiliga uu bilaawdo dhibaataadiisa

2 saac iyo wax ka badan

Duration of action xiliga uu soconkaro

Shan saac ilaa inta ay soconayso

Dietary advice cuntada haboon

Cuntada waa lagu isticmaali karaa

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato una qaadataa sidii lagu faray

Stoping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato intii dhaqtarku kuu soo qoray

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan lagama badin karo inta lagu qoray waxay keeni kartaa cervical cancer

Precaution: taxadarka

Hala isticmaalin alcol iyo waxyaabaha kale ee aashitada leh

U ismaalin hooyada uurka leh laakiin caruurta la nuujinaayo ha u isticmaalin

Over dose daawada oo qiyaasteeda layska badiyo

Majiro wax ladiiwaan galiyey ,

Drug intrection daawada aan lala qaadan karin

Zidovdine

Food intrection cuntada saamaynta ku leh

Ma jiro cunto ka hor imaanaysa daawadani

Disease intrection cuduro lid ku ma jiraan

wuxuu keeni karaa cudurada beerka iyo kalyo xanuun

Herpes simplex

Drug class: topical anti viral

penciclovir

Brand name: denavir

Penciclovir : topital cream

Daawadan waxay ku timaadaa boomaato

Side effect dhibaataada ay keeni karto

Malahan wax hibaato ah oo la xiriirta daawadan

Common dibaato caan ah oo ay keensato daawadan

Madax xanuun cuncun

Less comman dhibaato yar ayaa lagu arki karaa sida

Kabaabyo finan maqaarka ah

Indication or principal uses Tilmaanta

1 ku daawee viruska ee hargabku keeno

How to drug work sida daawadu u shaqeeyso

Daawadan waxay ku dhagtaa inseym waxaynanadiifiyaan jeermiska

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Daawadan waxaa loo isticmaalaa maqaarka waxaana loo mariyaa 2/4jeer maalintii mudo 2 isbuuc ah 250mg

Onset of effect xiliga uu bilaawdo dhibaataadiisa

2 /3 bilood

Duration of action xiliga uu soconkaro

Lama yaqaano xiliga

Dietary advice cuntada haboon

Cuntada waa lagu isticmaali karaa ama kadib

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato una qaadataa sidii lagu faray

Stoping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato
intii dhaqtarku kuu soo qoray

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan lagama badin karo inta lagu qoray waxay
keeni kartaa dhibaato

Precaution: taxadarka

Hala isticmaalin alcol iyo waxyaabaha kale ee aashitada
leh

Ha u isticmaalin hooyada uurka leh laakiin caruurta la
nuujinaayo ha u isticmaalin

Over dose daawada oo qiyaasteeda layska badiyo

Majiro wax ladiiwaan galiyey ,

Drug intrection daawada aan lala qaadan karin

Phenylbutazone

Food intrection cuntada saamaynta ku leh

Hala isticmaalin koko kolo iyo macmacaaada kale

Disease intrection cuduro lid ku ma jiraan

cudurada kalyaha iyo allerjikada

Genetal warts

Drug class: immunomodulator

imiquimod

Brand name: Aldara

Imiquimod : topical cream

Daawadan waxay ku timaadaa boomaato

Side effect dhibaataada ay keeni karto

Barar indhaha ah ,wajiga ama bushimaha finan iyo shanqar feera ah

Common dhibaato caan ah oo ay keensato daawadan

Gaduudasho maqaarka ah

Less common dhibaato yar ayaa lagu arki karaa sida

Maqaarka la daaweeyey oo adkaada

Indication or principal uses Tilmaanta

1 ku daawee wartska ama meelaha adag

How to drug work sida daawadu u shaqeeyso

Shaqadeeda ma ahan mid la diiwaan galigeey amd machanismka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Mari daawadan maqaarka jiran usbuucii 3 jeer kadibna ku dhaq biyo iyo saabuun

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Lama yaqaano

Duration of action xiliga uu soconkaro

Lama yaqaano xiliga

Dietary advice cuntada haboon

Cunto la aan ayaa la isticmaali karaa

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc ilaa 30 oc ama kayaryahay hana u ogolaan qabow saa id ah

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato una qaadataa sidii lagu faray

Stoping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato
intii dhaqtarku kuu soo qoray oo ah 16 todobaad

Prolonged use hadii daawada oo waqti dheer la qaato

Daawadan lagama badin karo inta lagu qoray waxay
keeni kartaa dhibaato 16 todobaad

Precaution: taxadarka

Daawadani alcolka wax ka hibayso

Daawadani marka la isticmaalayo waa inuu qaraa
dhaqtarka maqaarka waliba hooyada uurleyda ah

Over dose daawada oo qiyaasteeda layska badiyo

Majiro wax ladiiwaan galiyey ,

Drug intrection daawada aan lala qaadan karin

Majiro wax ladiiwaan galiyey ,

Food intrection cuntada saamaynta ku leh

Ma jiro cunto la diiwaan galiyey oo laga ilaalinaayo

Disease intrection cuduro lid ku ma jiraan

ha u isticmaalin dadka qaba cudurada maqaarka

Genetal warts

Drug class: immunomodulator

Interferon alfa/2b

Brand name: intron A

Imiquimod : topical injection

Daawadan waxay ku timaadaa cirbad

Side effect dhibaatada ay keeni karto

Wareer kaadida iyo saxarada oo soo gaduudata ,xabad xanuun ,qandho iyo qarqaryo finan gadgaduudan oo ka soo baxa maqaarka

Common dibaato caan ah oo ay keensato daawadan

Daal iyo muruqyo xanuun ,qandho iyo qarqaryo ,lalabo iyo matag

Less common dhibaato yar ayaa lagu arki karaa sida

Dhabar xanuun ,wareer ,afka oo qalala cuncun maqaarka ah ,kalagoys yada oo xanuuna

Indication or principal uses Tilmaanta

- 1 ku daawee leukemia, sarcoma or genital warts or hepatitis ama qaliinka malignant melanoma.

How to drug work sida daawadu u shaqeeyso

Waxay dhex gashaa oo burburisaa in la soodaayo
unugyada porotiinka waxayna la dagaalantaa in uguyada
viruska kansarka

Dosage =Daawada qiyaas ahaan la siin karo waa sidan

Leukemia 2 million units 3 jeer usbuucii ,hadii uu yahay
kansarka 30 million 3 waqti usbuucii

Onset of effect xiliga uu bilaawdo dhibaataadiisa

Lama yaqaano

Duration of action xiliga uu soconkaro

Lama yaqaano xiliga

Dietary advice cuntada haboon

Cabitaan badan cab si loo yareeyo qatarta hoos u dhaca
wareega dhiiga

Stroke kaydinta daawada

Daawadan ku kaydi meel heerkulkeedu u dhaxeeyo 25oc
ilaa 30 oc ama kayaryahay hana u ogolaan qabow saa id
ah

If you miss dose hadii aad daawada hilmaanto

Waa inaad qaadataa daawadan marka aad soo xasuusato
una qaadataa sidii lagu faray

Stopping the drug

Daawadan waxaad joojinkartaa marka ay kaa dhamaato intii dhaqtarku kuu soo qoray oo ah 16 todobaad

Prolonged use hadii daawada oo waqti dheer la qaato

Markay kuu dhamaato daawadan dhaqtarka ku celi si baaritaan labaad laguugu sameeyo

Precaution: taxadarka

Ka fogow cabitaanka alkoloda iyo hooyada uurleeyda ah

Over dose daawada oo qiyaasteeda layska badiyo

Majiro wax ladiiwaan galiyey ,

Drug intrection daawada aan lala qaadan karin

Daawadan hasiin dadka qaba waalida iyo maskax xanuunka iyo alcol

Food intrection cuntada saamaynta ku leh

Ma jiro cunto la diiwaan galiyey oo laga ilaalinaayo

Disease intrection cuduro lid ku ma jiraan

dadka qaba busbuska ,dadka qaba qalalka ,kalyo xanuunka ,wadno xanuunka