

ECE3103: Probability and Random Processes :
WORKSHEET

Problem 1.

Let X and Y be jointly continuous random variables with joint PDF

$$f_{X,Y}(x,y) = \begin{cases} cx + 1 & x, y \geq 0, x + y < 1 \\ 0 & \text{otherwise} \end{cases}$$

1. Find the constant C
2. Find the marginal PDFs $f_X(x)$ and $f_Y(y)$
3. Find $P(Y < 2X^2)$

Answer:

1.

To find the constant C

$$\begin{aligned} 1 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x,y) dx dy \\ &= \int_0^1 \int_0^{1-x} cx + 1 \, dy dx \\ &= \int_0^1 (cx + 1)(1 - x) \, dx \\ &= \frac{1}{2} + \frac{1}{6}c. \end{aligned}$$

Thus, we conclude $c = 3$

2.

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{XY}(x,y) dy \\ &= \int_0^{1-x} 3x + 1 \, dy \\ &= (3x + 1)(1 - x), \quad \text{for } x \in [0, 1]. \end{aligned}$$

ECE3103: Probability and Random Processes :
WORKSHEET

$$\begin{aligned}f_Y(y) &= \int_{-\infty}^{\infty} f_{XY}(x, y) dx \\&= \int_0^{1-y} (3x + 1) dx \\&= \frac{1}{2}(1-y)(5-3y), \quad \text{for } y \in [0, 1].\end{aligned}$$

$$f_Y(y) = \begin{cases} \frac{1}{2}(1-y)(5-3y) & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned}P(Y < 2X^2) &= \int_{-\infty}^{\infty} \int_{-\infty}^{2x^2} f_{XY}(x, y) dy dx \\&= \int_0^1 \int_0^{\min(2x^2, 1-x)} (3x + 1) dy dx \\&= \int_0^1 (3x + 1) \min(2x^2, 1-x) dx \\&= \int_0^{\frac{1}{2}} 2x^2(3x + 1) dx + \int_{\frac{1}{2}}^1 (3x + 1)(1-x) dx \\&= \frac{53}{96}.\end{aligned}$$

Problem 2.

Let X and Y be jointly continuous random variables with joint PDF

$$f_{X,Y}(x, y) = \begin{cases} 6e^{-(2x+3y)} & x, y \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

1. Are X and Y independent?
2. Find $E[Y | X > 2]$

ECE3103: Probability and Random Processes :
WORKSHEET

3. Find $P(X > Y)$

1. We can write

$$f_X(x) = 2e^{-2x}$$

and

$$f_Y(y) = 3e^{-3y}$$

Thus, X and Y are independent

2. Since X and Y are independent, we have

$$E[Y | X > 2] = E[Y]$$
$$EY = \frac{1}{3}$$

3. We have

$$\begin{aligned} P(X > Y) &= \int_0^{\infty} \int_y^{\infty} 6e^{-(2x+3y)} dx dy \\ &= \int_0^{\infty} 3e^{-5y} dy \\ &= \frac{3}{5}. \end{aligned}$$

Problem 3

Let X and Y be two jointly continuous random variables with joint PDF

$$f_{X,Y}(x,y) = \begin{cases} 6xy & 0 \leq x \leq 1, 0 \leq y \leq \sqrt{x} \\ 0 & \text{otherwise} \end{cases}$$

1.

Find $f_X(x)$ and $f_Y(y)$

2.

Are X and Y independent?

3.

ECE3103: Probability and Random Processes :
WORKSHEET

Find the conditional PDF of X given $Y = y$. $f_{X|Y}(x|y)$

4.

Find $E[X|Y = y]$

5.

Find $\text{Var}(X|Y = y)$

Q.N. 1

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{XY}(x, y) dy \\ &= \int_0^{\sqrt{x}} 6xy dy \\ &= 3x^2. \end{aligned}$$

$$f_X(x) = \begin{cases} 3x^2 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} f_Y(y) &= \int_{-\infty}^{\infty} f_{XY}(x, y) dx \\ &= \int_{y^2}^1 6xy dx \\ &= 3y(1 - y^4). \end{aligned}$$

$$f_Y(y) = \begin{cases} 3y(1 - y^4) & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

2.

X and Y are not independent, since $f_{XY}(x, y) \neq f_X(x)f_Y(y)$

3.

ECE3103: Probability and Random Processes :
WORKSHEET

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

$$= \begin{cases} \frac{2x}{1-y^4} & y^2 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

4.

$$E[X|Y = y] = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx$$

$$= \int_{y^2}^1 x \frac{2x}{1-y^4} dx$$

$$= \frac{2(1-y^6)}{3(1-y^4)}.$$

5.

$$E[X^2|Y = y] = \int_{-\infty}^{\infty} x^2 f_{X|Y}(x|y) dx$$

$$= \int_{y^2}^1 x^2 \frac{2x}{1-y^4} dx$$

$$= \frac{1-y^8}{2(1-y^4)}.$$

$$\text{Var}(X|Y = y) = E[X^2|Y = y] - (E[X|Y = y])^2$$

$$= \frac{1-y^8}{2(1-y^4)} - \left(\frac{2(1-y^6)}{3(1-y^4)} \right)^2.$$

Problem: 4

Let X and Y be two jointly continuous random variables with joint PDF

$$f_{XY}(x, y) = \begin{cases} 2 & y + x \leq 1, x > 0, y > 0 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Cov}(X, Y)$

ECE3103: Probability and Random Processes : **WORKSHEET**

- Solution
 - For $0 \leq x \leq 1$ $0 \leq x \leq 1$, we have

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{XY}(x, y) dy \\ &= \int_0^{1-x} 2 dy \\ &= 2(1-x). \end{aligned}$$

Thus,

$$f_X(x) = \begin{cases} 2(1-x) & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Similarly, we obtain

$$f_Y(y) = \begin{cases} 2(1-y) & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} EX &= \int_0^1 2x(1-x) dx \\ &= \frac{1}{3} = EY, \end{aligned}$$

$$\begin{aligned} EX^2 &= \int_0^1 2x^2(1-x) dx \\ &= \frac{1}{6} = EY^2. \end{aligned}$$

Thus,

$$\text{Var}(X) = \text{Var}(Y) = \frac{1}{18}.$$

ECE3103: Probability and Random Processes :
WORKSHEET

$$\begin{aligned} E_{XY} &= \int_0^1 \int_0^{1-x} 2xy \, dy \, dx \\ &= \int_0^1 x(1-x)^2 \, dx \\ &= \frac{1}{12}. \end{aligned}$$

$$\begin{aligned} \text{Cov}(X, Y) &= E_{XY} - E_X E_Y \\ &= \frac{1}{12} - \left(\frac{1}{3}\right)^2 \\ &= -\frac{1}{36}, \end{aligned}$$

Problem The random variable X has a range of $\{0, 1, 2\}$ and the random variable Y has a range of $\{1, 2\}$. The joint distribution of X and Y is given by the following table:

x	y	$P(X = x, Y = y)$
0	1	0.2
0	2	0.1
1	1	0.0
1	2	0.2
2	1	0.3
2	2	0.2

- Write down tables for the marginal distributions of X and of Y , i.e. give the values of $P(X = x)$ for all x , and of $P(Y = y)$ for all y .

x	$P(X = x)$
0	0.3
1	0.2
2	0.5

y	$P(Y = y)$
1	0.5
2	0.5

ECE3103: Probability and Random Processes :

WORKSHEET

2. Write down a table for the conditional distribution of X given that $Y = 2$, i.e. give the values of $P(X = x | Y = 2)$ for all x .

x	$P(X = x Y = 2)$
0	0.2
1	0.4
2	0.4

3. Compute $E(X)$ and $E(Y)$.

$$E(X) = 0 \times 0.3 + 1 \times 0.2 + 2 \times 0.5 = 1.2$$

$$E(Y) = 1 \times 0.5 + 2 \times 0.5 = 1.5$$

4. Compute $E(XY)$.

$$\begin{aligned} E(XY) &= (0 \times 1) \times 0.2 + (0 \times 2) \times 0.1 + (1 \times 1) \times 0.0 + (1 \times 2) \times 0.2 + (2 \times 1) \times 0.3 + (2 \times 2) \times 0.2 \\ &= 1.8 \end{aligned}$$

5. Are X and Y independent? Explain why or why not.

No, they are not independent, as can be seen from the fact that $P(X = 1, Y = 1) \neq P(X = 1)P(Y = 1)$. (The left side is 0, but the right side is $0.2 \times 0.5 = 0.1$.)

Note that the fact that $E(XY) = E(X)E(Y)$ does NOT imply that X and Y are independent. $E(XY) = E(X)E(Y)$ is implied by X and Y being independent, but not the other way around.

Problem

You roll one red die and one green die. Define the random variables X and Y as follows:

X = The number showing on the red die

Y = The number of dice that show the number six

For example, if the red and green dice show the numbers 6 and 4, then $X = 6$ and $Y = 1$.

Write down a table showing the joint probability mass function for X and Y , find the marginal distribution for Y , and compute $E(Y)$.

ECE3103: Probability and Random Processes : **WORKSHEET**

Here is a table showing the joint probability mass function, with the marginal distribution for Y on the right:

		$X =$						
		1	2	3	4	5	6	
$Y =$	0	$5/36$	$5/36$	$5/36$	$5/36$	$5/36$	0	$25/36$
	1	$1/36$	$1/36$	$1/36$	$1/36$	$1/36$	$5/36$	$10/36$
	2	0	0	0	0	0	$1/36$	$1/36$

The expectation of Y is

$$E(Y) = 0 \times 25/36 + 1 \times 10/36 + 2 \times 1/36 = 12/36 = 1/3$$

Question 3. Suppose you roll two fair, six-sided dice, one of which is red and the other of which is green. Define the following random variables:

X = The number shown on the red die

$$Y = \begin{cases} 0 & \text{if the two dice show the same number} \\ 1 & \text{if the number on the green die is bigger than the number on the red die} \\ 2 & \text{if the number on the red die is bigger than the number on the green die} \end{cases}$$

a) Write down a table showing the joint probability mass function for X and Y .

		$X =$					
		1	2	3	4	5	6
$Y =$	0	1/36	1/36	1/36	1/36	1/36	1/36
	1	5/36	4/36	3/36	2/36	1/36	0
	2	0	1/36	2/36	3/36	4/36	5/36

ECE3103: Probability and Random Processes :
WORKSHEET

b) Find the marginal probability mass function for Y , and compute its expected value.

y	$P(Y = y)$
0	6/36
1	15/36
2	15/36

$$E(Y) = 0 \times (6/36) + 1 \times (15/36) + 2 \times (15/36) = 45/36$$

c) Find the conditional probability mass function for X given $Y = 1$.

$X =$						
1	2	3	4	5	6	
5/15	4/15	3/15	2/15	1/15	0	

Correlation Coefficient:

$$\begin{aligned}\rho_{XY} &= \text{Cov}(U, V) = \text{Cov}\left(\frac{X - EX}{\sigma_X}, \frac{Y - EY}{\sigma_Y}\right) \\ &= \text{Cov}\left(\frac{X}{\sigma_X}, \frac{Y}{\sigma_Y}\right) \\ &= \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y}.\end{aligned}$$

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

Problem: 5

Let X be a continuous random variable with PDF

$$f_X(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

ECE3103: Probability and Random Processes : WORKSHEET

We know that given $X = x$ the random variable Y is uniformly distributed on $[-x, x]$

1. Find the joint PDF $f_{XY}(x, y)$
2. Find $f_Y(y)$
3. Find $P(|Y| < X^3)$

1. First note that, by the assumption

$$f_{Y|X}(y|x) = \begin{cases} \frac{1}{2x} & -x \leq y \leq x \\ 0 & \text{otherwise} \end{cases}$$

Thus, we have

$$f_{XY}(x, y) = f_{Y|X}(y|x)f_X(x) = \begin{cases} 1 & 0 \leq x \leq 1, -x \leq y \leq x \\ 0 & \text{otherwise} \end{cases}$$

Thus,

$$f_{XY}(x, y) = \begin{cases} 1 & |y| \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

2. First, note that $R_Y = [-1, 1]$. To find $f_Y(y)$ we can write

$$\begin{aligned} f_Y(y) &= \int_{-\infty}^{\infty} f_{XY}(x, y) dx \\ &= \int_{|y|}^1 1 dx \\ &= 1 - |y|. \end{aligned}$$

Thus,

$$f_Y(y) = \begin{cases} 1 - |y| & |y| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

ECE3103: Probability and Random Processes : WORKSHEET

3. To find $P(|Y| < X^3)$, we can use the law of total probability

$$\begin{aligned} P(|Y| < X^3) &= \int_0^1 P(|Y| < X^3 | X = x) f_X(x) dx \\ &= \int_0^1 P(|Y| < x^3 | X = x) 2x dx \\ &= \int_0^1 \left(\frac{2x^3}{2x} \right) 2x dx \quad \text{since } Y | X = x \sim \text{Uniform}(-x, x) \\ &= \frac{1}{2}. \end{aligned}$$

Problem: 6

Let the probability density function of X be given by

$$f(x) = \begin{cases} 6x(1-x), & 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

Now find the probability density of $Y = X^3$.

Let $G(y)$ denote the value of the distribution function of Y at y and write

$$\begin{aligned} G(y) &= P(Y \leq y) \\ &= P(X^3 \leq y) \\ &= P(X \leq y^{1/3}) \\ &= \int_0^{y^{1/3}} 6x(1-x) dx \\ &= \int_0^{y^{1/3}} (6x - 6x^2) dx \\ &= (3x^2 - 2x^3) \Big|_0^{y^{1/3}} \\ &= 3y^{2/3} - 2y \end{aligned}$$

Now differentiate $G(y)$ to obtain the density function $g(y)$

$$\begin{aligned} g(y) &= \frac{dG(y)}{dy} \\ &= \frac{d}{dy} (3y^{2/3} - 2y) \\ &= 2y^{-1/3} - 2 \\ &= 2(y^{-1/3} - 1), \quad 0 < y < 1 \end{aligned}$$

ECE3103: Probability and Random Processes :
WORKSHEET

Problem:7

Let X and Y be two jointly continuous random variables with joint PDF

$$f_{XY}(x, y) = \begin{cases} 2 & y + x \leq 1, x > 0, y > 0 \\ 0 & \text{otherwise} \end{cases}$$

Find $\text{Cov}(X, Y)$ and $\rho(X, Y)$

- Solution

- For $0 \leq x \leq 1$ $0 \leq x \leq 1$, we have

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{XY}(x, y) dy \\ &= \int_0^{1-x} 2 dy \\ &= 2(1-x). \end{aligned}$$

Thus,

$$f_X(x) = \begin{cases} 2(1-x) & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Similarly, we obtain

$$f_Y(y) = \begin{cases} 2(1-y) & 0 \leq y \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

Thus, we have

$$\begin{aligned} EX &= \int_0^1 2x(1-x) dx \\ &= \frac{1}{3} = EY, \end{aligned}$$

$$\begin{aligned} EX^2 &= \int_0^1 2x^2(1-x) dx \\ &= \frac{1}{6} = EY^2. \end{aligned}$$

ECE3103: Probability and Random Processes :
WORKSHEET

Thus,

$$\text{Var}(X) = \text{Var}(Y) = \frac{1}{18}.$$

$$\begin{aligned} E_{XY} &= \int_0^1 \int_0^{1-x} 2xy \, dy \, dx \\ &= \int_0^1 x(1-x)^2 \, dx \\ &= \frac{1}{12}. \end{aligned}$$

$$\begin{aligned} \text{Cov}(X, Y) &= E_{XY} - E_X E_Y \\ &= \frac{1}{12} - \left(\frac{1}{3}\right)^2 \\ &= -\frac{1}{36}, \end{aligned}$$

$$\begin{aligned} \rho(X, Y) &= \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= -\frac{1}{2}. \end{aligned}$$

Properties:

The covariance has the following properties:

1. $\text{Cov}(X, X) = \text{Var}(X)$
2. if X and Y are independent then $\text{Cov}(X, Y) = 0$
3. $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
4. $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$
5. $\text{Cov}(X + c, Y) = \text{Cov}(X, Y)$
6. $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$

ECE3103: Probability and Random Processes :
WORKSHEET

Let X and Y be two independent $N(0, 1)$ random variables and

$$Z = 1 + X + XY^2,$$

$$W = 1 + X.$$

Find $\text{Cov}(Z, W)$

- Solution

◦

$$\begin{aligned}\text{Cov}(Z, W) &= \text{Cov}(1 + X + XY^2, 1 + X) \\ &= \text{Cov}(X + XY^2, X) \\ &= \text{Cov}(X, X) + \text{Cov}(XY^2, X) \\ &= \text{Var}(X) + E[X^2Y^2] - E[XY^2]E[X] \\ &= 1 + E[X^2]E[Y^2] - E[X]^2E[Y^2] \\ &= 1 + 1 - 0 = 2.\end{aligned}$$

Problem: 8

A random Variable defined as $X(t) = A \cos(\omega t) + B \sin(\omega t)$, where A and B are random variables with density functions $f_A(a)$ and $f_B(b)$. We have to find the conditions under which $X(t)$ will be stationary:

1. Mean:

$$E[X(t)] = E[A \cos(\omega t) + B \sin(\omega t)] = \cos(\omega t)E[A] + \sin(\omega t)E[B]$$

If $E[X(t)]$ is to be independent of t , then $E[A] = E[B] = 0$, in which case $E[X(t)] = 0$.

ECE3103: Probability and Random Processes : **WORKSHEET**

2. Autocorrelation:

$$\begin{aligned} R_X(t_1, t_2) &= E[X(t_1)X(t_2)] = E\{[A \cos(\omega t_1) \\ &\quad + B \sin(\omega t_1)][A \cos(\omega t_2) + B \sin(\omega t_2)]\} \\ &= E\{A^2 \cos(\omega t_1) \cos(\omega t_2) + B^2 \sin(\omega t_1) \sin(\omega t_2) \\ &\quad + AB[\sin(\omega t_1) \cos(\omega t_2) + \cos(\omega t_1) \sin(\omega t_2)]\} \\ &= \frac{1}{2}E[A^2][\cos(\omega(t_2 - t_1)) + \cos(\omega(t_2 + t_1))] \\ &\quad + \frac{1}{2}E[B^2][\cos(\omega(t_2 - t_1)) - \cos(\omega(t_2 + t_1))] \\ &\quad + E[AB] \sin(\omega(t_2 + t_1)) \end{aligned}$$

The conditions under which this equation will be dependent only on $(t_2 - t_1)$ are $E[A^2] = E[B^2]$ and $E[AB] = 0$. In this case

$$R_X(t_1, t_2) = E[A^2][\cos(\omega(t_2 - t_1))] = E[A^2][\cos(\omega\tau)]$$

We can now summarize the conditions for stationarity:

- (a) $E[A] = E[B] = 0$
- (b) $E[A^2] = E[B^2]$
- (c) $E[AB] = 0$

Problem 9

Let a random variable $X(t)$ defined as $X(t) = A \sin(\omega_0 t + \Phi)$, where A and Φ are independent and A is uniformly distributed over $(0,1)$ and Φ is uniformly distributed over $(0, 2\pi)$ and find whether it is ergodic or not

ECE3103: Probability and Random Processes : **WORKSHEET**

We have already seen in Example 19.2.1 that

$$E[X(t)] = E[A]E[\sin(\omega_0 t + \Phi)] = 0$$

$$\begin{aligned} R_X(\tau) &= E[X(t)X(t + \tau)] = E[A^1]E[\sin(\omega_0 t + \Phi) \sin(\omega_0 t + \omega_0 \tau + \Phi)] \\ &= \frac{1}{2} E[A^2] \cos(\omega_0 \tau) \end{aligned}$$

and $X(t)$ is indeed stationary. In fact, it is also strict-sense stationary. We will check first whether it is mean-ergodic. Since the mean value is zero, $C_X(\tau) = \frac{1}{2} E[A^2] \cos(\omega_0 \tau)$. The variance of the time average $\sigma_{\hat{\mu}_X}^2$ is given by

$$\sigma_{\hat{\mu}_X}^2 = \frac{E[A^2]}{2T} \left\{ \int_{-2T}^{2T} \left[1 - \frac{|\tau|}{2T} \right] \frac{\cos(\omega_0 \tau)}{2} d\tau \right\}$$

We can use the result from Fourier transform that time multiplication is frequency convolution and evaluate the integral shown above. Substituting the Fourier transforms of the triangular and the cosine functions

$$\left[1 - \frac{|\tau|}{2T} \right] \Longleftrightarrow 2T \left[\frac{\sin(\omega T)}{\omega T} \right]^2 \quad \text{and} \quad \frac{\cos(\omega_0 \tau)}{2} \Longleftrightarrow \frac{\pi}{2} [\delta(\omega + \omega_0) + \delta(\omega - \omega_0)]$$

in the equation for the variance, we obtain

$$\begin{aligned} \sigma_{\hat{\mu}_X}^2 &= \frac{1}{2\pi} \frac{E[A^2]}{2T} \int_{-\infty}^{\infty} 2T \left[\frac{\sin(\omega T)}{\omega T} \right]^2 \frac{\pi}{2} [\delta(\omega + \omega_0) + \delta(\omega - \omega_0)] d\omega \\ &= \frac{E[A^2]}{2\pi} \left[\frac{\sin(\omega_0 T)}{\omega_0 T} \right]^2 \end{aligned}$$

and as $T \rightarrow \infty$, $\sigma_{\hat{\mu}_X}^2 \rightarrow 0$, for $\omega_0 \neq 0$. Thus $X(t)$ is mean-ergodic.

Substituting $E[A^2] = \frac{1}{3}$ in the variance equation, we obtain

$$\sigma_{\hat{\mu}_X}^2 = \frac{1}{6\pi} \left[\frac{\sin(\omega_0 T)}{\omega_0 T} \right]^2$$

To check for power ergodicity, it is simpler to use Eq. (19.3.16) rather than Eq. (19.3.17). Thus

$$\lim_{T \rightarrow \infty} \left[\frac{1}{2T} \int_{-T}^T X^2(t) dt \right] = \lim_{T \rightarrow \infty} \left[\frac{1}{2T} \int_{-T}^T A^2 \cos^2(\omega_0 t + \Phi) dt \right] = \frac{A^2}{2} \quad \text{for } \omega_0 \neq 0$$

Since this expression does not converge to $R_X(0) = \frac{1}{6}$, the process $X(t)$ is neither power-ergodic nor correlation-ergodic.

Problem 10

consider a random process $X(t) = B + A \sin(\omega_0 t + \Phi)$, where B , A , and Φ are independent random variables. A and B are uniformly distributed over $(0,1]$ and $(0,2]$ respectively. We will check whether $X(t)$ is mean-ergodic.

ECE3103: Probability and Random Processes : **WORKSHEET**

We first determine whether $X(t)$ is stationary. The mean value μ_X is given by

$$E[B + A \sin(\omega_0 t + \Phi)] = \mu_B + E[A \sin(\omega_0 t + \Phi)] = \mu_B$$

since $E[A \sin(\omega_0 t + \Phi)] = 0$ from previous examples. The autocorrelation function is

$$\begin{aligned} R_X(\tau) &= E\{[B + A \sin(\omega_0 t + \Phi)][B + A \sin(\omega_0 t + \omega_0 \tau + \Phi)]\} \\ &= E[B^2] + 0 + 0 + E[A^2]E[\sin(\omega_0 t + \Phi)A \sin(\omega_0 t + \omega_0 \tau + \Phi)] \\ &= E[B^2] + E[A^2] \frac{1}{2} E[\cos(\omega_0 \tau) - \cos(2\omega_0 t + \omega_0 \tau + 2\Phi)] \\ &= E[B^2] + \frac{1}{2} E[A^2] \cos(\omega_0 \tau) \end{aligned}$$

Hence the process $X(t)$ is stationary. The autocovariance function $C_X(\tau)$ is given by

$$\begin{aligned} C_X(\tau) &= R_X(\tau) - \mu_X^2 = E[B^2] + \frac{1}{2} E[A^2] \cos(\omega_0 \tau) - \mu_B^2 \\ &= \sigma_B^2 + \frac{1}{2} E[A^2] \cos(\omega_0 \tau) \end{aligned}$$

and substituting this equation in Eq. (19.3.7), we obtain the variance of the time average as

$$\begin{aligned} \sigma_{\hat{\mu}_X}^2 &= \frac{1}{2T} \left\{ \int_{-2T}^{2T} \left[\sigma_B^2 + \frac{1}{2} E[A^2] \cos(\omega_0 \tau) \right] \left[1 - \frac{|\tau|}{2T} \right] d\tau \right\} \\ &= \frac{1}{2T} \left[2T\sigma_B^2 + E[A^2] \frac{2T}{2\pi} \left[\frac{\sin(\omega_0 T)}{\omega_0 T} \right]^2 \right] = \sigma_B^2 + \frac{E[A^2]}{2\pi} \left[\frac{\sin(\omega_0 T)}{\omega_0 T} \right]^2 \end{aligned}$$

and

$$\lim_{T \rightarrow \infty} \sigma_{\hat{\mu}_X}^2 = \lim_{T \rightarrow \infty} \left\{ \sigma_B^2 + \frac{E[A^2]}{2\pi} \left[\frac{\sin(\omega_0 T)}{\omega_0 T} \right]^2 \right\} = \sigma_B^2 \quad \text{if } \omega_0 \neq 0$$

We find that even though $X(t)$ is stationary, it is not mean-ergodic. Hence the time average is not a good estimator of the ensemble average. We can now substitute $E[A^2] = \frac{1}{3}$ and $\sigma_B^2 = \frac{1}{3}$ in the variance equation and write

$$\sigma_{\hat{\mu}_X}^2 = \sigma_B^2 + \frac{E[A^2]}{2\pi} \left[\frac{\sin(\omega_0 T)}{\omega_0 T} \right]^2 = \frac{1}{3} \left[1 + \frac{\sin(\omega_0 T)}{2\pi\omega_0 T} \right]$$

Problem 11

A random process $Y(t)$ is given by $Y(t) = X(t) \cos(\omega t + \Phi)$, where $X(t)$, a zero mean wide-sense stationary random process with autocorrelation function $R_X(\tau) = 2e^{-2\lambda|\tau|}$ is modulating the carrier $\cos(\omega t + \Phi)$. The random variable Φ is uniformly distributed in the interval $(0, 2\pi)$, and is independent of $X(t)$. to find the mean, variance, and autocorrelation of $Y(t)$:

ECE3103: Probability and Random Processes : **WORKSHEET**

Mean. The independence of $X(t)$ and Φ allows us to write

$$E[Y(t)] = E[X(t)]E[\cos(\omega t + \Phi)]$$

and with $E[X(t)] = 0$ and $E[\cos(\omega t + \Phi)] = 0$ from Example 19.2.1, $E[Y(t)] = 0$.

Variance. Since $X(t)$ and Φ are independent, the variance can be given by

$$\sigma_Y^2 = E[Y^2(t)] = E[X^2(t) \cos^2(\omega t + \Phi)] = \sigma_X^2 E[\cos^2(\omega t + \Phi)]$$

However

$$E[\cos^2(\omega t + \Phi)] = \frac{1}{2} E[1 + \cos(2\omega t + 2\Phi)] = \frac{1}{2} \quad \text{and} \quad \sigma_X^2 = C_X(0) = R_X(0) = 2$$

and hence $\sigma_Y^2 = \sigma_X^2/2 = 1$.

Autocorrelation:

$$\begin{aligned} R_Y(\tau) &= E[Y(t)Y(t+\tau)] = E[X(t) \cos(\omega t + \Phi) X(t+\tau) \cos(\omega t + \omega\tau + \Phi)] \\ &= R_X(\tau) \frac{1}{2} E[\cos(\omega\tau) + \cos(2\omega t + \omega\tau + 2\Phi)] \\ &= \frac{R_X(\tau)}{2} \cos(\omega\tau) + \frac{R_X(\tau)}{2} E[\cos(2\omega t + \omega\tau + 2\Phi)] \end{aligned}$$

From Example 19.2.1 $E[\cos(2\omega t + \omega\tau + 2\Phi)] = 0$, and hence


$$R_Y(\tau) = \frac{R_X(\tau)}{2} \cos(\omega\tau) = e^{-2\lambda|\tau|} \cos(\omega\tau)$$

A graph of $R_Y(\tau)$ is shown in Fig. 19.2.7 with $\lambda = 0.5$ and $\omega = 2\pi$.

ECE3103: Probability and Random Processes : WORKSHEET

Problem 12

Consider a Gaussian random process $X(t)$ with autocorrelation function::



- Find the total average power, $E(X^2)$. $E(X^2) = R_X(0) = 4$
- $\text{var}(X) = E(X^2) - (E(X))^2 = 4 - 2 = 2$
- $E(X(3) \cdot X(4)) = R_X(t_1 = 3, t_2 = 4) = R_X(\tau = 1) = 3$ (from the graph, by inspection)

Problem 13

A wide-sense stationary (WSS) random process $X(t)$ has autocorrelation function

$$R_X(\tau) = 25 \exp(-4|\tau|) + 16 \cos(20\tau) + 36$$

- $E[X^2(t)] = R_X(0) = 25 + 16 + 36 = 77$
- $E[X(t)] = \pm 6$
- $\text{var}(X(t)) = E[X^2(t)] - (E(X))^2 = 77 - 36 = 41$

Problem: 14

Random variables X and Y have joint pdf

$$f_{XY}(x,y) = \begin{cases} 4xy & 0 \leq x \leq 1, 0 \leq y \leq 1 \\ 0 & \text{else} \end{cases}$$

Note: The joint pdf factors to give

$$\left. \begin{aligned} f_X(x) &= 2x \quad 0 \leq x \leq 1, \\ f_Y(y) &= 2y \quad 0 \leq y \leq 1 \end{aligned} \right\} \Rightarrow X, Y \text{ independent, identically distributed (IID)}$$

$$\text{a.} \quad E(X) = E(Y) = 2 \int_0^1 x^2 dx = \frac{2}{3} x^3 \Big|_{x=0}^1 = 2/3$$

$$\text{b.} \quad E(X^2) = E(Y^2) = 2 \int_0^1 x^3 dx = \frac{2}{4} x^4 \Big|_{x=0}^1 = \frac{1}{2}$$

$$\text{Var}(X) = \text{var}(Y) = E(X^2) - (E(X))^2 = (1/2) - 4/9 = 1/18$$

$$\text{c.} \quad \text{cov}(X, Y) = 0, \text{ since independent} \Rightarrow \text{uncorrelated}$$

$$E(X + Y) = E(X) + E(Y) = (2/3) + (2/3) = 4/3$$

$$\text{d.} \quad \text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y) = (1/18) + (1/18) + 0 = 1/9$$

Final Exam Questions

1. (a) Explain the term central limit theory (CLT)
(b) Define the term random process and random variable.
(c) Define monotonic and non-monotonic function.
(d) What are the conditions for any stochastic process will be Ergodic process.

2. A discrete random variable X and Y has joint PMF which is given **by**

$$p(x_i, y_j) = \begin{cases} k(4x_i + 2y_j), & x_i = 2, 3 \text{ and } y_j = 2, 3 \\ 0, & \text{otherwise} \end{cases}$$

Determine

- a. The constant K values
 - b. The marginal PMF of X and Y
 - c. The conditional PMF of X and Y
 - d. Are X and Y independent?
3. Let X and Y are two jointly distributed continuous random variables representing voltages measured in Volts at two different nodes of a circuit. X has the marginal density function

$$f(x) = \begin{cases} \frac{2x+1}{2}, & 0 \text{ volt} \leq x \leq 1 \text{ volts} \\ 0, & \text{otherwise} \end{cases}$$

You are also given the conditional density function

$$f(y|x) = \begin{cases} \frac{2(x+y)}{2x+1}, & 0 \text{ volt} \leq x \leq 1 \text{ volt and } 0 \text{ volt} \leq y \leq 1 \text{ volt} \\ 0, & \text{otherwise} \end{cases}$$

- a. Compute $P(Y \leq 0.5 \text{ volts} \mid X = 0.5 \text{ volts})$
 - b. Compute $P(0.25 \text{ volts} \leq X \leq 0.5 \text{ volts}, Y \leq 0.5 \text{ volts})$
 - c. Determine Mean of X and Y.
4. Let X has PMF given **by**

$$f_X(x_i) = \begin{cases} \frac{2}{3}, & x = 1 \\ \frac{1}{3}, & x = 2 \end{cases}$$

If $Y = X^2 + 1$, Find

- a. PMF of Y
- b. $E[Y]$

5. Two random variables X and Y have joint probability density function as

$$f_{X,Y}(x,y) = \begin{cases} 2 - x - y, & 0 < x < 1, \quad 0 < y < 1 \\ 0, & \text{otherwise} \end{cases}$$

Find Covariance of X and Y.

6. Let X be uniformly distributed random variable on [2, 4] and $Y = e^{2X}$

- Compute the CDF of Y.
- Compute pdf of Y.
- Find the mean of Y.

7. Consider the random process $X(t) = B \cos(2\pi t)$, random variable B follows the exponential distribution as $f_B(b) = 2e^{-2b}$ where $b > 0$, the mean and variance of B are $\frac{1}{2}$ and $\frac{1}{4}$ respectively. Find

- The mean value of $X(t)$.
- The autocorrelation function of the random process X (t).
- Is the random process X (t) wide-sense stationary (WSS)? If yes or no, explain.

