Question 1

A 16-year-old student presents to the clinic for a sport physical, she is accompanied by her mother. She is involved in gymnastics and has been since age 5. In addition to training for the gymnastics team, she runs 10 miles per day. She is proud of her athleticism and states that she is still too fat and wants to lose more weight. You note that her BMI is 15. She denies eating excessively or making herself vomits. The patients mother states she barely eats. This patient is at risk for which medical problem?

Select one:

a. Anorexia nervosa

b. Bulimia

c. Not at risk; normal adolescence

d. Obesity

Question 2

A 17 years old boy presented with a low grade fever and pain in the right lower quadrant. On palpation he has tenderness of the left lower quadrant of the abdomen. What is the most likely diagnosis

Select one:

a. Irritable bowel syndrome

b. Acute appendicitis

c. Psychogenic pain

d. Crohn disease

Question 3

A 23-year old patient was admitted to surgical ward after sustaining major trauma to the chest and abdomen following road traffic accident of 02 days. After insertion of chest tube, laparotomy was decided and the patient was kept NPO then after. Which of the following is the primary source of energy in this patient?

Select one:

a. Protein (muscle)

b. Fat

c. Ketone bodies

d. Glycogen

Question 4

A 25-year-old man was involved in a fight and had sustained machete injury to his scalp a week back. He was brought to the hospital because he developed generalized spasms since 24 hours. Physical examination revealed awake patient with a 4cm-sutured wound over the scalp with palpable depressed parietal bone underneath. He has weakness of his left upper limb. What is the diagnosis

Select one:

a. Epidural hematoma

b. Subdural hematoma

c. Tetanus

d. Increased ICP

Question 5

A 26 year old male presented to the emergency OPD suffering from severe allergic reaction to a bee sting. What is the most appropriate initial therapy

Select one:

a. Albuterol

b. Cimetidine

c. Nor epinephrine

d. Epinephrine

Question 6

A 29 years old female epileptic patient is brought to OPD seizing for more than 10 minutes and she was still having seizure.What is the first line of treatmentof this patient?

Select one:

a. Intravenous pentobarbital

b. Intravenous diazepam

c. Intravenous Paraldehyde

d. Intravenous phenytoin

Question 7

A 29 years old man presented with low grade fever, loss of appetite, weight loss, night sweating and cough productive of sputum of one month duration. Which one is the gold standard diagnostic test for this patients condition

Select one:

a. PPD test

b. Chest X-ray

c. ESR

d. AFB test

Question 8

A 30 year old man presented with Parkinsonian symptoms. The most probable cause is

Select one:

a. Parkinson disease

b. Drug/toxin induced

c. Idiopathic

d. Stroke

Question 9

A 30-year old patient develops high-grade fever on the fifth post laparotomy day. What is the most likely cause of the fever

Select one:

a. Surgical site infection

b. Metabolic crisis

c. Atelectasis

d. Malignant hyperthermia

Question 10

A 32-year-old female patient presented 4 hours after she sustained a human bite to her forearm. Physical examination revealed a 4cm linear laceration over her mid right forearm. What is the appropriate management

Select one:

a. Primary closure

b. Delayed primary closure

c. Secondary closure

d. Leave it to heal by secondary intention

Question 11

A 35-year-old otherwise healthy woman went into shock 6 hours after difficult cholecystectomy. What is the most likely cause of the shock

Select one:

a. Hemorrhage

b. Sepsis

c. Bowel perforation

d. Bile leak

Question 12

A 38-year-old daily laborer presented with right leg pain & swelling of 2 days duration. He has no history of trauma. Physical examination revealed BP=90/50mmHg, PR= 136/min, RR=28/min, TO= 35.4OC. His right leg is grossly swollen with blisters on the overlying skin, 2cm ulcer over the lateral malleolus discharging dishwater fluid. What is the appropriate definitive management of this patient?

Select one:

a. IV antibiotics & amputation

b. IV antibiotics & radical debridement

c. PO antibiotics & follow up

d. Antibiotic according to culture & sensitivity

Question 13

A 4 -year old female child presented initially with vaginal bleeding but later she also started to have foul smelling vaginal discharge for the last one month, no bleeding from other sites. The most likely cause of the vaginal bleeding is

Select one:

a. Neoplasm of the vagina

b. Urethral prolapse

c. Vulvovaginitis

d. Foreign body in the vagina

Question 14

Answer saved

Marked out of 1.00

Flag question

Question text

A 40 year old man presented with weakness of the upper and lower extremities of 1 year duration. Physical examination revealed predominantly distal weakness with absent reduced deep tender reflexes. He is most likely suffering from

Select one:

a. Myopathy

b. Upper motor neuron lesion

c. Neuropathy

d. Stroke

Question 15

Answer saved

Marked out of 1.00

Flag question

Question text

A 44-year old patient underwent a major surgical procedure, which resulted in significant blood loss. The patient was transferred with stable vital signs to the recovery where he was started on transfusion of cross-matched blood. Few minutes later, the patient started complaining of nausea, vomiting, fever, chills, chest and flank pain. On examination: BP=80/50mmHg, PR=109bpm, RR=35/min, T0=38. 90C. The urine appeared reddish in color. What is the most likely diagnosis?

Select one:

a. Delayed hemolytic transfusion reaction

b. Allergic transfusion reaction

c. Febrile non hemolytic transfusion reaction

d. Acute hemolytic transfusion reaction

Question 16

Answer saved

Marked out of 1.00

Remove flag

Question text

A 45-year-old man with severe non-bilious vomiting for 4 months presented to the OPD with marked dehydration. His urine output was 20ml/h, and the hematocrit is 48%. What is the predominant metabolic abnormality

Select one:

a. Hypochloremic alkalosis

b. Respiratory alkalosis

c. Metabolic acidosis

d. Intrinsic renal disease

Question 17

Answer saved

Marked out of 1.00

Remove flag

Question text

A 45-year-old man with severe non-bilious vomiting for 4 months presented to the OPD with marked dehydration. His urine output was 20ml/h, and the hematocrit is 48%. Initial treatment for this patient should include which of the following

Select one:

a. Normal saline with IV potassium

b. Antiemetic with NG tube insertion

c. Ringer lactate solution

d. Administration of DNS with IV potassium

Question 18

Answer saved

Marked out of 1.00

Remove flag

Question text

A 5 years old child sustained a fall down accident from a five meters tree. On arrival to emergency unit, his GCS 8/15. While on evaluation which technique do you use to open the airway

Select one:

a. Chin lift

b. Nose up

c. Jaw thrust

d. Neutral Position

Question 19

Answer saved

Marked out of 1.00

Remove flag

Question text

A 50 Years old woman came with a breast lamp and FNAC revealed breast cancer. Which one of the following can be considered as significant risk factor for this woman

Select one:

a. A history of using birth control pills

b. A history of endometrial carcinoma

c. A high fiber, low fat diet

d. Late menarche

Question 20

Answer saved

Marked out of 1.00

Flag question

Question text

A 55 years old man was diagnosis to have diabetes mellitus 6 months back. He was put on metformin and his blood glucose level is controlled. Which of the following least characterizes the diseases this patient is having?

Select one:

a. Genetic predisposition

b. Increased visceral fat

c. More insidious onset

d. Low basal insulin secretion early in the disease

Almaz comes to your clinic for advice about breast examination. What is the correct way of instructing about the self-breast examination?

Select one:

a. Recommend that the best time to perform BSE is Four to Seven days after the first day of the menstrual period

b. Tell the woman that if she is pregnant, she should not perform BSE until the baby is born

c. Urge the woman that she needs to do BSE bimonthly unless she has fibrocystic breast tissue

d. Tell the woman that the best time to perform BSE is in the middle of the menstrual cycle

Question 22

Answer saved

Marked out of 1.00

Flag question

Question text

A 6 month old infant was diagnosed with congenital rubella. Which of the following is the least likely presentation for her?

Select one:

a. Low set ears

b. Cataract

c. Microcephaly

d. Congenital heart disease

Question 23

Answer saved

Marked out of 1.00

Flag question

Question text

A male child presented with pain and discharge of one year duration at the leg. He sustained trauma at the site prior to onset. What is the likely diagnosis the patient is having

Select one:

a. Syphilis of the joint

b. Chronic Osteomyelitis

c. Cold abscess

d. Tuberculosis of bone

Question 24

Answer saved

Marked out of 1.00

Remove flag

Question text

A patient requests that you do not talk at all to husband about her medical care. Under which ethical concept does this request fall

Select one:

a. Informed consent

b. Confidentiality

c. Nonmaleficence

d. Advanced directive

Question 25

Answer saved

Marked out of 1.00

Remove flag

Question text

According to WHO classification of female genital cutting in 2008, which of the following describes type II?

Select one:

a. Narrowing of the vaginal orifice with creation of a covering seal by cutting with or without excision of the clitoris.

b. Partial or total removal of the clitoris and/or the prepuce

c. Partial or total removal of the clitoris and the labia minora, with or without excision of the labia majora.

d. Stitching the labia majora after cutting the labia minora

Question 26

Answer saved

Marked out of 1.00

Remove flag

Question text

At the start of a cohort study the exposure is determined with the help of a questionnaire. During the study there is no "loss to follow-up". At the end of the follow-up time the number of cases is known and is divided into exposed and un exposed. The odds ratio (OR) is used as the association measure. Which comment is the most appropriate here

Select one:

a. The OR has no useful interpretation.

b. Researchers should have better calculated the risk ratio

c. The OR approaches the incidence rate ratio.

d. Researchers should have better calculated the incidence rate ratio.

Question 27

Answer saved

Marked out of 1.00

Flag question

Question text

Helen is a 21 yrs old primigravida at 25wks, presented to your clinic because she noticed a small amount of yellowish discharge from her nipples. What is the best advice for her problem?

Select one:

a. An indication that the woman milk is forming

b. A sign of breast cancer

c. Too early in the pregnancy for lactation to begin the woman needs a referral to a specialist

d. Most likely to be colostrum and consider a normal finding this late in her pregnancy

Question 28

Answer saved

Marked out of 1.00

Flag question

Question text

In patients with shock, what is the most frequent cause of death?

Select one:

a. Hemorrhage

b. Minfarction

c. Ventilatory failure

d. Sepsis

Question 29

Answer saved

Marked out of 1.00

Flag question

Question text

Instrumental processing is one of the components of infection prevention. Which one of the following is the correct step in processing MVA

Select one:

a. Decontamination, cleaning, high level disinfection, storage.

b. High level disinfection, decontamination, cleaning, storage

c. High level dis infection, decontamination, cleaning, boiling

d. Cleaning, decontamination, high level disinfection and storage

Question 30

Answer saved

Marked out of 1.00

Flag question

Question text

On routine ANC follow up of a mother in her second trimester pregnancy obstetric ultrasound was done and neural tube defect of the fetus was suspected. Which one of the following investigation least likely supports the diagnosis of neural tube defect

Select one:

a. Elevated maternal serum alpha-fetoprotein level

b. Elevated amniotic fluid acetyl cholinesterase level

c. Elevated amniotic fluid alpha-fetoprotein level

d. Elevated maternal beta HCG level

One of the distinguishing features of Parkinsons disease as opposed to essential tremor is

Select one:

a. Age of onset

b. Presence of rigidity in essential tremor

c. Bilateral nature of the tremor in the initial phases of Parkinsons disease

d. Tremor at rest in Parkinsons disease

Question 32

Answer saved

Marked out of 1.00

Flag question

Question text

One of the following is an important family history that may suggest the possibility of neuropathy in a patient who presents with neurologic symptoms

Select one:

a. History of maternal exercise

b. Consanguineous marriage

c. Degree of love between parents

d. Paternal occupation

Question 33

Answer saved

Marked out of 1.00

Flag question

Question text

One of the following is appropriate measure for first aid of seizures?

Select one:

a. Preventing injuries and falls without restraining the patient

b. Immediately transferring the patient to the general ward

c. Leaving the patient loose so he can breath easily

d. Turning the person to the supine position

Question 34

Answer saved

Marked out of 1.00

Flag question

Question text

Sterilization is a process of destroying all microorganisms including endospores. Which one of the following is the correct way of sterilizing instruments

Select one:

a. Dry heat using an oven with a fan:170 0 C for 1 hour, or 160 0C for 2 hours

b. Steam sterilization using an autoclave: 121 0 C 106 kPa pressure: 20 minutes for unwrapped items, 30 minutes for wrapped items

c. Chemical sterilization using a glutaraldehyde, soak items for 10 hours or in formaldehyde for 24 hours.

d. All are correct way of sterilizing instruments.

Question 35

Answer saved

Marked out of 1.00

Flag question

Question text

The following is a factor that contributes to female susceptibility to vasomotor symptoms

Select one:

a. Higher cardiac baroreflex sensitivity

b. Higher blood volume

c. Higher baseline blood pressure

d. Less muscle mass

Question 36

Answer saved

Marked out of 1.00

Flag question

Question text

The most common cause of stroke in the young is

Select one:

a. Vasculitis

b. Cardio-embolism

c. HIV infection

d. Hypertension

Question 37

Answer saved

Marked out of 1.00

Flag question

Question text

What is the most effective measure that can be instituted to reduce the risk of chronic obstructive pulmonary disease is?

Select one:

a. Weight loss

b. Reduction of respiratory infections by hand washing

c. Smoking cessation

d. Exercise

Question 38

Answer saved

Marked out of 1.00

Flag question

Question text

What is the most important characteristic of diarrhea caused by Vibrio cholera?

Select one:

a. Massive bloody diarrhea

b. Severe abdominal pain

c. Mucoid diarrhea

d. Profound watery diarrhea

Question 39

Answer saved

Marked out of 1.00

Flag question

Question text

When observing the vestibule, you should be able to see which of the following anatomic structures?

Select one:

a. Urethral meatus and paraurethral (Skene) glands

b. Urethral meatus and vaginal orifice

c. Vaginal orifice and vestibular (Bartholin) glands

d. Paraurethral (Skene) and vestibular (Bartholin) glands

Question 40

Answer saved

Marked out of 1.00

Flag question

Question text

While performing CPR intubation for four year-old child, what is the required size of the uncuffed endotracheal tube

Select one:

a. 5

b. 4.5

c. 6

d. 4

Started on	**Sunday, 30 July 2017, 8:52 PM**
State	Finished
Completed on	Sunday, 30 July 2017, 9:45 PM
Time taken	53 mins 19 secs
Marks	29.00/40.00
Grade	**7.25** out of 10.00 (**73**%)

Question 1

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

A 16-year-old student presents to the clinic for a sport physical, she is accompanied by her mother. She is involved in gymnastics and has been since age 5. In addition to training for the gymnastics team, she runs 10 miles per day. She is proud of her athleticism and states that she is still too fat and wants to lose more weight. You note that her BMI is 15. She denies eating excessively or making herself vomits. The patients mother states she barely eats. This patient is at risk for which medical problem?

Select one:

a. Anorexia nervosa

b. Bulimia

c. Not at risk; normal adolescence

d. Obesity

Question 2

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 17 years old boy presented with a low grade fever and pain in the right lower quadrant. On palpation he has tenderness of the left lower quadrant of the abdomen. What is the most likely diagnosis

Select one:

a. Irritable bowel syndrome

b. Acute appendicitis

c. Psychogenic pain

d. Crohn disease

Question 3

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

A 23-year old patient was admitted to surgical ward after sustaining major trauma to the chest and abdomen following road traffic accident of 02 days. After insertion of chest tube, laparotomy was decided and the patient was kept NPO then after. Which of the following is the primary source of energy in this patient?

Select one:

a. Protein (muscle)

b. Fat

c. Ketone bodies

d. Glycogen

Question 4

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

A 25-year-old man was involved in a fight and had sustained machete injury to his scalp a week back. He was brought to the hospital because he developed generalized spasms since 24 hours. Physical examination revealed awake patient with a 4cm-sutured wound over the scalp with palpable depressed parietal bone underneath. He has weakness of his left upper limb. What is the diagnosis

Select one:

a. Epidural hematoma

b. Subdural hematoma

c. Tetanus

d. Increased ICP

Question 5

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 26 year old male presented to the emergency OPD suffering from severe allergic reaction to a bee sting. What is the most appropriate initial therapy

Select one:

a. Albuterol

b. Cimetidine

c. Nor epinephrine

d. Epinephrine

Question 6

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 29 years old female epileptic patient is brought to OPD seizing for more than 10 minutes and she was still having seizure.What is the first line of treatmentof this patient?

Select one:

a. Intravenous pentobarbital

b. Intravenous diazepam

c. Intravenous Paraldehyde

d. Intravenous phenytoin

Question 7

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 29 years old man presented with low grade fever, loss of appetite, weight loss, night sweating and cough productive of sputum of one month duration. Which one is the gold standard diagnostic test for this patients condition

Select one:

a. PPD test

b. Chest X-ray

c. ESR

d. AFB test

Question 8

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 30 year old man presented with Parkinsonian symptoms. The most probable cause is

Select one:

a. Parkinson disease

b. Drug/toxin induced

c. Idiopathic

d. Stroke

Question 9

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 30-year old patient develops high-grade fever on the fifth post laparotomy day. What is the most likely cause of the fever

Select one:

a. Surgical site infection

b. Metabolic crisis

c. Atelectasis

d. Malignant hyperthermia

Question 10

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

A 32-year-old female patient presented 4 hours after she sustained a human bite to her forearm. Physical examination revealed a 4cm linear laceration over her mid right forearm. What is the appropriate management

Select one:

a. Primary closure

b. Delayed primary closure

c. Secondary closure

d. Leave it to heal by secondary intention

Question 11

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 35-year-old otherwise healthy woman went into shock 6 hours after difficult cholecystectomy. What is the most likely cause of the shock

Select one:

a. Hemorrhage

b. Sepsis

c. Bowel perforation

d. Bile leak

Question 12

Correct

Mark 1.00 out of 1.00

Remove flag

Question text

A 38-year-old daily laborer presented with right leg pain & swelling of 2 days duration. He has no history of trauma. Physical examination revealed BP=90/50mmHg, PR= 136/min, RR=28/min, TO= 35.4OC. His right leg is grossly swollen with blisters on the overlying skin, 2cm ulcer over the lateral malleolus discharging dishwater fluid. What is the appropriate definitive management of this patient?

Select one:

a. IV antibiotics & amputation

b. IV antibiotics & radical debridement

c. PO antibiotics & follow up

d. Antibiotic according to culture & sensitivity

Question 13

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 4 -year old female child presented initially with vaginal bleeding but later she also started to have foul smelling vaginal discharge for the last one month, no bleeding from other sites. The most likely cause of the vaginal bleeding is

Select one:

a. Neoplasm of the vagina

b. Urethral prolapse

c. Vulvovaginitis

d. Foreign body in the vagina

Question 14

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

A 40 year old man presented with weakness of the upper and lower extremities of 1 year duration. Physical examination revealed predominantly distal weakness with absent reduced deep tender reflexes. He is most likely suffering from

Select one:

a. Myopathy

b. Upper motor neuron lesion

c. Neuropathy

d. Stroke

Question 15

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 44-year old patient underwent a major surgical procedure, which resulted in significant blood loss. The patient was transferred with stable vital signs to the recovery where he was started on transfusion of cross-matched blood. Few minutes later, the patient started complaining of nausea, vomiting, fever, chills, chest and flank pain. On examination: BP=80/50mmHg, PR=109bpm, RR=35/min, T0=38. 90C. The urine appeared reddish in color. What is the most likely diagnosis?

Select one:

a. Delayed hemolytic transfusion reaction

b. Allergic transfusion reaction

c. Febrile non hemolytic transfusion reaction

d. Acute hemolytic transfusion reaction

Question 16

Correct

Mark 1.00 out of 1.00

Remove flag

Question text

A 45-year-old man with severe non-bilious vomiting for 4 months presented to the OPD with marked dehydration. His urine output was 20ml/h, and the hematocrit is 48%. What is the predominant metabolic abnormality

Select one:

a. Hypochloremic alkalosis

b. Respiratory alkalosis

c. Metabolic acidosis

d. Intrinsic renal disease

Question 17

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 45-year-old man with severe non-bilious vomiting for 4 months presented to the OPD with marked dehydration. His urine output was 20ml/h, and the hematocrit is 48%. Initial treatment for this patient should include which of the following

Select one:

a. Normal saline with IV potassium

b. Antiemetic with NG tube insertion

c. Ringer lactate solution

d. Administration of DNS with IV potassium

Question 18

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

A 5 years old child sustained a fall down accident from a five meters tree. On arrival to emergency unit, his GCS 8/15. While on evaluation which technique do you use to open the airway

Select one:

a. Chin lift

b. Nose up

c. Jaw thrust

d. Neutral Position

Question 19

Incorrect

Mark 0.00 out of 1.00

Remove flag

Question text

A 50 Years old woman came with a breast lamp and FNAC revealed breast cancer. Which one of the following can be considered as significant risk factor for this woman

Select one:

a. A history of using birth control pills

b. A history of endometrial carcinoma

c. A high fiber, low fat diet

d. Late menarche

Question 20

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 55 years old man was diagnosis to have diabetes mellitus 6 months back. He was put on metformin and his blood glucose level is controlled. Which of the following least characterizes the diseases this patient is having?

Select one:

a. Genetic predisposition

b. Increased visceral fat

c. More insidious onset

d. Low basal insulin secretion early in the disease

Question 21

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

Almaz comes to your clinic for advice about breast examination. What is the correct way of instructing about the self-breast examination?

Select one:

a. Recommend that the best time to perform BSE is Four to Seven days after the first day of the menstrual period

b. Tell the woman that if she is pregnant, she should not perform BSE until the baby is born

c. Urge the woman that she needs to do BSE bimonthly unless she has fibrocystic breast tissue

d. Tell the woman that the best time to perform BSE is in the middle of the menstrual cycle

Question 22

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A 6 month old infant was diagnosed with congenital rubella. Which of the following is the least likely presentation for her?

Select one:

a. Low set ears

b. Cataract

c. Microcephaly

d. Congenital heart disease

Question 23

Correct

Mark 1.00 out of 1.00

Flag question

Question text

A male child presented with pain and discharge of one year duration at the leg. He sustained trauma at the site prior to onset. What is the likely diagnosis the patient is having

Select one:

a. Syphilis of the joint

b. Chronic Osteomyelitis

c. Cold abscess

d. Tuberculosis of bone

Question 24

Correct

Mark 1.00 out of 1.00

Remove flag

Question text

A patient requests that you do not talk at all to husband about her medical care. Under which ethical concept does this request fall

Select one:

a. Informed consent

b. Confidentiality

c. Nonmaleficence

d. Advanced directive

Question 25

Correct

Mark 1.00 out of 1.00

Remove flag

Question text

According to WHO classification of female genital cutting in 2008, which of the following describes type II?

Select one:

a. Narrowing of the vaginal orifice with creation of a covering seal by cutting with or without excision of the clitoris.

b. Partial or total removal of the clitoris and/or the prepuce

c. Partial or total removal of the clitoris and the labia minora, with or without excision of the labia majora.

d. Stitching the labia majora after cutting the labia minora

Question 26

Correct

Mark 1.00 out of 1.00

Remove flag

Question text

At the start of a cohort study the exposure is determined with the help of a questionnaire. During the study there is no "loss to follow-up". At the end of the follow-up time the number of cases is known and is divided into exposed and un exposed. The odds ratio (OR) is used as the association measure. Which comment is the most appropriate here

Select one:

a. The OR has no useful interpretation.

b. Researchers should have better calculated the risk ratio

c. The OR approaches the incidence rate ratio.

d. Researchers should have better calculated the incidence rate ratio.

Question 27

Correct

Mark 1.00 out of 1.00

Flag question

Question text

Helen is a 21 yrs old primigravida at 25wks, presented to your clinic because she noticed a small amount of yellowish discharge from her nipples. What is the best advice for her problem?

Select one:

a. An indication that the woman milk is forming

b. A sign of breast cancer

c. Too early in the pregnancy for lactation to begin the woman needs a referral to a specialist

d. Most likely to be colostrum and consider a normal finding this late in her pregnancy

Question 28

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

In patients with shock, what is the most frequent cause of death?

Select one:

a. Hemorrhage

b. Minfarction

c. Ventilatory failure

d. Sepsis

Question 29

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

Instrumental processing is one of the components of infection prevention. Which one of the following is the correct step in processing MVA

Select one:

a. Decontamination, cleaning, high level disinfection, storage.

b. High level disinfection, decontamination, cleaning, storage

c. High level dis infection, decontamination, cleaning, boiling

d. Cleaning, decontamination, high level disinfection and storage

Question 30

Correct

Mark 1.00 out of 1.00

Flag question

Question text

On routine ANC follow up of a mother in her second trimester pregnancy obstetric ultrasound was done and neural tube defect of the fetus was suspected. Which one of the following investigation least likely supports the diagnosis of neural tube defect

Select one:

a. Elevated maternal serum alpha-fetoprotein level

b. Elevated amniotic fluid acetyl cholinesterase level

c. Elevated amniotic fluid alpha-fetoprotein level

d. Elevated maternal beta HCG level

Question 31

Correct

Mark 1.00 out of 1.00

Remove flag

Question text

One of the distinguishing features of Parkinsons disease as opposed to essential tremor is

Select one:

a. Age of onset

b. Presence of rigidity in essential tremor

c. Bilateral nature of the tremor in the initial phases of Parkinsons disease

d. Tremor at rest in Parkinsons disease

Question 32

Correct

Mark 1.00 out of 1.00

Flag question

Question text

One of the following is an important family history that may suggest the possibility of neuropathy in a patient who presents with neurologic symptoms

Select one:

a. History of maternal exercise

b. Consanguineous marriage

c. Degree of love between parents

d. Paternal occupation

Question 33

Correct

Mark 1.00 out of 1.00

Flag question

Question text

One of the following is appropriate measure for first aid of seizures?

Select one:

a. Preventing injuries and falls without restraining the patient

b. Immediately transferring the patient to the general ward

c. Leaving the patient loose so he can breath easily

d. Turning the person to the supine position

Question 34

Correct

Mark 1.00 out of 1.00

Flag question

Question text

Sterilization is a process of destroying all microorganisms including endospores. Which one of the following is the correct way of sterilizing instruments

Select one:

a. Dry heat using an oven with a fan:170 0 C for 1 hour, or 160 0C for 2 hours

b. Steam sterilization using an autoclave: 121 0 C 106 kPa pressure: 20 minutes for unwrapped items, 30 minutes for wrapped items

c. Chemical sterilization using a glutaraldehyde, soak items for 10 hours or in formaldehyde for 24 hours.

d. All are correct way of sterilizing instruments.

Question 35

Incorrect

Mark 0.00 out of 1.00

Flag question

Question text

The following is a factor that contributes to female susceptibility to vasomotor symptoms

Select one:

a. Higher cardiac baroreflex sensitivity

b. Higher blood volume

c. Higher baseline blood pressure

d. Less muscle mass

Question 36

Correct

Mark 1.00 out of 1.00

Flag question

Question text

The most common cause of stroke in the young is

Select one:

a. Vasculitis

b. Cardio-embolism

c. HIV infection

d. Hypertension

Question 37

Correct

Mark 1.00 out of 1.00

Flag question

Question text

What is the most effective measure that can be instituted to reduce the risk of chronic obstructive pulmonary disease is?

Select one:

a. Weight loss

b. Reduction of respiratory infections by hand washing

c. Smoking cessation

d. Exercise

Question 38

Correct

Mark 1.00 out of 1.00

Flag question

Question text

What is the most important characteristic of diarrhea caused by Vibrio cholera?

Select one:

a. Massive bloody diarrhea

b. Severe abdominal pain

c. Mucoid diarrhea

d. Profound watery diarrhea

Question 39

Correct

Mark 1.00 out of 1.00

Flag question

Question text

When observing the vestibule, you should be able to see which of the following anatomic structures?

Select one:

a. Urethral meatus and paraurethral (Skene) glands

b. Urethral meatus and vaginal orifice

c. Vaginal orifice and vestibular (Bartholin) glands

d. Paraurethral (Skene) and vestibular (Bartholin) glands

Question 40

Correct

Mark 1.00 out of 1.00

Flag question

Question text

While performing CPR intubation for four year-old child, what is the required size of the uncuffed endotracheal tube

Select one:

a. 5

b. 4.5

c. 6

d. 4