STERILIZATION AND DISINFECTION

B y uz:
A isha Ibrahim Mohammed

INTRODUCTION

- **Sterilization**

  A physical or chemical process that completely destroys or removes all microbial life, including spores.

- **Disinfection**

  It is killing or removing of harmful microorganisms

- **Disinfectant**

  Products used to kill microorganisms on inanimate objects or surfaces. Disinfectants are not necessarily sporicidal, but may be sporostatic, inhibiting germination or outgrowth

- **Antiseptic**

  A product that destroys or inhibits the growth of microorganisms in or on living tissue.

- **Aseptic**

  Characterized by the absence of pathogenic microbes.

METHODS OF STERILIZATION

1. Physical methods

1/Radiation

- U.V. light
- Ionizing radiation

2 / Filtration

3/ Heat

- Dry
- Moist

2. Chemical Methods



METHODS OF STERILIZATION

1. Physical methods

1/ Radiation:


A-U.V. light- Has limited sterilizing power because of its poor penetration into most materials. Generally used in irradiation of air in certain areas eg. Operating Rooms and T.B. laboratories.

B-Ionizing radiation- *e.g.* Gamma radiation: Source Cobalt^{60} has greater energy than U.V. light, therefore more effective. Used mainly in industrial facilities e.g. sterilization of disposable plastic syringes, gloves, specimens containers and Petri Dishes.

METHODS OF STERILIZATION

- **2/ Filtration**
- May be done under either negative or positive pressure. Best known example is the membrane filter made from cellulose acetate. Generally removes most bacteria but viruses and some small bacteria *e.g.* Chlamydias & Mycoplasmas may pass through. Thus filtration does not technically sterilize items but it is adequate for circumstances under which it is used.





METHODS OF STERILIZATION

- 3/Sterilization by Heat: Most common method
- A-Dry Heat



1/ Bunsen burner-

- Simplest method is exposing the item to be sterilized to the naked flame e.g. Bunsen burner- for sterilizing bacteriological loops, knives, blades.

2/ Hot air oven



- expose items to 160°C for 1 hour. It has electric element in the chamber as source of heat plus a fan to circulate air for even distribution of heat in chamber. Oven without fan is dangerous.

- Used for Metals, Glassware, Ointment, Oils, Waxes, Powders *i.e.* items that are lacking water

- B-Moist Heat:Uses hot water. Moist heat kills microorganisms by denaturing proteins.
- types
- **1: Boiling**
- **2: Tyndallization**
- **3: Pasteurization**
- **4: Autoclaving**



METHODS OF STERILIZATION

1: Boiling – quite common especially in domestic circumstances.

2: Tyndallization named after John Tyndall

- Lengthy process designed to reduce the level of activity of sporulating bacteria that are left by a simple boiling water method.

CON…
METHODS OF STERILIZATION**MOIST HEAT:**

Tyndallization

- The process involves boiling for a period (typically 20 minutes) at atmospheric pressure, cooling, incubating for a day, boiling, cooling, incubating for a day, boiling, cooling, incubating for a day, and finally boiling again.

- The three incubation periods are to allow heat-resistant spores surviving the previous boiling period to germinate to form the heat-sensitive vegetative (growing) stage, which can be killed by the next boiling step.

- The procedure only works for media that can support bacterial growth - it will not sterilize plain water.

CON…
METHODS OF STERILIZATION**MOIST HEAT:**

3: Pasteurization

- It aims to reduce the number of viable pathogens in liquids so they are unlikely to cause disease
- It uses heat at temperatures sufficient to inactivate harmful organism in milk. Does not achieve sterilization.
- **Temperature may be**
- 138°C for a fraction of a second (flash method),
- 71.7°C for 15-20 seconds
- or 62°C for 30 minutes.


CON…
METHODS OF STERILIZATION**MOIST HEAT:**

4: Autoclaving –

Standard sterilization method in hospitals.

- The Autoclave works under the same principle as the pressure cooker where water boils at increased atmospheric pressure *i.e.* because of increased pressure the boiling point of water is >100°C.
- The autoclave is a tough double walled chamber in which air is replaced by pure saturated steam under pressure.

- The air in the chamber is evacuated andfilled with saturated steam. The chamber is closed tightly the steam keeps on filling into it and the pressure gradually increases.
- The items to be sterilized get completely surrounded by saturated steam (moist heat) which on contact with the surface of material to be sterilized condenses to release its latent heat of condensation which adds to already raised temperature of steam so that eventually all the microorganisms in what ever form –are killed.
- The usual temperature achieved is 121 °C at a pressure of 15 pps.i. at exposure time of only 15-20 mins. By increasing the temperature, the time for sterilizing is further reduced.


steam exhaust pipe
steam from jacket into chamber
inner chamber
door
steam in
culture
medium
jacket
to drain
steam supply



METHODS OF STERILIZATION

Advantages of Autoclave

- Temperature is > 100°C therefore spores are killed.
- Condensation of steam generates extra heat (latent heat of condensation).
- The condensation also allows the steam to penetrate rapidly into porous materials.
- Note: that autoclavable items must be steam permeable. Can not be used for items that are lacking water.


MONITORING OF AUTOCLAVES

- A- Physical- use of thermocouple to measure accurately the temperature.




CON…

- B- Chemical- it consists of heat sensitive chemical that changes color at the right temperature and exposure time.
  - Autoclave tape
  - Browne’s tube.



CON…

- C- Biological – where a spore-bearing organism is added during the sterilization process and then cultured later to ensure that it has been killed.



METHODS OF STERILIZATION

2/ Chemical Methods

- Useful for heat sensitive materials e.g. plastics and lensed instruments endoscopes).eg:-
- **(1) Ethylene Oxide Chamber:**
  - Ethylene oxide alkylates DNA molecules and thereby inactivates microorganisms.
  - Ethylene oxide may cause explosion if used pure so it is mixed with an inert gas e.g. Neon, Freon at a ratio of 10:90
  - It requires high humidity and is used at relative humidity 50-60% Temperature : 55-60°C and exposure period 4-6 hours.
- **(2) Activated alkaline Glutaraldehyde 2%:**
  - Immerse item in solution for about 20 minutes if organism is TB. In case of spores, the immersion period is extended to 2-3 hours.

DISINFECTANTS

Types of Disinfectants

Phenol and phenolics

- Phenol (carbolic acid) is seldom used today. Derivatives of the phenol molecule, however, are widely used.
- Phenolics injure plasma membrane, inactivate enzymes, or denature proteins.
- They are stable, persistent, and are not sensitive to organic matter.



DISINFECTANTS

Hexachlorophene

- It is main ingredient of a prescription lotion, pHisoHex, used in nurseries and for surgical and hospital microbial control procedures to control gram positive skin bacteria such as staphylococci and streptococci.
- Excessive use can cause neurological damage.

Triclosan

- It is a widely used found in many household products. It has broad spectrum of activity, especially against gram positive bacteria. It is also effective against gram negative bacteria and fungi.

DISINFECTANTS

Biguanides

- Chlorhexidine, a member of the biguanide group, is not a phenol, but its structure and applications resemble hexachlorophene. It is frequently used for surgical skin preparation and surgical hand scrubs.

Halogens

- **Iodine** is effective against all kinds of bacteria, many endospores, fungi, and some viruses. Its mechanism of activity may be its combination with the amino acid tyrosine in enzyme and cellular proteins.



- **Chlorine**
- is used as a gas or in combination with other chemicals. Chlorine gas is used for disinfecting municipal water supplies, swimming pools, and sewage.
- Sodium hypochlorite – ordinary household bleach- is good disinfectant.
- Chloramines consist of chlorine and ammonia. They are more stable than most chlorine. The U.S. military uses tablets for field disinfection of water.

DISINFECTANTS

Alcohols

- Both ethanol and isopropanol (rubbing alcohol) are widely used, normally at a concentration of about 70%.
- They are bactericidal and fungicidal but are not effective against endospores or non-enveloped viruses.

- Alcohols enhance the effectiveness of other chemical agents.



DISINFECTANTS

Heavy metals and their compounds

- Tiny amount of heavy metals (e.g. silver and copper) are effective antimicrobials. A silver coin on an inoculated nutrient medium will inhibit growth for some distance.

- Silver-sulfadiazine is used in wound dressings. Available as topical cream for use on burns.



Mercuric chloride is highly bactericidal, but is toxic and corrosive and is inactivated by organic matter. Organic mercury compounds such as Mercurochrome are less irritating and less toxic than inorganic mercury.

- Copper sulfate is often used to destroy green algae in reservoirs or other water.

- Zinc chloride is used in mouthwashes, and zinc oxide is used in paints as antifungal

