

GULELE SUB-CITY EDUCATION OFFICE, ADDIS ABEBA

GRADE 12 MATHEMATICS MODEL EXAMINATION

SENE 2012/MAY 2020

NUMBER OF QUESTIONS: 65

TIME ALLOWED:- 3 HOURS

GENERAL DIRECTIONS

THIS BOOKLET CONTAINS **MATHEMATICS (FOR SOCIAL SCIENCE STUDENTS)** EXAMINATION. IN THIS EXAMINATION, THERE ARE A TOTAL OF 65 **MULTIPLE CHOICE QUESTIONS**. CAREFULLY SELECT THE BEST ANSWER AND **BLACKEN** ONLY THE LETTER OF YOUR CHOICE ON THE SEPARATE ANSWER SHEET PROVIDED. FOLLOW THE INSTRUCTIONS ON THE ANSWER SHEET AND THE EXAMINATION PAPER CAREFULLY. USE ONLY **PENCIL** TO MARK YOUR ANSWERS. YOUR ANSWER MARK SHOULD BE **HEAVY** AND **DARK**, COVERING THE ANSWER SPACE COMPLETELY. PLEASE ERASE ALL UNNECESSARY MARKS COMPLETELY FROM YOUR ANSWER SHEET.

YOU ARE ALLOWED TO WORK ON THE EXAM FOR **3 HOURS**. WHEN TIME IS CALLED, YOU MUST IMMEDIATELY STOP WORKING, PUT YOUR PENCIL DOWN, AND WAIT FOR FURTHER INSTRUCTIONS.

ANY FORM OF CHEATING OR AN ATTEMPT TO CHEAT IN THE EXAMINATION WILL RESULT IN AN AUTOMATIC DISMISSAL FROM THE EXAMINATION HALL AND CANCELLATION OF YOUR SCORE (S).

PLEASE MAKE SURE THAT YOU HAVE WRITTEN ALL THE REQUIRED INFORMATION ON THE ANSWER SHEET BEFORE YOU START TO WORK ON THE EXAMINATION.

DO NOT TURN THIS PAGE UNTIL YOU ARE TOLD TO DO SO.

DIRECTION: Each of the following questions is followed by four possible alternatives. Read each question carefully and **BLACKEN** the letter of your choice on the answer sheet provided.

- Which one of the following is a one to one function?
 - $f = \{(1, 6), (2, 7), (3, 6), (4, 8)\}$
 - $k : \mathbb{R} \rightarrow \mathbb{R}$ given by $h(x) = |x - 3|$
 - $k : (0, \infty) \rightarrow \mathbb{R}$ given by $h(x) = \log x$
 - $g = \{(x, y) : x \text{ is a student and } y \text{ is his / her rank}\}$
- The domain of the function f defined by $f(x) = \frac{1}{\sqrt{x - |x|}}$ is:
 - \mathbb{R}
 - $(-\infty, 0)$
 - $(0, \infty)$
 - f is not defined for any $x \in \mathbb{R}$
- If $f(x) = \ln\left(\frac{x}{x-1} + 2\right)$, for $x > 1$, then which one of the following is the inverse of f ?
 - $g(x) = \frac{e^x - 2}{e^x - 3}$
 - $g(x) = \frac{e^x}{e^x + 1} - 2$
 - $g(x) = \frac{e^x - 2}{e^x + 1}$
 - $g(x) = e^{\frac{x}{x-1}} - 2$
- Given the functions $f(x) = x^2 - 2x + 8$ and $g(x) = 0.5x - 3$, then $(f \circ g^{-1})(2)$ is equal to:
 - 88
 - 100
 - 20
 - 68
- The simplified form of $\frac{x^{n+2} + yx^{n+1} + x^n y^2}{x^{3-n} - x^{-n} y^3} \div \frac{x^{2n}}{3}$, for $x \neq y$ and $x \neq 0$
 - $\frac{x+y}{3}$
 - $\frac{x-y}{3}$
 - $\frac{3}{x-y}$
 - $\frac{3}{x-y}$
- The solution set of the equation $\frac{x^2}{x+3} + \frac{3x}{x^2+5x+6} = \frac{2x}{x+3}$ is:
 - $\{0, 1\}$
 - $\{-1, 1\}$
 - $\{-1, 0, 1\}$
 - $\{-1, 0\}$
- At what value(s) of x does the graph of $f(x) = \frac{x^4 + 1}{x^4 + x^2}$ crosses its horizontal asymptote?
 - $x = 0$ and $x = 1$
 - $x = -1$ and $x = 1$
 - $x = 1$
 - $x = -1$

8. Consider the sequence $\{a_n\}_{n \geq 1}$, where the first term is $a_1 = 2$ and $a_{n+1} = a_n + 4$ for all $n \geq 1$.

Then the sum $a_3 + a_4 + a_5 + \dots + a_{35}$ is equal to:

- A. 2460 B. 2458 C. 2442 D. 2450

9. What is the fourth term of a geometric sequence whose third term is 1 and whose eighth term is $\frac{1}{32}$?

- A. 4 B. $\frac{1}{2}$ C. $\frac{1}{8}$ D. $\frac{1}{16}$

10. What is the sum of the series $\sum_{n=1}^{\infty} (-4) \left(\frac{5}{3}\right)^{1-n}$?

- A. $-\frac{5}{2}$ B. $-\frac{12}{5}$ C. -4 D. -10

11. Let $f(x) = \begin{cases} 3k - x, & \text{if } x < 1 \\ b + 1, & \text{if } x = 1 \\ kx, & \text{if } x > 1 \end{cases}$. What is the value of b if f is continuous at $x = 1$?

- A. $\frac{1}{2}$ B. $-\frac{1}{2}$ C. -1 D. 3

12. The left hand side limit, $\lim_{x \rightarrow 2^+} \frac{x|x-2|}{x-2}$ equals to:

- A. 1 B. 2 C. -2 D. Does not exist

13. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that $f(1) = \frac{3}{2}$ and $f\left(\frac{1}{\sqrt{2}}\right) = 3$. Then the value of

$\lim_{x \rightarrow 0} f\left(\frac{\sqrt{1+x} - \sqrt{1-x}}{x}\right)$ is equal to:

- A. $\frac{3}{2}$ B. 3 C. $\frac{9}{2}$ D. Can not be determined

14. Which of the following is not true?

- A. $\lim_{n \rightarrow \infty} \left(\frac{2n-3}{3n+7}\right)^4 = \frac{16}{81}$ C. $\lim_{n \rightarrow \infty} \left(\frac{1+2 \times 10^n}{5+3 \times 10^n}\right) = \frac{2}{3}$
- B. $\lim_{n \rightarrow \infty} \left(\frac{\sqrt{1+9n^2}}{3n+5}\right) = 1$ D. $\lim_{n \rightarrow \infty} \left(\frac{\sqrt{16n^2+9}}{3n-1}\right)^4 = \frac{16}{3}$

15. Evaluate $\lim_{n \rightarrow \infty} \frac{(-1)^n n^3 + 1}{2n^3 + 3}$.
- A. -1 B. ∞ C. 1 D. does not exist
16. If in a hyperbola, the distance between the foci is 10 and the transverse axis has length 8, then the length of its latus rectum is
- A. 9 B. $\frac{9}{2}$ C. $\frac{32}{2}$ D. $\frac{64}{3}$
17. The equation of the directrix of the parabola $y^2 + x + 2y + 1 = 0$ is:
- A. $x = \frac{-1}{4}$ B. $y = \frac{-1}{4}$ C. $x = \frac{1}{4}$ D. $y = \frac{1}{4}$
18. What is the equation of an ellipse with centre at (3,2), vertices at (5,2) and (3,5)
- A. $9(x-3)^2 + 4(y-2)^2 = 36$ C. $4(x-3)^2 + (y-2)^2 = 1$
 B. $4(x-3)^2 + 9(y-2)^2 = 36$ D. $9(x-3)^2 + 4(y-2)^2 = 1$
19. A parabolic arch has a height of 20m and width of 36m at the base. If the vertex of the parabola is at the top of the arch, at what height above the base is it 18m wide?
- A. 18m B. 15m C. 20m D. 24m
20. One of the following is not true about the ellipse whose equation is given by $2x^2 + 8y^2 = 32$
- A. the length of the minor axis is 4 C. the foci are $(0, -2\sqrt{3})$ and $(0, 2\sqrt{3})$
 B. the major axis is parallel to the x – axis D. $(-4, 0)$ and $(0, 2)$ are vertices
21. Which one of the following is equation of a circle whose center is on y-axis and radius is 3?
- A. $x^2 + y^2 + 6y = 0$ C. $(x-2)^2 + y^2 = 9$
 B. $x^2 + (y-2)^2 = 3$ D. $x^2 - 2x + y^2 = 8$
22. Which of the following is equivalent to $\neg[(p \Rightarrow \neg q) \vee \neg r]$?
- A. $(\neg p \vee q) \wedge r$ C. $\neg p \wedge q \wedge r$
 B. $p \wedge q \wedge r$ D. $(q \Rightarrow \neg p) \wedge r$
23. Which of the following is **not** true statement on the set of real numbers?
- A. $(\forall x)(\exists y) (y = x + 2)$ C. $(\exists x)(\exists y) (x^2 + y^2 = 5)$
 B. $(\exists y)(\forall x) (y + x = 3)$ D. $(\exists x)(\exists y) (y < x)$
24. If each of the compound propositions $P \wedge \neg Q$, $R \Rightarrow Q$, $\neg R \Leftrightarrow S$ is true, then which one of the following is true?
- A. Q B. S C. $P \wedge \neg S$ D. $S \Rightarrow R$

25. Suppose the following are the premises of an argument.

“He was lazy or he did not like the classroom.

He could not pass the exam, if he was lazy.

He passed the exam.”

Which of the following can be a conclusion that makes the argument valid?

A. He did like the classroom

B. If he was not lazy, he did like the classroom

C. He did not like the classroom.

D. He was not lazy and he did like the classroom.

26. If the mean of the frequency distribution given below is 2, then what is the value of x?

V	-7	-2	3	9	13
F	10	6	2	5	x

A. 3

B. 6

C. 4

D. 7

27. Below are the marks of grade 12 students in mathematics out of 100. What is the variance for the frequency distribution given below

Marks	20-40	40-60	60-80	80-100
Number of students	4	10	4	2

A. 107.2

B. 4

C. 304

D. 17.5

28. Which one of the following is the constant term of the expansions of $\left(x^2 + \frac{2}{x^3}\right)^5$?

A. 40

C. 80

B. 60

D. 20

29. In a random arrangement of the word “SOCIETY” what is the total number of words in which all the three vowels come together?

A. 120

C. 420

B. 720

D. 840

30. Items produced by a certain company are subject to kinds of defects D_1 and D_2 . Out of the total production, if 5% have defect D_1 , 10% have defect D_2 and 2% have both defects, then what is the probability that an item to have defect D_2 , given that it has defect D_1 .

A. 0.4

B. 0.2

C. 0.1

D. 0.5

31. A three digit library identification card is to be printed from the numbers 0, 1, 2, 3, 4 and 5 in such a way that the first is non-zero and **no** number is to be repeated, how many such cards can be printed?

- A. 100 B. 120 C. 180 D. 150

32. Consider the experiment of rolling a die whose sample space is $\{1, 2, 3, 4, 5, 6\}$. If two dice are rolled simultaneously, what is the probability that a prime number turns up on one of the dice and a composite number on the other?

- A. $\frac{1}{3}$ B. $\frac{2}{3}$ C. $\frac{1}{6}$ D. $\frac{5}{6}$

33. Different codes each consisting of five characters are to be generated in each code the first two characters are either A or B and the remaining three characters are any of the digits 0, 1, 2, ..., 9. how many distinct codes can be generated so?

- A. 4000 B. 3600 C. 3000 D. 2400

34. Which of the following matrix is **not** in reduced row echelon form?

A. $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

C. $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

B. $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$

D. $\begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

35. Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 \\ 1 & c \end{bmatrix}$. For what value of c is $AB = \begin{bmatrix} 3 & 8 \\ 7 & 18 \end{bmatrix}$?

- A. 3 B. 2 C. 4 D. 1

36. Which of the following is the solution set of the system of equation $\begin{cases} x + y + 2z = 1 \\ x + 2y + z = 1 \\ 3x + 4y + 5z = 7 \end{cases}$?

- A. $\{(1, 0, 0)\}$ C. $\{(-3k + 1, k, k) | k \in \mathcal{R}\}$
 B. $\{(k + 1, -k, 0) | k \in \mathcal{R}\}$ D. \emptyset

37. IF $A = \begin{bmatrix} 3 & 3 & 5 \\ 0 & -1 & 2 \\ 4 & 2 & 1 \end{bmatrix}$, then the cofactors of a_{23} and a_{31} respectively are:

- A. 11 and 6
B. -6 and 11
C. 6 and -11
D. 6 and 11

38. One of the following is false about matrices?

- A. If A is a square matrix with order n , then $\frac{(A+A^T)}{3}$ is symmetric matrix.
B. If B is a square matrix with order m , then the matrix $D = 2B - 3B^T$ is skew symmetric.
C. For any two matrices $A = (a_{ij})_{mn}$ and $B = (b_{ij})_{pq}$, BA exists if and only if $m = q$.
D. For any two none zero matrices A and B their product AB may be zero matrix.

39. Let $A = \begin{bmatrix} 2 & 3 & 5 \\ 4 & 5 & 7 \\ 8 & -3 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} -4 & 3 & -5 \\ 2 & 1 & 7 \\ 6 & 5 & 0 \end{bmatrix}$. If $C = AB + A^T$, what is the (2, 3) entry of C ?

- A. 18
B. 42
C. 46
D. 12

40. If z is a complex number such $\frac{-5z}{4+3i} = 3i(\sqrt{3}-i)$, then $|z|$ is equal to:

- A. 6
B. 9
C. 15
D. $\sqrt{15}$

41. The polar form of $2 - 2\sqrt{3}i$

- A. $4(\cos \frac{4}{3}\pi + i\sin \frac{4}{3}\pi)$
B. $4(\cos \frac{\pi}{3} + i\sin \frac{\pi}{3})$
C. $4(\cos \frac{5}{6}\pi + i\sin \frac{5}{6}\pi)$
D. $4\left(\cos\left(\frac{\pi}{3}\right) - i\sin\left(\frac{\pi}{3}\right)\right)$

42. If $z = 2 + 3i$, complex number, then $|z|^2 - z^2 =$ _____

- A. 0
B. $18 - 12i$
C. $8 - 12i$
D. $12i$

43. The locus represented by $|z - 1| = |z + i|$ where $z = x + iy$ $x, y \in \mathbb{R}$ is

- A. a circle of radius 1
B. an ellipse with foci at (1, 0) and (-1, 0)
C. a straight line through the origin
D. a circle with centre at $\left(\frac{1}{2}, \frac{1}{2}\right)$

44. If $y = x^y$, then $\frac{dy}{dx}$ equals:

- A. $\frac{x^2}{y(1 - y \ln x)}$
B. $\frac{y^2}{x(1 - x \ln y)}$
C. $\frac{y^2}{x(1 - x \ln x)}$
D. $\frac{y^2}{x(1 - y \ln x)}$

45. Which of the following is the equation of the tangent line to the graph of $f(x) = \frac{3x}{x-1}$ at

$(2, f(2))$?

A. $3y + x = 20$

B. $3y - x = 16$

C. $y - 3x = 0$

D.

$y + 3x = 12$

46. Let $f(x) = \ln(x^2 - 3x)$. Then, $f''(-1)$ is equal to :

A. $\frac{-17}{16}$

B. $\frac{-7}{16}$

C. $\frac{-5}{16}$

D. $\frac{-19}{16}$

47. If $f(2) = -3$, $f'(2) = 4$, $g(1) = -5$, $g'(1) = 1$ and $F(x) = f(2x+2) \cdot g(1-x^2)$, then what is the value of $F'(0)$?

A. 19

B. 0

C. -20

D. -40

48. Let $f(x) = x^5 - 20x$ on $[-2, 2]$. Which of the following is the local maximum value of f ?

A. $f(-\sqrt{2})$

B. $f(\sqrt{2})$

C. $f(2)$

D. $f(-1)$

49. A point moves around a circle $x^2 + y^2 = 9$. When the point is at $(-\sqrt{3}, \sqrt{6})$ its x-coordinate is increasing at the rate of 20 unit per second. How fast is its y-coordinate changing in unit per second at that instant?

A. $5\sqrt{2}$

B. $10\sqrt{2}$

C. $10\sqrt{3}$

D. $20\sqrt{3}$

50. Which of the following is false about the graph of $f(x) = \frac{x}{x^2 + 1}$

A. It is strictly increasing on $[-1, 1]$

B. one of the inflection point is $(0, 0)$

C. It is concave down ward on $(-\infty, -\sqrt{3}) \cup (0, \sqrt{3})$

D. Its relative maximum value is $\frac{-1}{2}$

51. Two concentric circles are such that the radius of one is always three times that of the other. If the radius of the smaller is increasing at the rate of 2m/sec. find the rate at which the area between the circles is increasing when the radius of the larger is 3m.

A. $36\pi m^2 / \text{sec}$

B. $24\pi m^2 / \text{sec}$

C. $32\pi m^2 / \text{sec}$

D. $18\pi m^2 / \text{sec}$

52. For what values of a and b $h(x) = a + bx - x^2$ have maximum value of 5 at $x = 3$

- A. -4, 6 B. 4, 6 C. 4, -6 D. -2, 6

53. Which one of the following is equal to $\int \frac{x + \ln(x+1)}{(x+1)^2} dx$?

- A. $\ln(x+1) + \frac{x}{x+1} + c$ C. $(x+1)^2 - \frac{1}{x+1} + c$
 B. $(x+1)^2 + \frac{1}{x+1} + c$ D. $\frac{x \ln(x+1)}{x+1} + c$

54. If $F(x)$ is an antiderivative of $f(x) = 1 - \frac{2}{x^2}$ and $F(1) = 0$, then $F(2)$ is equal to:

- A. 0 B. $\frac{1}{2}$ C. $-\frac{1}{2}$ D. 3

55. If $f(x) = \begin{cases} x^2 + 2, & x \geq 2 \\ 5x - 4, & x < 2 \end{cases}$, then $\int_{-1}^3 f(x) dx$ is equal to:

- A. $\frac{35}{2}$ B. $\frac{23}{6}$ C. $\frac{6}{23}$ D. $\frac{2}{35}$

56. Three partners P_1 , P_2 , and P_3 , have setup a trading center by contributing Birr 4000, 5000 and 2000, respectively. If the profit of birr 3432 obtained from the trading center is to be divided between the partners in proportion to their contributions, how much birr will P_1 get from the profit?

- A. 1342.75 B. 1248 C. 1224.5 D. 1144

57. What are the values of x and y that minimize $z = -2x + y$

$$\begin{aligned} \text{subject to } & x + y \leq 7 \\ & x - y \leq 3 \quad ? \\ & y \leq 4 \\ & x \geq 0, y \geq 0 \end{aligned}$$

- A. $x = 0, y = 0$ C. $x = 7, y = 4$
 B. $x = 7, y = 0$ D. $x = 5, y = 2$

58. An item is sold for birr 192 after 20% discount from its regular price. How much was its regular price?

- A. Birr 212 B. Birr 232 C. Birr 238 D. Birr 240

59. A grinder purchased 3 years ago for birr 75,000 is now appraised at birr 54,000. If the rate of depreciation is determined by the fixed installment method, after how many years from now will the grinder be depreciated to Birr 12,000?
- A. 9 B. 8 C. 6 D. 5
60. Suppose z is directly proportional to x and inversely proportional to y . If $z = 0.3$ when the ratio of y to x is $2:3$, how much is z when $x = 5$ and $y = 4$?
- A. $\frac{5}{4}$ B. $\frac{4}{5}$ C. $\frac{3}{8}$ D. $\frac{1}{4}$
61. Which one of the following is a type of investment issued by government or corporate usually known as fixed income security?
- A. Saving account C. Stock
B. Certificate of deposit D. Bond
62. A publishing company has purchased a new machine at a cost of birr 12,000. the machine has a useful life of 8 years. If the salvage value at the end of 8 years is Birr 2,000, what is its book value at the end of 5 years?
- A. Birr 6820 B. Birr 5,685 C. Birr 5,750 D. Birr 3,925
63. What the maximum value of $z = 3x - 2y$ on S , where
 $S = \{(x, y) \mid x + y \geq 2, 2x - y, -x + 2x \leq 4\}$?
- A. 12 B. 6 C. 4 D. 8
64. A saving and a loan wants to offer a certificate of deposit with a monthly compound interest that has an effective rate of 4%. What annual nominal rate compounded monthly should they use?
- [you may use the information: $(1.04)^{\frac{1}{12}} = 1.003273$ $(1.04)^{12} = 1.601032$
- A. 3.9% B. 4.8% C. 5.2% D. 7.2%
65. The population of Addis Ababa increased by 5% each year. At this rate, after how many years will the current population be doubled?
- A. 25 B. 35 C. 20 D. 40